

Searching for Holes in the Matrix Universe

Lucas Kerbs

Spring 2022

2022-06-03

Searching for Holes in the Matrix Universe

Searching for Holes in the Matrix Universe

Lucas Kerbs

Spring 2022

- Eventual goal: lift the tools of algebraic topology to spaces of matrices
- If we only consider 2×2 matrices we can use classical theory
- The moment we want more than one size, things the classical theory breaks down
- Today we will develop some *fairly heftly* tools to do just that
- Along the way, hopefully I can convince you that this is an interesting question.
- To do so, we need to go back to our mathematical roots

Part I: Objects and Maps

A Naive Attempt

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

Part I: Objects and Maps
A Naive Attempt

- That’s right—objects and maps.
- Our Naive attempt involves that looking at lifting functions on \mathbb{R} or \mathbb{C} to accept matrices as their input.
- An operator theorist would call this a “functional calculus”

Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ($A = A^*$) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix

Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.
 A is diagonalizable as $A = U\Lambda U^*$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ($A = A^*$) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.

Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.
 A is diagonalizable as $A = U \Lambda U^*$

$$f(A) = a_n A^n + \cdots + a_1 A + a_0 I_k$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.
 A is diagonalizable as $A = U \Lambda U^*$

$$f(A) = a_n A^n + \cdots + a_1 A + a_0 I_k$$

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ($A = A^*$) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.
- Watch what happens when we plug this into our polynomial — need to be careful with the constant term.

Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.

A is diagonalizable as $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \end{aligned}$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.
 A is diagonalizable as $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \end{aligned}$$

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ($A = A^*$) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.
- Watch what happens when we plug this into our polynomial — need to be careful with the constant term.

Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.

A is diagonalizable as $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \end{aligned}$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.
 A is diagonalizable as $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \end{aligned}$$

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ($A = A^*$) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.
- Watch what happens when we plug this into our polynomial — need to be careful with the constant term.

Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.

A is diagonalizable as $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \end{aligned}$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.
 A is diagonalizable as $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \end{aligned}$$

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ($A = A^*$) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.
- Watch what happens when we plug this into our polynomial — need to be careful with the constant term.

Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.

A is diagonalizable as $A = U \Lambda U^*$

$$\begin{aligned}
 f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\
 &= a_n (U \Lambda U^*)^n + \cdots + a_1 U \Lambda U^* + a_0 I_k \\
 &= a_n U \Lambda^n U^* + \cdots + a_1 U \Lambda U^* + a_0 I_k \\
 &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \\
 &= U (f(\Lambda)) U^*
 \end{aligned}$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.
 A is diagonalizable as $A = U \Lambda U^*$

$$\begin{aligned}
 f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\
 &= a_n (U \Lambda U^*)^n + \cdots + a_1 U \Lambda U^* + a_0 I_k \\
 &= a_n U \Lambda^n U^* + \cdots + a_1 U \Lambda U^* + a_0 I_k \\
 &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \\
 &= U (f(\Lambda)) U^*
 \end{aligned}$$

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ($A = A^*$) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.
- Watch what happens when we plug this into our polynomial — need to be careful with the constant term.

Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.

A is diagonalizable as $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \\ &= U (f(\Lambda)) U^* \end{aligned}$$

$$f \left(\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} \right)$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.
 A is diagonalizable as $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \\ &= U (f(\Lambda)) U^* \end{aligned}$$

$$f \left(\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} \right)$$

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ($A = A^*$) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.
- Watch what happens when we plug this into our polynomial — need to be careful with the constant term.

Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.

A is diagonalizable as $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \\ &= U (f(\Lambda)) U^* \end{aligned}$$

$$f\left(\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}\right) = \begin{bmatrix} f(\lambda_1) & & \\ & \ddots & \\ & & f(\lambda_n) \end{bmatrix}$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.
 A is diagonalizable as $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \\ &= U (f(\Lambda)) U^* \end{aligned}$$

$$f\left(\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}\right) = \begin{bmatrix} f(\lambda_1) & & \\ & \ddots & \\ & & f(\lambda_n) \end{bmatrix}$$

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ($A = A^*$) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.
- Watch what happens when we plug this into our polynomial — need to be careful with the constant term.
- This application along the diagonal is precisely the behavior we want to emulate in the functional calculus.

Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.

A is diagonalizable as $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \\ &= U (f(\Lambda)) U^* \end{aligned}$$

$$f\left(\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}\right) = \begin{bmatrix} f(\lambda_1) & & \\ & \ddots & \\ & & f(\lambda_n) \end{bmatrix}$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

Let $f \in \mathbb{R}[x]$ and $A \in M_k(\mathbb{C})$ be self adjoint.
 A is diagonalizable as $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \\ &= U (f(\Lambda)) U^* \end{aligned}$$

$$f\left(\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}\right) = \begin{bmatrix} f(\lambda_1) & & \\ & \ddots & \\ & & f(\lambda_n) \end{bmatrix}$$

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ($A = A^*$) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.
- Watch what happens when we plug this into our polynomial — need to be careful with the constant term.
- This application along the diagonal is precisely the behavior we want to emulate in the functional calculus.

Let \mathbb{H}_n be the set of $n \times n$ self adjoint matrices, and define

$$\mathbb{H} = \bigcup_{n \in \mathbb{N}} \mathbb{H}_n, \quad \mathcal{M} = \bigcup_{n \in \mathbb{N}} M_n(\mathbb{C})$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

Let \mathbb{H}_n be the set of $n \times n$ self adjoint matrices, and define

$$\mathbb{H} = \bigcup_{n \in \mathbb{N}} \mathbb{H}_n, \quad \mathcal{M} = \bigcup_{n \in \mathbb{N}} M_n(\mathbb{C})$$

- With the polynomial case in mind, we can extend a general function. First, a piece of notation
- Lets grab a function on the real line and the self adjoint matrices with their spectrum in that domain
- Then we can lift g by emulating the behavior of polynomials.
- unwrap a self adjoint matrix, apply g to the diagonal, then wrap it back up
- Something to notice about this functional calculus—it treats direct sums *very* well
- This is all well and good, but can we do anything with these new functions?

Let \mathbb{H}_n be the set of $n \times n$ self adjoint matrices, and define

$$\mathbb{H} = \bigcup_{n \in \mathbb{N}} \mathbb{H}_n, \quad \mathcal{M} = \bigcup_{n \in \mathbb{N}} M_n(\mathbb{C})$$

Definition:

Let $g : [a, b] \rightarrow \mathbb{C}$ and $D \subset \mathbb{H}$ denote the set of self adjoint matrices with their spectrum in $[a, b]$.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

Let \mathbb{H}_n be the set of $n \times n$ self adjoint matrices, and define

$$\mathbb{H} = \bigcup_{n \in \mathbb{N}} \mathbb{H}_n, \quad \mathcal{M} = \bigcup_{n \in \mathbb{N}} M_n(\mathbb{C})$$

Definition:

Let $g : [a, b] \rightarrow \mathbb{C}$ and $D \subset \mathbb{H}$ denote the set of self adjoint matrices with their spectrum in $[a, b]$.

- With the polynomial case in mind, we can extend a general function. First, a piece of notation
- Lets grab a function on the real line and the self adjoint matrices with their spectrum in that domain
- Then we can lift g by emulating the behavior of polynomials.
- unwrap a self adjoint matrix, apply g to the diagonal, then wrap it back up
- Something to notice about this functional calculus—it treats direct sums *very* well
- This is all well and good, but can we do anything with these new functions?

Let \mathbb{H}_n be the set of $n \times n$ self adjoint matrices, and define

$$\mathbb{H} = \bigcup_{n \in \mathbb{N}} \mathbb{H}_n, \quad \mathcal{M} = \bigcup_{n \in \mathbb{N}} M_n(\mathbb{C})$$

Definition:

Let $g : [a, b] \rightarrow \mathbb{C}$ and $D \subset \mathbb{H}$ denote the set of self adjoint matrices with their spectrum in $[a, b]$. Then

$$g : D \longrightarrow \mathcal{M}$$

$$X = U \Lambda U^* \longmapsto U \begin{bmatrix} g(\lambda_1) & & \\ & \ddots & \\ & & g(\lambda_n) \end{bmatrix} U^*.$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

- With the polynomial case in mind, we can extend a general function. First, a piece of notation
- Lets grab a function on the real line and the self adjoint matrices with their spectrum in that domain
- Then we can lift g by emulating the behavior of polynomials.
- unwrap a self adjoint matrix, apply g to the diagonal, then wrap it back up
- Something to notice about this functional calculus—it treats direct sums *very* well
- This is all well and good, but can we do anything with these new functions?

Let \mathbb{H}_n be the set of $n \times n$ self adjoint matrices, and define

$$\mathbb{H} = \bigcup_{n \in \mathbb{N}} \mathbb{H}_n, \quad \mathcal{M} = \bigcup_{n \in \mathbb{N}} M_n(\mathbb{C})$$

Definition:

Let $g : [a, b] \rightarrow \mathbb{C}$ and $D \subset \mathbb{H}$ denote the set of self adjoint matrices with their spectrum in $[a, b]$. Then

$$g : D \longrightarrow \mathcal{M}$$

$$X = U \Lambda U^* \longmapsto U \begin{bmatrix} g(\lambda_1) & & \\ & \ddots & \\ & & g(\lambda_n) \end{bmatrix} U^*.$$

Let \mathbb{H}_n be the set of $n \times n$ self adjoint matrices, and define

$$\mathbb{H} = \bigcup_{n \in \mathbb{N}} \mathbb{H}_n, \quad \mathcal{M} = \bigcup_{n \in \mathbb{N}} M_n(\mathbb{C})$$

Definition:

Let $g : [a, b] \rightarrow \mathbb{C}$ and $D \subset \mathbb{H}$ denote the set of self adjoint matrices with their spectrum in $[a, b]$. Then

$$g : D \longrightarrow \mathcal{M}$$

$$X = U \Lambda U^* \longmapsto U \begin{bmatrix} g(\lambda_1) & & \\ & \ddots & \\ & & g(\lambda_n) \end{bmatrix} U^*.$$

Important: In this functional calculus,

$$g(X \oplus Y) = g(X) \oplus g(Y)$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

- With the polynomial case in mind, we can extend a general function. First, a piece of notation
- Lets grab a function on the real line and the self adjoint matrices with their spectrum in that domain
- Then we can lift g by emulating the behavior of polynomials.
- unwrap a self adjoint matrix, apply g to the diagonal, then wrap it back up
- Something to notice about this functional calculus—it treats direct sums *very* well
- This is all well and good, but can we do anything with these new functions?

Let \mathbb{H}_n be the set of $n \times n$ self adjoint matrices, and define

$$\mathbb{H} = \bigcup_{n \in \mathbb{N}} \mathbb{H}_n, \quad \mathcal{M} = \bigcup_{n \in \mathbb{N}} M_n(\mathbb{C})$$

Definition:

Let $g : [a, b] \rightarrow \mathbb{C}$ and $D \subset \mathbb{H}$ denote the set of self adjoint matrices with their spectrum in $[a, b]$. Then

$$g : D \longrightarrow \mathcal{M}$$

$$X = U \Lambda U^* \longmapsto U \begin{bmatrix} g(\lambda_1) & & \\ & \ddots & \\ & & g(\lambda_n) \end{bmatrix} U^*.$$

Important: In this functional calculus,

$$g(X \oplus Y) = g(X) \oplus g(Y)$$

Directional Derivative

Definition: *Directional Derivative*

Fix some $X \in \mathbb{H}_n$. The derivative of f at X in the direction $H \in M_n(\mathbb{C})$ is

$$Df(X)[H] = \lim_{t \rightarrow 0} \frac{f(X + tH) - f(X)}{t}$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Directional Derivative

Definition: Directional Derivative
Fix some $X \in \mathbb{H}_n$. The derivative of f at X in the direction $H \in M_n(\mathbb{C})$ is

$$Df(X)[H] = \lim_{t \rightarrow 0} \frac{f(X + tH) - f(X)}{t}$$

- We can define a directional derivative—as long as we are careful to have the direction in the same “level-wise” slice.
- Notice that, with some special attention to what operation we are carrying out, this is the exact same definition as classic multivariable calculus.
- There is another formulation that is (generally) more useful for computation

Directional Derivative

Definition: *Directional Derivative*

Fix some $X \in \mathbb{H}_n$. The derivative of f at X in the direction $H \in M_n(\mathbb{C})$ is

$$Df(X)[H] = \lim_{t \rightarrow 0} \frac{f(X + tH) - f(X)}{t}$$

Alternatively,

$$Df(X)[H] = \left. \frac{df(X + tH)}{dt} \right|_{t=0}$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Directional Derivative

- We can define a directional derivative—as long as we are careful to have the direction in the same “level-wise” slice.
- Notice that, with some special attention to what operation we are carrying out, this is the exact same definition as classic multivariable calculus.
- There is another formulation that is (generally) more useful for computation

Directional Derivative

Definition: Directional Derivative
Fix some $X \in \mathbb{H}_n$. The derivative of f at X in the direction $H \in M_n(\mathbb{C})$ is

$$Df(X)[H] = \lim_{t \rightarrow 0} \frac{f(X + tH) - f(X)}{t}$$

Alternatively,

$$Df(X)[H] = \left. \frac{df(X + tH)}{dt} \right|_{t=0}$$

Example: $g(x) = x^3$

$$g(X + tH) =$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example: $g(x) = x^3$ Example: $g(x) = x^3$

$$g(X + tH) =$$

- Now we consider an example. Since $Df(X)[H]$ is linear, we can just work with a single monomial
- First we expand $(x + th)^3$ —but we can't use the binomial theorem since x and h don't commute
- Once we expand, we take standard derivatives w.r.t t —treating X and H as formal symbols.

Example: $g(x) = x^3$

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example: $g(x) = x^3$ Example: $g(x) = x^3$

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

- Now we consider an example. Since $Df(X)[H]$ is linear, we can just work with a single monomial
- First we expand $(x + th)^3$ —but we can't use the binomial theorem since x and h don't commute
- Once we expand, we take standard derivatives w.r.t t —treating X and H as formal symbols.

Example: $g(x) = x^3$

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

From here, we can calculate:

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example: $g(x) = x^3$ Example: $g(x) = x^3$

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

From here, we can calculate:

- Now we consider an example. Since $Df(X)[H]$ is linear, we can just work with a single monomial
- First we expand $(x + th)^3$ —but we can't use the binomial theorem since x and h don't commute
- Once we expand, we take standard derivatives w.r.t t —treating X and H as formal symbols.

Example: $g(x) = x^3$

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

From here, we can calculate:

$$\frac{d}{dt}g(X + tH) = X^2H + XHX + 2tXH^2 + HX^2 \\ + 2tHXX + 2tH^2X + 3t^2H^3$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example: $g(x) = x^3$ Example: $g(x) = x^3$

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

From here, we can calculate:

$$\frac{d}{dt}g(X + tH) = X^2H + XHX + 2tXH^2 + HX^2 \\ + 2tHXX + 2tH^2X + 3t^2H^3$$

- Now we consider an example. Since $Df(X)[H]$ is linear, we can just work with a single monomial
- First we expand $(x + th)^3$ —but we can't use the binomial theorem since x and h don't commute
- Once we expand, we take standard derivatives w.r.t t —treating X and H as formal symbols.

Example: $g(x) = x^3$

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

From here, we can calculate:

$$\frac{d}{dt}g(X + tH) = X^2H + XHX + 2tXH^2 + HX^2 \\ + 2tHXX + 2tH^2X + 3t^2H^3$$

$$\frac{d^2}{dt^2}g(X + tH) = 2XH^2 + 2HXX + 2H^2X + 6tH^3$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example: $g(x) = x^3$ Example: $g(x) = x^3$

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

From here, we can calculate:

$$\frac{d}{dt}g(X + tH) = X^2H + XHX + 2tXH^2 + HX^2 \\ + 2tHXX + 2tH^2X + 3t^2H^3$$

$$\frac{d^2}{dt^2}g(X + tH) = 2XH^2 + 2HXX + 2H^2X + 6tH^3$$

- Now we consider an example. Since $Df(X)[H]$ is linear, we can just work with a single monomial
- First we expand $(x + th)^3$ —but we can't use the binomial theorem since x and h don't commute
- Once we expand, we take standard derivatives w.r.t t —treating X and H as formal symbols.

Example: $g(x) = x^3$

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

From here, we can calculate:

$$\frac{d}{dt}g(X + tH) = X^2H + XHX + 2tXH^2 + HX^2 \\ + 2tHXX + 2tH^2X + 3t^2H^3$$

$$\frac{d^2}{dt^2}g(X + tH) = 2XH^2 + 2HXX + 2H^2X + 6tH^3$$

$$\frac{d^3}{dt^3}g(X + tH) = 6H^3.$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example: $g(x) = x^3$ Example: $g(x) = x^3$

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

From here, we can calculate:

$$\frac{d}{dt}g(X + tH) = X^2H + XHX + 2tXH^2 + HX^2 \\ + 2tHXX + 2tH^2X + 3t^2H^3$$

$$\frac{d^2}{dt^2}g(X + tH) = 2XH^2 + 2HXX + 2H^2X + 6tH^3$$

$$\frac{d^3}{dt^3}g(X + tH) = 6H^3.$$

- Now we consider an example. Since $Df(X)[H]$ is linear, we can just work with a single monomial
- First we expand $(x + th)^3$ —but we can't use the binomial theorem since x and h don't commute
- Once we expand, we take standard derivatives w.r.t t —treating X and H as formal symbols.

Example: $g(x) = x^3$

And so the first 3 directional derivatives are:

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example: $g(x) = x^3$

Example: $g(x) = x^3$

And so the first 3 directional derivatives are:

- Now all we have to do is set $t = 0$ in the previous expressions and we get the first three directional derivatives
- Note that they are all in the same direction—mixed derivatives are possible but we won't need them.

Example: $g(x) = x^3$

And so the first 3 directional derivatives are:

$$Df(X)[H] = X^2H + XHX + HX^2$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example: $g(x) = x^3$ Example: $g(x) = x^3$

And so the first 3 directional derivatives are:

$$Df(X)[H] = X^2H + XHX + HX^2$$

- Now all we have to do is set $t = 0$ in the previous expressions and we get the first three directional derivatives
- Note that they are all in the same direction—mixes derivatives are possible but we won't need them.

Example: $g(x) = x^3$

And so the first 3 directional derivatives are:

$$Df(X)[H] = X^2H + XHX + HX^2$$

$$D^{(2)}f(X)[H] = 2XH^2 + 2HXX + 2H^2X$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example: $g(x) = x^3$ Example: $g(x) = x^3$

And so the first 3 directional derivatives are:

$$Df(X)[H] = X^2H + XHX + HX^2$$

$$D^{(2)}f(X)[H] = 2XH^2 + 2HXX + 2H^2X$$

- Now all we have to do is set $t = 0$ in the previous expressions and we get the first three directional derivatives
- Note that they are all in the same direction—mixes derivatives are possible but we won't need them.

Example: $g(x) = x^3$

And so the first 3 directional derivatives are:

$$Df(X)[H] = X^2H + XHX + HX^2$$

$$D^{(2)}f(X)[H] = 2XH^2 + 2HXXH + 2H^2X$$

$$D^{(3)}f(X)[H] = 6H^3$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example: $g(x) = x^3$

- Now all we have to do is set $t = 0$ in the previous expressions and we get the first three directional derivatives
- Note that they are all in the same direction—mixes derivatives are possible but we won't need them.

Example: $g(x) = x^3$

And so the first 3 directional derivatives are:

$$Df(X)[H] = X^2H + XHX + HX^2$$

$$D^{(2)}f(X)[H] = 2XH^2 + 2HXXH + 2H^2X$$

$$D^{(3)}f(X)[H] = 6H^3$$

Part I.5: Objects and Maps

A Second Attempt

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

Part I.5: Objects and Maps
A Second Attempt

- In seeking a more general theory we need to leave the world of this “SA functional calculus” behind.
- Rather than lifting functions to be matrix valued, we will define *new* objects that behave like those we just looked at.

Some Definitions

Definition:

The g -dimensional **Matrix Universe** is

$$\mathcal{M}^g = \bigcup_{n \in \mathbb{N}} (M_n(\mathbb{C}))^g$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

└ Some Definitions

Definition:

The g -dimensional **Matrix Universe** is

$$\mathcal{M}^g = \bigcup_{n \in \mathbb{N}} (M_n(\mathbb{C}))^g$$

- Definition of the matrix universe — g tuples of matrices of all sizes
- By convetion, tuples are likes size

Some Definitions

Definition:

The g -dimensional **Matrix Universe** is

$$\mathcal{M}^g = \bigcup_{n \in \mathbb{N}} (M_n(\mathbb{C}))^g$$

By convention, $X \in \mathcal{M}^g$ is a tuple of like-size matrices

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

└ Some Definitions

Definition:

The g -dimensional **Matrix Universe** is

$$\mathcal{M}^g = \bigcup_{n \in \mathbb{N}} (M_n(\mathbb{C}))^g$$

By convention, $X \in \mathcal{M}^g$ is a tuple of like-size matrices

- Definition of the matrix universe — g tuples of matrices of all sizes
- By convetion, tuples are likes size

Definition:

We say $D \subset \mathcal{M}^g$ is a **free set** if it is closed with respect to direct sums and unitary conjugation. That is,

.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

Definition:
We say $D \subset \mathcal{M}^g$ is a **free set** if it is closed with respect to direct sums and unitary conjugation. That is,

- In math we often think about substructures that capture the implicit structure our space (subgroup, subspace, etc)
- In the nc setting, this is a *free set*, also called nc set
- direct sums and unitary conjugation are component wise
- If you see a D , you can assume that it is a free set.
- A subscript denotes a level-wise slice
- Note that this requires a lot of structure on free sets—we want to put a name to these structure.

Definition:

We say $D \subset \mathcal{M}^g$ is a **free set** if it is closed with respect to direct sums and unitary conjugation. That is,

$$\textcircled{1} \quad X, Y \in D \text{ means } X \oplus Y = (X_1 \oplus Y_1, \dots, X_g \oplus Y_g) \in D.$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

Definition:
We say $D \subset \mathcal{M}^g$ is a **free set** if it is closed with respect to direct sums and unitary conjugation. That is,
 $\bullet \quad X, Y \in D \text{ means } X \oplus Y = (X_1 \oplus Y_1, \dots, X_g \oplus Y_g) \in D.$

- In math we often think about substructures that capture the implicit structure our space (subgroup, subspace, etc)
- In the nc setting, this is a *free set*, also called nc set
- direct sums and unitary conjugation are component wise
- If you see a D , you can assume that it is a free set.
- A subscript denotes a level-wise slice
- Note that this requires a lot of structure on free sets—we want to put a name to these structure.

Definition:

We say $D \subset \mathcal{M}^g$ is a **free set** if it is closed with respect to direct sums and unitary conjugation. That is,

- ① $X, Y \in D$ means $X \oplus Y = (X_1 \oplus Y_1, \dots, X_g \oplus Y_g) \in D$.
- ② For X, U like-size matrices with U unitary and $X \in D$, then $UXU^* = (UX_1 U^*, \dots, UX_g U^*) \in D$.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

Definition:

We say $D \subset \mathcal{M}^g$ is a **free set** if it is closed with respect to direct sums and unitary conjugation. That is,

- $X, Y \in D$ means $X \oplus Y = (X_1 \oplus Y_1, \dots, X_g \oplus Y_g) \in D$.
- For X, U like-size matrices with U unitary and $X \in D$, then $UXU^* = (UX_1 U^*, \dots, UX_g U^*) \in D$.

- In math we often think about substructures that capture the implicit structure our space (subgroup, subspace, etc)
- In the nc setting, this is a *free set*, also called nc set
- direct sums and unitary conjugation are component wise
- If you see a D , you can assume that it is a free set.
- A subscript denotes a level-wise slice
- Note that this requires a lot of structure on free sets—we want to put a name to these structure.

Definition:

We say $D \subset \mathcal{M}^g$ is a **free set** if it is closed with respect to direct sums and unitary conjugation. That is,

- ① $X, Y \in D$ means $X \oplus Y = (X_1 \oplus Y_1, \dots, X_g \oplus Y_g) \in D$.
- ② For X, U like-size matrices with U unitary and $X \in D$, then $UXU^* = (UX_1U^*, \dots, UX_gU^*) \in D$.

For D a free set, define $D_n = D \cap (M_n(\mathbb{C}))^g$.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

Definition:
We say $D \subset \mathcal{M}^g$ is a **free set** if it is closed with respect to direct sums and unitary conjugation. That is,
 • $X, Y \in D$ means $X \oplus Y = (X_1 \oplus Y_1, \dots, X_g \oplus Y_g) \in D$.
 • For X, U like-size matrices with U unitary and $X \in D$, then $UXU^* = (UX_1U^*, \dots, UX_gU^*) \in D$.
 For D a free set, define $D_n = D \cap (M_n(\mathbb{C}))^g$.

- In math we often think about substructures that capture the implicit structure our space (subgroup, subspace, etc)
- In the nc setting, this is a *free set*, also called nc set
- direct sums and unitary conjugation are component wise
- If you see a D , you can assume that it is a free set.
- A subscript denotes a level-wise slice
- Note that this requires a lot of structure on free sets—we want to put a name to these structure.

Definition:

Given $X \in \mathcal{M}^g$, a tuple of $n \times n$ matrices, the **fiber** of X is the set

$$\{X^{\oplus k} \mid k \in \mathbb{N}\}.$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

Definition:

Given $X \in \mathcal{M}^g$, a tuple of $n \times n$ matrices, the **fiber** of X is the set

$$\{X^{\oplus k} \mid k \in \mathbb{N}\}.$$

- The fiber is all the points “above” a given point. Conceptually, we imagine identification along the fiber—this will become important when we start doing topology
- The envelope (which will be less important to us) is the unitary smearing of the fiber at each level.
- This is a good spot for questions.

Definition:

Given $X \in \mathcal{M}^g$, a tuple of $n \times n$ matrices, the **fiber** of X is the set

$$\{X^{\oplus k} \mid k \in \mathbb{N}\}.$$

Definition:

Given $X \in \mathcal{M}^g$, a tuple of $n \times n$ matrices, the **envelope** of X is the set

$$\{U^* X^{\oplus k} U \mid k \in \mathbb{N}, U \text{ Unitary}\}.$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

Definition:

Given $X \in \mathcal{M}^g$, a tuple of $n \times n$ matrices, the **fiber** of X is the set

$$\{X^{\oplus k} \mid k \in \mathbb{N}\}.$$

Definition:

Given $X \in \mathcal{M}^g$, a tuple of $n \times n$ matrices, the **envelope** of X is the set

$$\{U^* X^{\oplus k} U \mid k \in \mathbb{N}, U \text{ Unitary}\}.$$

- The fiber is all the points “above” a given point. Conceptually, we imagine identification along the fiber—this will become important when we start doing topology
- The envelope (which will be less important to us) is the unitary smearing of the fiber at each level.
- This is a good spot for questions.

A bit of topology

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

└ A bit of topology

- If we are going to look for holes and build up the algebraic topology, we need a point set topology first—so there is a natural question.
- Bad news: there isn't a natural choice
- There are a handful of candidates (fine, fat, free, nc Zariski). I wish we had time to go into detail.
- For us, free sets are open if their level-wise restriction is open
- Other point-set characterizations are similar.

A bit of topology

What does it mean for $D \subset \mathcal{M}^g$ to be open?

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

└ A bit of topology

- If we are going to look for holes and build up the algebraic topology, we need a point set topology first—so there is a natural question.
- Bad news: there isn't a natural choice
- There are a handful of candidates (fine, fat, free, nc Zariski). I wish we had time to go into detail.
- For us, free sets are open if their level-wise restriction is open
- Other point-set characterizations are similar.

A bit of topology

What does it mean for $D \subset \mathcal{M}^g$ to be open?

- There is not a canonical topology

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

└ A bit of topology

- There is not a canonical topology

- If we are going to look for holes and build up the algebraic topology, we need a point set topology first—so there is a natural question.
- Bad news: there isn't a natural choice
- There are a handful of candidates (fine, fat, free, nc Zariski). I wish we had time to go into detail.
- For us, free sets are open if their level-wise restriction is open
- Other point-set characterizations are similar.

A bit of topology

What does it mean for $D \subset \mathcal{M}^g$ to be open?

- There is not a canonical topology
- For us, we will say that D is open if each D_n is open.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

└ A bit of topology

What does it mean for $D \subset \mathcal{M}^g$ to be open?

- There is not a canonical topology
- For us, we will say that D is open if each D_n is open.

- If we are going to look for holes and build up the algebraic topology, we need a point set topology first—so there is a natural question.
- Bad news: there isn't a natural choice
- There are a handful of candidates (fine, fat, free, nc Zariski). I wish we had time to go into detail.
- For us, free sets are open if their level-wise restriction is open
- Other point-set characterizations are similar.

A bit of topology

What does it mean for $D \subset \mathcal{M}^g$ to be open?

- There is not a canonical topology
- For us, we will say that D is open if each D_n is open.
- Simply connected, connected, bounded, etc. are defined similarly.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

└ A bit of topology

What does it mean for $D \subset M^g$ to be open?

- There is not a canonical topology
- For us, we will say that D is open if each D_n is open.
- Simply connected, connected, bounded, etc. are defined similarly.

- If we are going to look for holes and build up the algebraic topology, we need a point set topology first—so there is a natural question.
- Bad news: there isn't a natural choice
- There are a handful of candidates (fine, fat, free, nc Zariski). I wish we had time to go into detail.
- For us, free sets are open if their level-wise restriction is open
- Other point-set characterizations are similar.

What the natural functions on \mathcal{M}^g ?

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

└ What the natural functions on \mathcal{M}^g ?

- We have our objects, but what are the maps?
- Free functions are defined to be anything that behaves like a polynomial
- tracial functions look like traces
- For both of these maps, the directional derivative is defined identically as before—but tracial functions get something extra.

What the natural functions on \mathcal{M}^g ?**Definition:**

A function $f : D \rightarrow \mathcal{M}^{\hat{g}}$ is called **free** if

- ① $f(X \oplus Y) = f(X) \oplus f(Y)$
- ② $f(UXU^*) = f(U)f(X)f(U^*)$ where X and U are like-size and U is unitary.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

└ What the natural functions on \mathcal{M}^g ?

- We have our objects, but what are the maps?
- Free functions are defined to be anything that behaves like a polynomial
- tracial functions look like traces
- For both of these maps, the directional derivative is defined identically as before—but tracial functions get something extra.

Definition:

A function $f : D \rightarrow \mathcal{M}^{\hat{g}}$ is called free if

- $f(X \oplus Y) = f(X) \oplus f(Y)$
- $f(UXU^*) = f(U)f(X)f(U^*)$ where X and U are like-size and U is unitary.

What the natural functions on \mathcal{M}^g ?**Definition:**

A function $f : D \rightarrow \mathcal{M}^{\hat{g}}$ is called **free** if

- ① $f(X \oplus Y) = f(X) \oplus f(Y)$
- ② $f(UXU^*) = f(U)f(X)f(U^*)$ where X and U are like-size and U is unitary.

Definition:

A function $f : D \rightarrow \mathbb{C}$ is a **tracial function** if

- ① $f(X \oplus Y) = f(X) + f(Y)$
- ② $f(UXU^*) = f(X)$ where X and U are like-size and U is unitary.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

└ What the natural functions on \mathcal{M}^g ?

- We have our objects, but what are the maps?
- Free functions are defined to be anything that behaves like a polynomial
- tracial functions look like traces
- For both of these maps, the directional derivative is defined identically as before—but tracial functions get something extra.

Definition:

A function $f : D \rightarrow \mathcal{M}^{\hat{g}}$ is called **free** if

- $f(X \oplus Y) = f(X) \oplus f(Y)$
- $f(UXU^*) = f(U)f(X)f(U^*)$ where X and U are like-size and U is unitary.

Definition:

A function $f : D \rightarrow \mathbb{C}$ is a **tracial function** if

- $f(X \oplus Y) = f(X) + f(Y)$
- $f(UXU^*) = f(X)$ where X and U are like-size and U is unitary.

Definition:

Given a tracial function f , the free gradient, ∇f , is the unique free function satisfying

$$\text{tr}(H \cdot \nabla f(X)) = Df(X)[H],$$

where, if $A = (A_1, \dots, A_g)$ and $B = (B_1, \dots, B_g)$ are tuples of like-size matrices then $A \cdot B = \sum_{i=1}^g A_i B_i$.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Uniqueness of the Gradient

Definition:
Given a tracial function f , the free gradient, ∇f , is the unique free function satisfying

$$\text{tr}(H \cdot \nabla f(X)) = Df(X)[H],$$

where, if $A = (A_1, \dots, A_g)$ and $B = (B_1, \dots, B_g)$ are tuples of like-size matrices then $A \cdot B = \sum_{i=1}^g A_i B_i$.

- The ∇ of a free function is the unique free function satisfy this equation—where the \cdot is just like the dot product
- Whenever you see $\text{tr}(\cdot)$ I want you to think of the inner product—it is slightly distinct but it will make a lot of things make more sense
- Some of you may be hesitant at the fact that I claim ∇ is unique. Why should this be true?

Why should ∇f be unique?

Theorem (Trace Duality)

Let f, g be free functions $\mathcal{M}^g \rightarrow \mathcal{M}^{\tilde{g}}$. If $\text{tr}(H \cdot f) = \text{tr}(H \cdot g)$ for all tuples H , then $f = g$.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Uniqueness of the Gradient

└ Why should ∇f be unique?

- $f = g$ whenever the domains overlap
- In the vector space setting—with an inner product—this is a fairly immediate result. You would show it by picking vectors of all 0's and a single 1.
- You prove this identically—but with coordinate matrices instead of coordinate vectors.
- Before we look at the algebraic topology, we need to take a brief trip to complex variable land

Part II: Analytic Continuation and Monodromy

2022-06-03


Searching for Holes in the Matrix Universe

└ Part II: Monodromy

Part II: Analytic Continuation and Monodromy

- analytic continuation and monodromy is the link between complex analysis and topology.

Analytic Continuation




2022-06-03

Searching for Holes in the Matrix Universe

└ Part II: Monodromy


└ Analytic Continuation

└ Analytic Continuation



- Lets say we have some analytic function defined on Ω and a curve γ taking α to β .
- We can expand a power series about α with some radius of convergence. But since f is analytic, we can expand about some any point on γ that is still in the red disk

Analytic Continuation




2022-06-03

Searching for Holes in the Matrix Universe

└ Part II: Monodromy


└ Analytic Continuation

└ Analytic Continuation



- But now we have a new radius of convergence! Importantly this will agree with the original function on that initial overlap
- We can keep doing this!

Analytic Continuation




2022-06-03

Searching for Holes in the Matrix Universe

└ Part II: Monodromy


└ Analytic Continuation

└ Analytic Continuation



- Now we have a third power series representation for f —once again, it will agree with our last expansions where those two disks overlap.
- As long as γ stays away from any potential poles, we can keep doing this all the way to β

Analytic Continuation




2022-06-03

Searching for Holes in the Matrix Universe


└ Part II: Monodromy

└ Analytic Continuation

└ Analytic Continuation



- After repeatedly expanding, we finally have an analytic function at β !
- As we said, we need γ to avoid poles, but what can we say about the uniqueness of the analytic function at β ?


Example: Analytically continuing $\text{Log } z$ 

2022-06-03


Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Analytic Continuation

└ Example: Analytically continuing $\text{Log } z$ Example: Analytically continuing $\text{Log } z$ 

- Now we have another example: consider the principle branch of the complex logarithm.
- if we analytically continue along γ_1 , then we can evaluate $\text{Log}(-1)$.


Example: Analytically continuing $\text{Log } z$ 

2022-06-03


Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Analytic Continuation

└ Example: Analytically continuing $\text{Log } z$ Example: Analytically continuing $\text{Log } z$ 

- when we do this, we see that the $\text{Log}(-1) = -\pi i$.
- But what about the other way? What if we continued along a path that went through the UHP?


Example: Analytically continuing $\text{Log } z$ 

2022-06-03


Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Analytic Continuation

└ Example: Analytically continuing $\text{Log } z$ Example: Analytically continuing $\text{Log } z$ 

- when we do this, we get $\text{Log}(-1) = \pi i$! They disagree!
- what's even stranger is what happens when we keep going on a circle around the origin.


Example: Analytically continuing $\text{Log } z$ 

2022-06-03

Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Analytic Continuation

└ Example: Analytically continuing $\text{Log } z$ Example: Analytically continuing $\text{Log } z$ 

- amazingly, when you continue all the way around and compute $f(1)$, you get $2\pi i$ —not 1.
- What is going on here? when are two analytic continuations equal?

When are two analytic continuations equal?

2022-06-03

Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Monodromy

└ When are two analytic continuations equal?

- It is in answering this question that we see the deep link between analytic continuation and topology.
- Monodromy 1: the paths are homotopic and the function analytically continues along all of the intermediate paths, then you are golden!
- a picture for those who like that!
- Note that this tells us that analytic continuation searches for holes
- when seek to lift this idea to a nc case, it will serve much better to consider an alternate characterization

When are two analytic continuations equal?

Theorem (Monodromy I)

Let γ_1, γ_2 be two paths from α to β and Γ_s be a fixed-endpoint homotopy between them. If f can be continued along Γ_s for all $s \in [0, 1]$, then the continuations along γ_1 and γ_2 agree at β .

2022-06-03

Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Monodromy


└ When are two analytic continuations equal?

- It is in answering this question that we see the deep link between analytic continuation and topology.
- Monodromy 1: the paths are homotopic and the function analytically continues along all of the intermediate paths, then you are golden!
- a picture for those who like that!
- Note that this tells us that analytic continuation searches for holes
- when seek to lift this idea to a nc case, it will serve much better to consider an alternate characterization

When are two analytic continuations equal?

Theorem (Monodromy I)

Let γ_1, γ_2 be two paths from α to β and Γ_s be a fixed-endpoint homotopy between them. If f can be continued along Γ_s for all $s \in [0, 1]$, then the continuations along γ_1 and γ_2 agree at β .



2022-06-03

Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Monodromy

└ When are two analytic continuations equal?

- It is in answering this question that we see the deep link between analytic continuation and topology.
- Monodromy 1: the paths are homotopic and the function analytically continues along all of the intermediate paths, then you are golden!
- a picture for those who like that!
- Note that this tells us that analytic continuation searches for holes
- when seek to lift this idea to a nc case, it will serve much better to consider an alternate characterization

When are two analytic continuations equal?

Theorem (Monodromy I)

Let γ_1, γ_2 be two paths from α to β and Γ_s be a fixed-endpoint homotopy between them. If f can be continued along Γ_s for all $s \in [0, 1]$, then the continuations along γ_1 and γ_2 agree at β .



When are two analytic continuations equal?

Theorem (Monodromy II)

Let $U \subset \mathbb{C}$ be a disk in \mathbb{C} centered at z_0 and $f : U \rightarrow \mathbb{C}$ an analytic function. If W is an open, simply connected set containing U and f continues along any path $\gamma \subset W$ starting at z_0 , then f has a unique extension to all of W .

2022-06-03

Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Monodromy

└ When are two analytic continuations equal?

When are two analytic continuations equal?

Theorem (Monodromy II)

Let $U \subset \mathbb{C}$ be a disk in \mathbb{C} centered at z_0 and $f : U \rightarrow \mathbb{C}$ an analytic function. If W is an open, simply connected set containing U and f continues along any path $\gamma \subset W$ starting at z_0 , then f has a unique extension to all of W .

- We require simply connected in the big set! Thus, holes get in the way of a unique extension.
- Obviously we don't have the time to go into explicit detail about why, but it turn out that you can realize the fundamental group of some open, connected, subset of \mathbb{C} simply by looking at the analytic continuation of functions!

What about the nc case?

Before we look at a free analogue of the monodromy theorem, we need to ask an important question: What does it mean for a free function to be analytic?

2022-06-03

Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Free Monodromy

└ What about the nc case?

What about the nc case?

Before we look at a free analogue of the monodromy theorem, we need to ask an important question: What does it mean for a free function to be analytic?

- Now that we have this (very power) theorem from classical complex analysis, if we want an nc analogue—what is an “analytic” free function.
- As with our other characterizations, a free function is analytic if it is analytic as a function on each D_n .
- Even more surprisingly, we have a *wild* characterization due to Agler and McCarthy
- This is incredibly powerful—I will let you draw your own conclusion as to what it says about the underlying point set topology.
- With our question answered, we can present the free analogue.

What about the nc case?

Before we look at a free analogue of the monodromy theorem, we need to ask an important question: What does it mean for a free function to be analytic?

A: $f : D \rightarrow \mathcal{M}^{\hat{g}}$ is analytic if it is analytic as a function of each D_n .

2022-06-03

Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Free Monodromy

└ What about the nc case?

- Now that we have this (very power) theorem from classical complex analysis, if we want an nc analogue—what is an “analytic” free function.
- As with our other characterizations, a free function is analytic if it is analytic as a function on each D_n .
- Even more surprisingly, we have a *wild* characterization due to Agler and McCarthy
- This is incredibly powerful—I will let you draw your own conclusion as to what it says about the underlying point set topology.
- With our question answered, we can present the free analogue.

What about the nc case?

Before we look at a free analogue of the monodromy theorem, we need to ask an important question: What does it mean for a free function to be analytic?

A: $f : D \rightarrow \mathcal{M}^{\hat{g}}$ is analytic if it is analytic as a function of each D_n .

What about the nc case?

Before we look at a free analogue of the monodromy theorem, we need to ask an important question: What does it mean for a free function to be analytic?

A: $f : D \rightarrow \mathcal{M}^{\hat{g}}$ is analytic if it is analytic as a function of each D_n .

Theorem (Agler, McCarthy (2016))

Let $f : D \rightarrow \mathcal{M}^{\hat{g}}$ be a free function. If f is locally bounded on each D_n , then f is an analytic free function.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Free Monodromy

└ What about the nc case?

- Now that we have this (very power) theorem from classical complex analysis, if we want an nc analogue—what is an “analytic” free function.
- As with our other characterizations, a free function is analytic if it is analytic as a function on each D_n .
- Even more surprisingly, we have a *wild* characterization due to Agler and McCarthy
- This is incredibly powerful—I will let you draw your own conclusion as to what it says about the underlying point set topology.
- With our question answered, we can present the free analogue.

What about the nc case?

Before we look at a free analogue of the monodromy theorem, we need to ask an important question: What does it mean for a free function to be analytic?

A: $f : D \rightarrow \mathcal{M}^{\hat{g}}$ is analytic if it is analytic as a function of each D_n .

Theorem (Agler, McCarthy (2016))

Let $f : D \rightarrow \mathcal{M}^{\hat{g}}$ be a free function. If f is locally bounded on each D_n , then f is an analytic free function.

Theorem (Free Universal Monodromy, Pascoe 2020)

Let f be an analytic free function defined on some ball $B \subset D$, for D an open, connected free set. If f analytically continues along every path in D , then f has a unique analytic continuation to all of D .

2022-06-03

Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Free Monodromy

Theorem (Free Universal Monodromy, Pascoe 2020)

Let f be an analytic free function defined on some ball $B \subset D$, for D an open, connected free set. If f analytically continues along every path in D , then f has a unique analytic continuation to all of D .

- all you need to have a unique extension is to continue along every path!
- The larger set doesn't have to be simply connected! This is huge!
- While this theorem is amazing, it is the bearer of bad news

Consequences of Free Monodromy

2022-06-03

Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Free Monodromy

└ Consequences of Free Monodromy

- There are two major (and related) consequences to free monodromy
- First, free functions cannot detect holes via analytic continuation
- Therefore, if we want a fundamental group that is governed by analytic continuation, we need to look elsewhere
- before we transition to the fundamental group, any questions.

Consequences of Free Monodromy

- 1 Free functions can't detect holes!

2022-06-03

Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Free Monodromy

└ Consequences of Free Monodromy

- Free functions can't detect holes!

- There are two major (and related) consequences to free monodromy
- First, free functions cannot detect holes via analytic continuation
- Therefore, if we want a fundamental group that is governed by analytic continuation, we need to look elsewhere
- before we transition to the fundamental group, any questions.

Consequences of Free Monodromy

- ① Free functions can't detect holes!
- ② If we want a fundamental group governed by analytic continuation, we need to look elsewhere.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Free Monodromy

└ Consequences of Free Monodromy

● Free functions can't detect holes!

● If we want a fundamental group governed by analytic continuation, we need to look elsewhere.

- There are two major (and related) consequences to free monodromy
- First, free functions cannot detect holes via analytic continuation
- Therefore, if we want a fundamental group that is governed by analytic continuation, we need to look elsewhere
- before we transition to the fundamental group, any questions.

Part III: Homotopy

- In part III we look at two types of fundamental group.
- Since free monodromy says that free functions won't give us a π_1 we start by looking at a fundamental group that is divorced from analytic continuation

Definition:

A continuous function $\gamma : [0, 1] \rightarrow D$ **essentially takes** X to Y if

$$\gamma(0) = X^{\oplus \ell}, \text{ for some } \ell \in \mathbb{N}$$

$$\gamma(1) = Y^{\oplus k}, \text{ for some } k \in \mathbb{N}.$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part III: Homotopy

└ A First Fundamental Group

Definition:
A continuous function $\gamma : [0, 1] \rightarrow D$ **essentially takes** X to Y if

$$\gamma(0) = X^{\oplus \ell}, \text{ for some } \ell \in \mathbb{N}$$

$$\gamma(1) = Y^{\oplus k}, \text{ for some } k \in \mathbb{N}.$$


- recall that the fiber of a point in \mathcal{M}^g is all direct sum copies of that point—and futher that we consider everything in the fiber somehow “the same”
- An essential path it a traditon path between the fibers!
- In order for us to create a group out of these paths, we a concatenation product.

Definition:

A continuous function $\gamma : [0, 1] \rightarrow D$ **essentially takes** X to Y if

$$\gamma(0) = X^{\oplus \ell}, \text{ for some } \ell \in \mathbb{N}$$

$$\gamma(1) = Y^{\oplus k}, \text{ for some } k \in \mathbb{N}.$$



2022-06-03

Searching for Holes in the Matrix Universe

└ Part III: Homotopy

└ A First Fundamental Group

Definition:
A continuous function $\gamma : [0, 1] \rightarrow D$ **essentially takes** X to Y if

$$\gamma(0) = X^{\oplus \ell}, \text{ for some } \ell \in \mathbb{N}$$

$$\gamma(1) = Y^{\oplus k}, \text{ for some } k \in \mathbb{N}.$$



- recall that the fiber of a point in \mathcal{M}^g is all direct sum copies of that point—and further that we consider everything in the fiber somehow “the same”
- An essential path is a traditon path between the fibers!
- In order for us to create a group out of these paths, we a concatenation product.

Given γ essentially taking X to Y and β taking Z to W , define

$$\gamma \oplus \beta(t) = \begin{bmatrix} \gamma(t) & 0 \\ 0 & \beta(t) \end{bmatrix}.$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part III: Homotopy

└ A First Fundamental Group

Given γ essentially taking X to Y and β taking Z to W , define

$$\gamma \oplus \beta(t) = \begin{bmatrix} \gamma(t) & 0 \\ 0 & \beta(t) \end{bmatrix}.$$

- First, we can take the direct sum of paths the exact way that you would expect.
- To concatenate the paths, the definition is almost identical—first you do one path twice as fast, then you do the other
- Except you direct sum “enough” times to make it continuous

Given γ essentially taking X to Y and β taking Z to W , define

$$\gamma \oplus \beta(t) = \begin{bmatrix} \gamma(t) & 0 \\ 0 & \beta(t) \end{bmatrix}.$$

Definition:

Let γ and β be paths taking X to Y and Y to Z respectively. We define their product to be the path essentially taking X to Z given by

$$\beta\gamma(t) = \begin{cases} \gamma^{\oplus k}(2t) & t \in [0, 0.5) \\ \beta^{\oplus \ell}(2t - 1) & t \in [0.5, 1] \end{cases}$$

where k and ℓ are positive integers chosen to make $\gamma^{\oplus k}$ and $\beta^{\oplus \ell}$ like size matrices for each $t \in [0, 1]$.

Searching for Holes in the Matrix Universe

└ Part III: Homotopy

└ A First Fundamental Group

2022-06-03

Given γ essentially taking X to Y and β taking Z to W , define

$$\gamma \oplus \beta(t) = \begin{bmatrix} \gamma(t) & 0 \\ 0 & \beta(t) \end{bmatrix}.$$

Definition:

Let γ and β be paths taking X to Y and Y to Z respectively. We define their product to be the path essentially taking X to Z given by

$$\beta\gamma(t) = \begin{cases} \gamma^{\oplus k}(2t) & t \in [0, 0.5) \\ \beta^{\oplus \ell}(2t - 1) & t \in [0.5, 1] \end{cases}$$

where k and ℓ are positive integers chosen to make $\gamma^{\oplus k}$ and $\beta^{\oplus \ell}$ like size matrices for each $t \in [0, 1]$.

- First, we can take the direct sum of paths the exact way that you would expect.
- To concatenate the paths, the definition is almost identical—first you do one path twice as fast, then you do the other
- Except you direct sum “enough” times to make it continuous

The Full Fundamental Group

For $D \subset \mathcal{M}^g$ a connected free set, the **full fundamenal group**, $\pi_1(D)$, is the group of paths essentially taking X to X up to homotopy equivalence and the relation $\gamma = \gamma^{\oplus k}$.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part III: Homotopy

└ A First Fundamental Group

└ The Full Fundamental Group

- That was all we needed to create the “full” fundamental group.
- group of paths up to homotopy equivalence and direct sum of paths
- You can show that this is abelian and divisible—but computationally we are totally stuck—we don’t have any tools to compute π *full*
- Instead, we can look at analytic continuation of *tracial* functions and see if that can get us anything.

Let $D \subset \mathcal{M}^g$ be a connected, open, free set. If there exists a nonempty, simply-connected, open, free $B \subset D$, then we say that D is **anchored**.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part III: Homotopy

└ A Second Fundamental Group

Let $D \subset \mathcal{M}^g$ be a connected, open, free set. If there exists a nonempty, simply-connected, open, free $B \subset D$, then we say that D is **anchored**.

- Since we are going to look at analytic continuation, we need a place to start the functions—this is the anchor
- the functions in question are the “global germs”, which are defined on the anchor but analytically along any path in the free set.
- to make sense of $f(\gamma)$ —analytically continue along γ , compute f of the endpoint, then normalize

Let $D \subset \mathcal{M}^g$ be a connected, open, free set. If there exists a nonempty, simply-connected, open, free $B \subset D$, then we say that D is **anchored**.

For D an anchored set, and $B \subset D$ its anchor, we call a tracial function $f : B \rightarrow \mathbb{C}$ a **global germ** if it analytically continues along every path in D which starts in B .

2022-06-03

Searching for Holes in the Matrix Universe

└ Part III: Homotopy

└ A Second Fundamental Group

- Since we are going to look at analytic continuation, we need a place to start the functions—this is the anchor
- the functions in question are the “global germs”, which are defined on the anchor but analytically along any path in the free set.
- to make sense of $f(\gamma)$ —analytically continue along γ , compute f of the endpoint, then normalize

Let $D \subset \mathcal{M}^g$ be a connected, open, free set. If there exists a nonempty, simply-connected, open, free $B \subset D$, then we say that D is **anchored**.

For D an anchored set, and $B \subset D$ its anchor, we call a tracial function $f : B \rightarrow \mathbb{C}$ a **global germ** if it analytically continues along every path in D which starts in B .

Let $D \subset \mathcal{M}^g$ be a connected, open, free set. If there exists a nonempty, simply-connected, open, free $B \subset D$, then we say that D is **anchored**.

For D an anchored set, and $B \subset D$ its anchor, we call a tracial function $f : B \rightarrow \mathbb{C}$ a **global germ** if it analytically continues along every path in D which starts in B .

For our purposes, we view γ as coupled with its endpoint. Thus, if γ essentially takes X to Y , then

$$f(\gamma) = \frac{1}{k} f(Y^{\oplus k}).$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part III: Homotopy

└ A Second Fundamental Group

- Since we are going to look at analytic continuation, we need a place to start the functions—this is the anchor
- the functions in question are the “global germs”, which are defined on the anchor but analytically along any path in the free set.
- to make sense of $f(\gamma)$ —analytically continue along γ , compute f of the endpoint, then normalize

Let $D \subset \mathcal{M}^g$ be a connected, open, free set. If there exists a nonempty, simply-connected, open, free $B \subset D$, then we say that D is **anchored**.

For D an anchored set, and $B \subset D$ its anchor, we call a tracial function $f : B \rightarrow \mathbb{C}$ a **global germ** if it analytically continues along every path in D which starts in B .

For our purposes, we view γ as coupled with its endpoint. Thus, if γ essentially takes X to Y , then

$$f(\gamma) = \frac{1}{k} f(Y^{\oplus k}).$$

Trace Equivalence

Definition:

Let $B \subset D$ be an anchor and fix $X \in B$. If γ and β both essentially take X to Y , we say they are **trace equivalent** if, for every global germ f and every path δ taking Y to Z ,

$$f(\delta\gamma) = f(\delta\beta).$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part III: Homotopy

└ A Second Fundamental Group

└ Trace Equivalence

Definition:

Let $B \subset D$ be an anchor and fix $X \in B$. If γ and β both essentially take X to Y , we say they are **trace equivalent** if, for every global germ f and every path δ taking Y to Z ,

$$f(\delta\gamma) = f(\delta\beta).$$

- For two paths essentially taking X to Y , we say they are **tr eq** if for all global germs $f(\delta\gamma) = f(\delta\beta)$.
- Both of these are computing $f(Z)$, but with different analytic contuations.
- Note that (from classical monodromy) this captures endpoint homotopy and the direct sum identity.

Trace Equivalence

Definition:

Let $B \subset D$ be an anchor and fix $X \in B$. If γ and β both essentially take X to Y , we say they are **trace equivalent** if, for every global germ f and every path δ taking Y to Z ,

$$f(\delta\gamma) = f(\delta\beta).$$

That is, trace equivalent paths are those which cannot be told apart via analytic continuation of global germ.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part III: Homotopy

└ A Second Fundamental Group

└ Trace Equivalence

Trace Equivalence

Definition:

Let $B \subset D$ be an anchor and fix $X \in B$. If γ and β both essentially take X to Y , we say they are **trace equivalent** if, for every global germ f and every path δ taking Y to Z ,

$$f(\delta\gamma) = f(\delta\beta).$$

That is, trace equivalent paths are those which cannot be told apart via analytic continuation of global germ.

- For two paths essentially taking X to Y , we say they are tr eq if for all global germs $f(\delta\gamma) = f(\delta\beta)$.
- Both of these are computing $f(Z)$, but with different analytic contuations.
- Note that (from classical monodromy) this captures endpoint homotopy and the direct sum identity.

The Tracial Fundamental Group

Let $D \subset \mathcal{M}^g$ be an anchored space with B is anchor. For $X \in B$ define $\pi_1^{\text{tr}}(D)$ to be the group of trace equivalent paths essentially taking X to X .

2022-06-03

Searching for Holes in the Matrix Universe

└ Part III: Homotopy

└ A Second Fundamental Group

└ The Tracial Fundamental Group

- This is our second fundamental group—entirely goverened by the analytic continuation of global germs.
- Since tr eq captures fixed end pt homotopy and the direct sum identity, π_1^{tr} is a quotient of π_1
- There is still a big downside to this definition—it gets us no closer to computation.

The Tracial Fundamental Group

Let $D \subset \mathcal{M}^g$ be an anchored space with B is anchor. For $X \in B$ define $\pi_1^{\text{tr}}(D)$ to be the group of trace equivalent paths essentially taking X to X .

Computationally, we are still stuck.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part III: Homotopy

└ A Second Fundamental Group

└ The Tracial Fundamental Group

- This is our second fundamental group—entirely goverened by the analytic continuation of global germs.
- Since tr eq captures fixed end pt homotopy and the direct sum identity, π_1^{tr} is a quotient of π_1
- There is still a big downside to this definition—it gets us no closer to computation.

Part IV: Cohomology

- Cohomology is going to help us a lot when it comes to characterizing—and eventually computing π_1^{tr} .
- Just a note, this is by far the most technical section—so bear with me.
- before we jump into the particular cohomological theory we are going to be using, lets speedrun a review of cohomology.

- In traditional homology, the boundary homomorphisms *decrease the index*.
- For cohomology, the boundary morphism go the other way—the index *increases*
- Generally, we consider the chain groups to be groups of functions into some abelian group, but that isn't always the case
- once you have this co-chain complex, you compute the cohomology groups exactly the same way you did before.
- the **kernel** of one map mod the **image** of the previous one.

Traditional homology considers a complex of the form

$$\cdots \xleftarrow{\partial_{n-1}} C_{n-1} \xleftarrow{\partial_n} C_n \xleftarrow{\partial_{n+1}} C_{n+1} \xleftarrow{\partial_{n+2}} \cdots$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ A Short Review of Cohomology

Traditional homology considers a complex of the form

$$\cdots \xrightarrow{\partial_{n-1}} C_{n-1} \xrightarrow{\partial_n} C_n \xrightarrow{\partial_{n+1}} C_{n+1} \xrightarrow{\partial_{n+2}} \cdots$$

- In traditional homology, the boundary homomorphisms *decrease the index*.
- For cohomology, the boundary morphism go the other way—the index *increases*
- Generally, we consider the chain groups to be groups of functions into some abelian group, but that isn't always the case
- once you have this co-chain complex, you compute the cohomology groups exactly the same way you did before.
- the **kernel** of one map mod the **image** of the previous one.

Traditional homology considers a complex of the form

$$\cdots \xleftarrow{\partial_{n-1}} C_{n-1} \xleftarrow{\partial_n} C_n \xleftarrow{\partial_{n+1}} C_{n+1} \xleftarrow{\partial_{n+2}} \cdots$$

While cohomology considers a dual complex

$$\cdots \xrightarrow{d_{n-2}} C^{n-1} \xrightarrow{d_{n-1}} C^n \xrightarrow{d_n} C^{n+1} \xrightarrow{d_{n+1}} \cdots$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ A Short Review of Cohomology

Traditional homology considers a complex of the form

$$\cdots \xleftarrow{\partial_{n-1}} C_{n-1} \xleftarrow{\partial_n} C_n \xleftarrow{\partial_{n+1}} C_{n+1} \xleftarrow{\partial_{n+2}} \cdots$$

While cohomology considers a dual complex

$$\cdots \xrightarrow{d_{n-2}} C^{n-1} \xrightarrow{d_{n-1}} C^n \xrightarrow{d_n} C^{n+1} \xrightarrow{d_{n+1}} \cdots$$

- In traditional homology, the boundary homomorphisms *decrease the index*.
- For cohomology, the boundary morphism go the other way—the index *increases*
- Generally, we consider the chain groups to be groups of functions into some abelian group, but that isn't always the case
- once you have this co-chain complex, you compute the cohomology groups exactly the same way you did before.
- the **kernel** of one map mod the **image** of the previous one.

Traditional homology considers a complex of the form

$$\cdots \xleftarrow{\partial_{n-1}} C_{n-1} \xleftarrow{\partial_n} C_n \xleftarrow{\partial_{n+1}} C_{n+1} \xleftarrow{\partial_{n+2}} \cdots$$

While cohomology considers a dual complex

$$\cdots \xrightarrow{d_{n-2}} C^{n-1} \xrightarrow{d_{n-1}} C^n \xrightarrow{d_n} C^{n+1} \xrightarrow{d_{n+1}} \cdots$$

In general, C^k is a group of functions into some abelian group.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ A Short Review of Cohomology

Traditional homology considers a complex of the form

$$\cdots \xleftarrow{\partial_{n-1}} C_{n-1} \xleftarrow{\partial_n} C_n \xleftarrow{\partial_{n+1}} C_{n+1} \xleftarrow{\partial_{n+2}} \cdots$$

While cohomology considers a dual complex

$$\cdots \xrightarrow{d_{n-2}} C^{n-1} \xrightarrow{d_{n-1}} C^n \xrightarrow{d_n} C^{n+1} \xrightarrow{d_{n+1}} \cdots$$

In general, C^k is a group of functions into some abelian group.

- In traditional homology, the boundary homomorphisms *decrease the index*.
- For cohomology, the boundary morphism go the other way—the index *increases*
- Generally, we consider the chain groups to be groups of functions into some abelian group, but that isn't always the case
- once you have this co-chain complex, you compute the cohomology groups exactly the same way you did before.
- the **kernel** of one map mod the **image** of the previous one.

Traditional homology considers a complex of the form

$$\cdots \xleftarrow{\partial_{n-1}} C_{n-1} \xleftarrow{\partial_n} C_n \xleftarrow{\partial_{n+1}} C_{n+1} \xleftarrow{\partial_{n+2}} \cdots$$

While cohomology considers a dual complex

$$\cdots \xrightarrow{d_{n-2}} C^{n-1} \xrightarrow{d_{n-1}} C^n \xrightarrow{d_n} C^{n+1} \xrightarrow{d_{n+1}} \cdots$$

In general, C^k is a group of functions into some abelian group.

The k -th cohomology group is

$$H^k = \frac{\ker d_i}{\operatorname{Im} d_{i-1}}$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ A Short Review of Cohomology

- In traditional homology, the boundary homomorphisms *decrease the index*.
- For cohomology, the boundary morphism go the other way—the index *increases*
- Generally, we consider the chain groups to be groups of functions into some abelian group, but that isn't always the case
- once you have this co-chain complex, you compute the cohomology groups exactly the same way you did before.
- the **kernel** of one map mod the **image** of the previous one.

Traditional homology considers a complex of the form

$$\cdots \xrightarrow{\partial_{n-1}} C_{n-1} \xrightarrow{\partial_n} C_n \xrightarrow{\partial_{n+1}} C_{n+1} \xrightarrow{\partial_{n+2}} \cdots$$

While cohomology considers a dual complex

$$\cdots \xrightarrow{d_{n-2}} C^{n-1} \xrightarrow{d_{n-1}} C^n \xrightarrow{d_n} C^{n+1} \xrightarrow{d_{n+1}} \cdots$$

In general, C^k is a group of functions into some abelian group.The k -th cohomology group is

$$H^k = \frac{\ker d_i}{\operatorname{Im} d_{i-1}}$$

Let D be an anchored set. Denote the set of (globally defined) tracial functions on D by $\mathcal{T}(D)$ and the set of free functions by $\mathcal{F}(D)$.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Tracial Cohomology

Let D be an anchored set. Denote the set of (globally defined) tracial functions on D by $\mathcal{T}(D)$ and the set of free functions by $\mathcal{F}(D)$.

- Lets build a complex! We start with two sets of functions.
- Since the gradient of a **tracial** function is a free one, we have the first map!
- Now this isn't particularly impressive, it is (almost) enough if we are careful.

Let D be an anchored set. Denote the set of (globally defined) tracial functions on D by $\mathcal{T}(D)$ and the set of free functions by $\mathcal{F}(D)$. For $f \in \mathcal{T}(D)$, $\nabla f \in \mathcal{F}(D)$ —so we have the beginnings of a chain complex!

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Tracial Cohomology

Let D be an anchored set. Denote the set of (globally defined) tracial functions on D by $\mathcal{T}(D)$ and the set of free functions by $\mathcal{F}(D)$. For $f \in \mathcal{T}(D)$, $\nabla f \in \mathcal{F}(D)$ —so we have the beginnings of a chain complex!

- Lets build a complex! We start with two sets of functions.
- Since the gradient of a **tracial** function is a free one, we have the first map!
- Now this isn't particularly impressive, it is (almost) enough if we are careful.

Let D be an anchored set. Denote the set of (globally defined) tracial functions on D by $\mathcal{T}(D)$ and the set of free functions by $\mathcal{F}(D)$. For $f \in \mathcal{T}(D)$, $\nabla f \in \mathcal{F}(D)$ —so we have the beginnings of a chain complex!

$$0 \rightarrow \mathcal{T}(D) \xrightarrow{\nabla} \mathcal{F}(D) \rightarrow \cdots .$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Tracial Cohomology

Let D be an anchored set. Denote the set of (globally defined) tracial functions on D by $\mathcal{T}(D)$ and the set of free functions by $\mathcal{F}(D)$. For $f \in \mathcal{T}(D)$, $\nabla f \in \mathcal{F}(D)$ —so we have the beginnings of a chain complex!

$$0 \rightarrow \mathcal{T}(D) \xrightarrow{\nabla} \mathcal{F}(D) \rightarrow \cdots .$$

- Lets build a complex! We start with two sets of functions.
- Since the gradient of a **tracial** function is a free one, we have the first map!
- Now this isn't particularly impressive, it is (almost) enough if we are careful.

$$0 \rightarrow \mathcal{T}(D) \xrightarrow{\nabla} \mathcal{F}(D) \rightarrow \dots .$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Tracial Cohomology

$$0 \rightarrow \mathcal{T}(D) \xrightarrow{\nabla} \mathcal{F}(D) \rightarrow \dots .$$

- Since we are looking at π_1 , we need to create H^1 —which is the homology of *this* part.
- The image of ∇ is very easy—a free function is exact if it there is some *global* tracial function such that $\nabla f = g$
- The kernel is a bit harder—we say closed if it follows this condition.
- if you view $\text{tr}(\cdot)$ as the inner product and $Dg(X)[H]$ as something like the jacobian evaluated on a direction, then this is just the classical condition of a closed!
- Now we define the first *tracial* fundamental group as the closed functions mod the exact ones

$$0 \rightarrow \mathcal{T}(D) \xrightarrow{\nabla} \mathcal{F}(D) \rightarrow \dots$$

A free function $g : D \rightarrow \mathcal{M}^g$ is **exact** if there exists a tracial function $f : D \rightarrow \mathbb{C}$ such that $\nabla f = g$.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Tracial Cohomology

$$0 \rightarrow \mathcal{T}(D) \xrightarrow{\nabla} \mathcal{F}(D) \rightarrow \dots$$

A free function $g : D \rightarrow \mathcal{M}^g$ is **exact** if there exists a tracial function $f : D \rightarrow \mathbb{C}$ such that $\nabla f = g$.

- Since we are looking at π_1 , we need to create H^1 —which is the homology of *this* part.
- The image of ∇ is very easy—a free function is exact if it there is some *global* tracial function such that $\nabla f = g$
- The kernel is a bit harder—we say closed if it follows this condition.
- if you view $\text{tr}(\cdot)$ as the inner product and $Dg(X)[H]$ as something like the jacobian evaluated on a direction, then this is just the classical condition of a closed!
- Now we define the first *tracial* fundamental group as the closed functions mod the exact ones

$$0 \rightarrow \mathcal{T}(D) \xrightarrow{\nabla} \mathcal{F}(D) \rightarrow \dots$$

A free function $g : D \rightarrow \mathcal{M}^g$ is **exact** if there exists a tracial function $f : D \rightarrow \mathbb{C}$ such that $\nabla f = g$.

A free function $g : D \rightarrow \mathcal{M}^g$ is **closed** if

$$\text{tr}(K \cdot Dg(X)[H]) = \text{tr}(H \cdot Dg(X)[K])$$

for all directions H, K .

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Tracial Cohomology

- Since we are looking at π_1 , we need to create H^1 —which is the homology of *this* part.
- The image of ∇ is very easy—a free function is exact if there is some *global* tracial function such that $\nabla f = g$
- The kernel is a bit harder—we say closed if it follows this condition.
- if you view $\text{tr}(\cdot)$ as the inner product and $Dg(X)[H]$ as something like the jacobian evaluated on a direction, then this is just the classical condition of a closed!
- Now we define the first *tracial* fundamental group as the closed functions mod the exact ones

$$0 \rightarrow \mathcal{T}(D) \xrightarrow{\nabla} \mathcal{F}(D) \rightarrow \dots$$

A free function $g : D \rightarrow \mathcal{M}^g$ is **exact** if there exists a tracial function $f : D \rightarrow \mathbb{C}$ such that $\nabla f = g$.

A free function $g : D \rightarrow \mathcal{M}^g$ is **closed** if

$$\text{tr}(K \cdot Dg(X)[H]) = \text{tr}(H \cdot Dg(X)[K])$$

for all directions H, K .

$$0 \rightarrow \mathcal{T}(D) \xrightarrow{\nabla} \mathcal{F}(D) \rightarrow \dots$$

A free function $g : D \rightarrow \mathcal{M}^{\theta}$ is **exact** if there exists a tracial function $f : D \rightarrow \mathbb{C}$ such that $\nabla f = g$.

A free function $g : D \rightarrow \mathcal{M}^{\theta}$ is **closed** if

$$\text{tr}(K \cdot Dg(X)[H]) = \text{tr}(H \cdot Dg(X)[K])$$

for all directions H, K .

Definition:

The **first tracial cohomology group** is the vector space of closed free functions modulo the exact free function. We write $H_{\text{tr}}^1(D)$.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Tracial Cohomology

$$0 \rightarrow \mathcal{T}(D) \xrightarrow{\nabla} \mathcal{F}(D) \rightarrow \dots$$

A free function $g : D \rightarrow \mathcal{M}^{\theta}$ is **exact** if there exists a tracial function $f : D \rightarrow \mathbb{C}$ such that $\nabla f = g$.

A free function $g : D \rightarrow \mathcal{M}^{\theta}$ is **closed** if

$$\text{tr}(K \cdot Dg(X)[H]) = \text{tr}(H \cdot Dg(X)[K])$$

for all directions H, K .

Definition:

The **first tracial cohomology group** is the vector space of closed free functions modulo the exact free function. We write $H_{\text{tr}}^1(D)$.

- Since we are looking at π_1 , we need to create H^1 —which is the homology of *this* part.
- The image of ∇ is very easy—a free function is exact if there is some *global* tracial function such that $\nabla f = g$
- The kernel is a bit harder—we say closed if it follows this condition.
- if you view $\text{tr}(\cdot)$ as the inner product and $Dg(X)[H]$ as something like the jacobian evaluated on a direction, then this is just the classical condition of a closed!
- Now we define the first *tracial* fundamental group as the closed functions mod the exact ones

What about global germs?

- For $f : B \rightarrow \mathbb{C}$ a global germ, since f analytically continues along every path, so does ∇f .

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Tracial Cohomology

└ What about global germs?

- Now I want to pause for a second and think about the global germs that govern π_1^{tr} —where do they fit in with everything?
- Since f continues along every path, so does ∇f .
- By free monodromy then, ∇f has a unique extension to the whole space—it is a global free function.
- But! ∇f is **not** necessarily exact bc f doesn't necessarily have a global extension.

What about global germs?

- For $f : B \rightarrow \mathbb{C}$ a global germ, since f analytically continues along every path, so does ∇f .
- Free Monodromy means that ∇f has a global extension.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Tracial Cohomology

└ What about global germs?

What about global germs?

- For $f : B \rightarrow \mathbb{C}$ a global germ, since f analytically continues along every path, so does ∇f .
- Free Monodromy means that ∇f has a global extension.

- Now I want to pause for a second and think about the global germs that govern π_1^{tr} —where do they fit in with everything?
- Since f continues along every path, so does ∇f .
- By free monodromy then, ∇f has a unique extension to the whole space—it is a global free function.
- But! ∇f is *not* necessarily exact bc f doesn't necessarily have a global extension.

What about global germs?

- For $f : B \rightarrow \mathbb{C}$ a global germ, since f analytically continues along every path, so does ∇f .
- Free Monodromy means that ∇f has a global extension.
- **Important:** If f is a global germ, then ∇f is not necessarily exact since $\mathcal{T}(D)$ is the set of tracial functions defined on *all of* D .

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Tracial Cohomology

└ What about global germs?

- Now I want to pause for a second and think about the global germs that govern π_1^{tr} —where do they fit in with everything?
- Since f continues along every path, so does ∇f .
- By free monodromy then, ∇f has a unique extension to the whole space—it is a global free function.
- But! ∇f is **not** necessarily exact bc f doesn't necessarily have a global extension.

- For $f : B \rightarrow \mathbb{C}$ a global germ, since f analytically continues along every path, so does ∇f .
- Free Monodromy means that ∇f has a global extension.
- **Important:** If f is a global germ, then ∇f is not necessarily exact since $\mathcal{T}(D)$ is the set of tracial functions defined on *all of* D .

Goal: Show that $\pi_1^{\text{tr}}(D)$ injects into \mathbb{C} .

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Injecting into \mathbb{C} Goal: Show that $\pi_1^{\text{tr}}(D)$ injects into \mathbb{C} .

- Before we continue, I want to give a look at the light at the end of the tunnel. We are going to use the tracial cohomology group to show that π_1^{tr} injects in \mathbb{C} and prove a major structure theorem.
- Like any good theorem, it all depends on some technical lemma—here is ours
- So what is this saying? $f(\alpha\beta) - f(\alpha)$ measure how analytic continuation changes the value of $f(\alpha)$. With this in mind, it isn't hard to believe the lemma.

Goal: Show that $\pi_1^{\text{tr}}(D)$ injects into \mathbb{C} .

Lemma (Kerbs)

Let D be an anchored set. For any $\alpha, \beta \in \pi_1^{\text{tr}}(D)$ and global germ f ,

$$f(\alpha\beta) - f(\alpha) = f(\beta) - f(\tau)$$

where τ is the constant path.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Injecting into \mathbb{C}

Goal: Show that $\pi_1^{\text{tr}}(D)$ injects into \mathbb{C} .

Lemma (Kerbs)

Let D be an anchored set. For any $\alpha, \beta \in \pi_1^{\text{tr}}(D)$ and global germ f ,

$$f(\alpha\beta) - f(\alpha) = f(\beta) - f(\tau)$$

where τ is the constant path.

- Before we continue, I want to give a look at the light at the end of the tunnel. We are going to use the tracial cohomology group to show that π_1^{tr} injects in \mathbb{C} and prove a major structure theorem.
- Like any good theorem, it all depends on some technical lemma—here is ours
- So what is this saying? $f(\alpha\beta) - f(\alpha)$ measure how analytic continuation changes the value of $f(\alpha)$. With this in mind, it isn't hard to believe the lemma.

For D an anchored set, $X \in B_1$ the base point, f a global germ, and $\gamma \in \pi_1^{\text{tr}}(D)$, define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Injecting into \mathbb{C}

For D an anchored set, $X \in B_1$ the base point, f a global germ, and $\gamma \in \pi_1^{\text{tr}}(D)$, define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

- Lets define $c^f(\gamma)$ to be the measure of how analytic continuation along γ changes f .
- What is amazing is that $c^f(\gamma)$ is a homomorphism into \mathbb{C}
- From the definition, we use the ol’ “add, subtract” trick and then our technical lemma
- In fact, $c^f(\gamma)$ gives us something stronger than a homomorphism into \mathbb{C} .

For D an anchored set, $X \in B_1$ the base point, f a global germ, and $\gamma \in \pi_1^{\text{tr}}(D)$, define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

c^f gives us a homomorphism into \mathbb{C} !

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Injecting into \mathbb{C}

For D an anchored set, $X \in B_1$ the base point, f a global germ, and $\gamma \in \pi_1^{\text{tr}}(D)$, define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

c^f gives us a homomorphism into \mathbb{C} !

- Lets define $c^f(\gamma)$ to be the measure of how analytic continuation along γ changes f .
- What is amazing is that $c^f(\gamma)$ is a homomorphism into \mathbb{C}
- From the definition, we use the ol’ “add, subtract” trick and then our technical lemma
- In fact, $c^f(\gamma)$ gives us something stronger than a homomorphism into \mathbb{C} .

For D an anchored set, $X \in B_1$ the base point, f a global germ, and $\gamma \in \pi_1^{\text{tr}}(D)$, define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

c^f gives us a homomorphism into \mathbb{C} !

$$c^f(\gamma_1\gamma_2) = f(\gamma_1\gamma_2) - f(\tau)$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Injecting into \mathbb{C}

For D an anchored set, $X \in B_1$ the base point, f a global germ, and $\gamma \in \pi_1^{\text{tr}}(D)$, define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

c^f gives us a homomorphism into \mathbb{C} !

$$c^f(\gamma_1\gamma_2) = f(\gamma_1\gamma_2) - f(\tau)$$

- Lets define $c^f(\gamma)$ to be the measure of how analytic continuation along γ changes f .
- What is amazing is that $c^f(\gamma)$ is a homomorphism into \mathbb{C}
- From the definition, we use the ol’ “add, subtract” trick and then our technical lemma
- In fact, $c^f(\gamma)$ gives us something stronger than a homomorphism into \mathbb{C} .

For D an anchored set, $X \in B_1$ the base point, f a global germ, and $\gamma \in \pi_1^{\text{tr}}(D)$, define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

c^f gives us a homomorphism into \mathbb{C} !

$$\begin{aligned} c^f(\gamma_1\gamma_2) &= f(\gamma_1\gamma_2) - f(\tau) \\ &= f(\gamma_1\gamma_2) - f(\gamma_1) + f(\gamma_1) - f(\tau) \end{aligned}$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Injecting into \mathbb{C}

For D an anchored set, $X \in B_1$ the base point, f a global germ, and $\gamma \in \pi_1^{\text{tr}}(D)$, define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

c^f gives us a homomorphism into \mathbb{C} !

$$\begin{aligned} c^f(\gamma_1\gamma_2) &= f(\gamma_1\gamma_2) - f(\tau) \\ &= f(\gamma_1\gamma_2) - f(\gamma_1) + f(\gamma_1) - f(\tau) \end{aligned}$$

- Lets define $c^f(\gamma)$ to be the measure of how analytic continuation along γ changes f .
- What is amazing is that $c^f(\gamma)$ is a homomorphism into \mathbb{C}
- From the definition, we use the ol' “add, subtract” trick and then our technical lemma
- In fact, $c^f(\gamma)$ gives us something stronger than a homomorphism into \mathbb{C} .

For D an anchored set, $X \in B_1$ the base point, f a global germ, and $\gamma \in \pi_1^{\text{tr}}(D)$, define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

c^f gives us a homomorphism into \mathbb{C} !

$$\begin{aligned} c^f(\gamma_1\gamma_2) &= f(\gamma_1\gamma_2) - f(\tau) \\ &= f(\gamma_1\gamma_2) - f(\gamma_1) + f(\gamma_1) - f(\tau) \\ &= f(\gamma_2) - f(\tau) + f(\gamma_1) - f(\tau) \end{aligned}$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Injecting into \mathbb{C}

For D an anchored set, $X \in B_1$ the base point, f a global germ, and $\gamma \in \pi_1^{\text{tr}}(D)$, define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

c^f gives us a homomorphism into \mathbb{C} !

$$\begin{aligned} c^f(\gamma_1\gamma_2) &= f(\gamma_1\gamma_2) - f(\tau) \\ &= f(\gamma_1\gamma_2) - f(\gamma_1) + f(\gamma_1) - f(\tau) \\ &= f(\gamma_2) - f(\tau) + f(\gamma_1) - f(\tau) \end{aligned}$$

- Lets define $c^f(\gamma)$ to be the measure of how analytic continuation along γ changes f .
- What is amazing is that $c^f(\gamma)$ is a homomorphism into \mathbb{C}
- From the definition, we use the ol' “add, subtract” trick and then our technical lemma
- In fact, $c^f(\gamma)$ gives us something stronger than a homomorphism into \mathbb{C} .

For D an anchored set, $X \in B_1$ the base point, f a global germ, and $\gamma \in \pi_1^{\text{tr}}(D)$, define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

c^f gives us a homomorphism into \mathbb{C} !

$$\begin{aligned} c^f(\gamma_1\gamma_2) &= f(\gamma_1\gamma_2) - f(\tau) \\ &= f(\gamma_1\gamma_2) - f(\gamma_1) + f(\gamma_1) - f(\tau) \\ &= f(\gamma_2) - f(\tau) + f(\gamma_1) - f(\tau) \\ &= c^f(\gamma_2) + c^f(\gamma_1). \end{aligned}$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Injecting into \mathbb{C}

For D an anchored set, $X \in B_1$ the base point, f a global germ, and $\gamma \in \pi_1^{\text{tr}}(D)$, define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

c^f gives us a homomorphism into \mathbb{C} !

$$\begin{aligned} c^f(\gamma_1\gamma_2) &= f(\gamma_1\gamma_2) - f(\tau) \\ &= f(\gamma_1\gamma_2) - f(\gamma_1) + f(\gamma_1) - f(\tau) \\ &= f(\gamma_2) - f(\tau) + f(\gamma_1) - f(\tau) \\ &= c^f(\gamma_2) + c^f(\gamma_1). \end{aligned}$$

- Lets define $c^f(\gamma)$ to be the measure of how analytic continuation along γ changes f .
- What is amazing is that $c^f(\gamma)$ is a homomorphism into \mathbb{C}
- From the definition, we use the ol' “add, subtract” trick and then our technical lemma
- In fact, $c^f(\gamma)$ gives us something stronger than a homomorphism into \mathbb{C} .

Lemma (Pascoe, 2020)

The map

$$\Phi : \pi_1^{\text{tr}}(D) \longrightarrow \prod_{\substack{\nabla f \in H_{\text{tr}}^1(D) \\ f \text{ a global germ}}} \mathbb{C}$$

$$\gamma \longmapsto \prod c^f(\gamma)$$

is an injective homomorphism.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Injecting into \mathbb{C}

Lemma (Pascoe, 2020)

The map

$$\Phi : \pi_1^{\text{tr}}(D) \longrightarrow \prod_{\substack{\nabla f \in H_{\text{tr}}^1(D) \\ f \text{ a global germ}}} \mathbb{C}$$

$$\gamma \longmapsto \prod c^f(\gamma)$$

is an injective homomorphism.

- So this lemma claims to be an injective homomorphism, but what it going on with the map?
- We map π_1 into a product of \mathbb{C} 's with one for every unique image of a global germ under ∇ .
- To get injectivity, you do need that many products of \mathbb{C} . You end up using a very similar arguement to what we did before of “add, subtract” then using the technical lemma.
- This gives us two really important characterizations of π_1^{tr} —we are commutative *and* torsion free

Lemma (Pascoe, 2020)

The map

$$\Phi : \pi_1^{\text{tr}}(D) \longrightarrow \prod_{\substack{\nabla f \in H_{\text{tr}}^1(D) \\ f \text{ a global germ}}} \mathbb{C}$$

$$\gamma \longmapsto \prod c^f(\gamma)$$

*is an injective homomorphism.*So $\pi_1^{\text{tr}}(D)$ is commutative and torsion free!

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Injecting into \mathbb{C}

Lemma (Pascoe, 2020)

The map

$$\Phi : \pi_1^{\text{tr}}(D) \longrightarrow \prod_{\substack{\nabla f \in H_{\text{tr}}^1(D) \\ f \text{ a global germ}}} \mathbb{C}$$

$$\gamma \longmapsto \prod c^f(\gamma)$$

*is an injective homomorphism.*So $\pi_1^{\text{tr}}(D)$ is commutative and torsion free!

- So this lemma claims to be an injective homomorphism, but what it going on with the map?
- We map π_1 into a product of \mathbb{C} 's with one for every unique image of a global germ under ∇ .
- To get injectivity, you do need that many products of \mathbb{C} . You end up using a very similar arguement to what we did before of “add, subtract” then using the technical lemma.
- This gives us two really important characterizations of π_1^{tr} —we are commutative *and* torsion free

$\pi_1^{\text{tr}}(D)$ is divisible

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Characterizing $\pi_1^{\text{tr}}(D)$ └ $\pi_1^{\text{tr}}(D)$ is divisible

- But we can go further, and show that π_1^{tr} is divisible. Since we are writing our group multiplicatively, this is equivalent to saying we can take n -th roots
- The first thing we need is that, in π_1 you can direct sum a τ on either side. All you need for this is a fixed end pt homotopy of some rotation matrices
- so if we have a path on one side of a direct sum and an identity on the other, we are free to flip the order—this is how we will show divisibility

$\pi_1^{\text{tr}}(D)$ is divisible

For any $\gamma \in \pi_1^{\text{tr}}(D)$,

$$\gamma \oplus \tau = \tau \oplus \gamma.$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Characterizing $\pi_1^{\text{tr}}(D)$ └ $\pi_1^{\text{tr}}(D)$ is divisible

- But we can go further, and show that π_1^{tr} is divisible. Since we are writing our group multiplicatively, this is equivalent to saying we can take n -th roots
- The first thing we need is that, in π_1 you can direct sum a τ on either side. All you need for this is a fixed end pt homotopy of some rotation matrices
- so if we have a path on one side of a direct sum and an identity on the other, we are free to flip the order—this is how we will show divisibility

$\pi_1^{\text{tr}}(D)$ is divisible

For any $\gamma \in \pi_1^{\text{tr}}(D)$,

$$\gamma \oplus \tau = \tau \oplus \gamma.$$

Why?

$$H(t, \theta) = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} (\gamma \oplus \tau) \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}^*$$

is a homotopy between the paths.

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Characterizing $\pi_1^{\text{tr}}(D)$ └ $\pi_1^{\text{tr}}(D)$ is divisible

- But we can go further, and show that π_1^{tr} is divisible. Since we are writing our group multiplicatively, this is equivalent to saying we can take n -th roots
- The first thing we need is that, in π_1 you can direct sum a τ on either side. All you need for this is a fixed end pt homotopy of some rotation matrices
- so if we have a path on one side of a direct sum and an identity on the other, we are free to flip the order—this is how we will show divisibility

 $\pi_1^{\text{tr}}(D)$ is divisibleFor any $\gamma \in \pi_1^{\text{tr}}(D)$,

$$\gamma \oplus \tau = \tau \oplus \gamma.$$

Why?

$$H(t, \theta) = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} (\gamma \oplus \tau) \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}^*$$

is a homotopy between the paths.

$$\gamma = \underbrace{\begin{bmatrix} \gamma & & \\ & \gamma & \\ & & \ddots \\ & & & \gamma \end{bmatrix}}_{k\text{-times}}$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Characterizing $\pi_1^{\text{tr}}(D)$

$$\gamma = \begin{bmatrix} \gamma & & \\ & \ddots & \\ & & \gamma \end{bmatrix}_{k\text{-times}}$$

- Under trace equivalence, we can direct sum as many γ 's we want, so we grab k of them
- Since diagonal, we can separate this to be a single γ and a whole lot of τ 's
- But now we can go through and flip these one by one to put the γ first
- If we collect terms, we get a formula for k -th roots!

$$\gamma = \underbrace{\begin{bmatrix} \gamma & & \\ & \gamma & \\ & & \ddots \\ & & & \gamma \end{bmatrix}}_{k\text{-times}}$$

$$= \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix} \begin{bmatrix} \tau & & \\ & \gamma & \\ & & \ddots \\ & & & \tau \end{bmatrix} \cdots \begin{bmatrix} \tau & & \\ & \tau & \\ & & \ddots \\ & & & \gamma \end{bmatrix}$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Characterizing $\pi_1^{\text{tr}}(D)$

$$\gamma - \underbrace{\begin{bmatrix} \gamma & \\ & \gamma \end{bmatrix}}_{k\text{-times}}$$

$$= \begin{bmatrix} \gamma & \\ & \tau \end{bmatrix} \begin{bmatrix} \tau & \\ & \gamma \end{bmatrix} - \begin{bmatrix} \gamma & \\ & \gamma \end{bmatrix}$$

- Under trace equivalence, we can direct sum as many γ 's we want, so we grab k of them
- Since diagonal, we can separate this to be a single γ and a whole lot of τ 's
- But now we can go through and flip these one by one to put the γ first
- If we collect terms, we get a formula for k -th roots!

$$\begin{aligned}
 \gamma &= \underbrace{\begin{bmatrix} \gamma & & \\ & \gamma & \\ & & \ddots \\ & & & \gamma \end{bmatrix}}_{k\text{-times}} \\
 &= \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix} \begin{bmatrix} \tau & & \\ & \gamma & \\ & & \ddots \\ & & & \tau \end{bmatrix} \cdots \begin{bmatrix} \tau & & \\ & \tau & \\ & & \ddots \\ & & & \gamma \end{bmatrix} \\
 &= \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix} \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix} \cdots \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix}
 \end{aligned}$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Characterizing $\pi_1^{\text{tr}}(D)$

$$\gamma - \underbrace{\begin{bmatrix} \gamma & \\ & \ddots \\ & & \gamma \end{bmatrix}}_{k\text{-times}}$$

- Under trace equivalence, we can direct sum as many γ 's we want, so we grab k of them
- Since diagonal, we can separate this to be a single γ and a whole lot of τ 's
- But now we can go through and flip these one by one to put the γ first
- If we collect terms, we get a formula for k -th roots!

$$\begin{aligned}
 \gamma &= \underbrace{\begin{bmatrix} \gamma & & \\ & \gamma & \\ & & \ddots \\ & & & \gamma \end{bmatrix}}_{k\text{-times}} \\
 &= \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix} \begin{bmatrix} \tau & & \\ & \gamma & \\ & & \ddots \\ & & & \tau \end{bmatrix} \cdots \begin{bmatrix} \tau & & \\ & \tau & \\ & & \ddots \\ & & & \gamma \end{bmatrix} \\
 &= \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix} \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix} \cdots \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix} \\
 &= \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix}^k
 \end{aligned}$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Characterizing $\pi_1^{\text{tr}}(D)$

$$\gamma = \underbrace{\begin{bmatrix} \gamma & & \\ & \gamma & \\ & & \ddots \\ & & & \gamma \end{bmatrix}}_{k\text{-times}} \begin{bmatrix} \tau & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix} \cdots \begin{bmatrix} \tau & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix}$$

- Under trace equivalence, we can direct sum as many γ 's we want, so we grab k of them
- Since diagonal, we can separate this to be a single γ and a whole lot of τ 's
- But now we can go through and flip these one by one to put the γ first
- If we collect terms, we get a formula for k -th roots!

Theorem (Pascoe 2020)

For D an anchored free set, $\pi_1^{\text{tr}}(D)$ is a torsion free, abelian, divisible group. That is,

$$\pi_1^{\text{tr}}(D) \simeq \bigoplus_{i \in I} \mathbb{Q} = \mathbb{Q}^I$$

for some set I .

2022-06-03

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Characterizing $\pi_1^{\text{tr}}(D)$

Theorem (Pascoe 2020)

For D an anchored free set, $\pi_1^{\text{tr}}(D)$ is a torsion free, abelian, divisible group. That is,

$$\pi_1^{\text{tr}}(D) \simeq \bigoplus_{i \in I} \mathbb{Q} = \mathbb{Q}^I$$

for some set I .

- Putting this all together, we know that π_1^{tr} is abelian, divisible, and torsion free. Thanks to a “fundamental structure theorem” this means that π_1^{tr} is isomorphic to some number of copies of \mathbb{Q} !
- “But I promised we would be able to compute! This is just a structure theorem!”

Part V: Computing $\pi_1^{\text{tr}}(D)$

- In classical theory, we have VC and MV—neither of these exist here. But we have something better: universal properties.

Let D be an anchored, path connected set such that each D_n is nonempty and choose an anchor $B \subset D$ such that each B_n is also nonempty.

2022-06-03

Searching for Holes in the Matrix Universe

└ Computing $\pi_1^{\text{tr}}(D)$

└ The Direct Limit Approach

Let D be an anchored, path connected set such that each D_n is nonempty and choose an anchor $B \subset D$ such that each B_n is also nonempty.

- Here are our niceness condition for our computation to work.
- as before, the $_n$ denotes restricting to the $n \times n$ level.
- We are a quotient bc these are all paths in D_n but there might be some paths that are distinct in $\pi_1(D_n)$ but not in $\pi_1^{\text{tr}}(D)_n$.
- Via direct sums, we have inclusion into any level with contains n as a factor

Let D be an anchored, path connected set such that each D_n is nonempty and choose an anchor $B \subset D$ such that each B_n is also nonempty.

Let $\pi_1^{\text{tr}}(D)_n$ is the subgroup of paths in D_n . Note that $\pi_1^{\text{tr}}(D)_n$ is a quotient of $\pi_1(D_n)$.

2022-06-03

Searching for Holes in the Matrix Universe

└ Computing $\pi_1^{\text{tr}}(D)$

└ The Direct Limit Approach

- Here are our niceness condition for our computation to work.
- as before, the $_n$ denotes restricting to the $n \times n$ level.
- We are a quotient bc these are all paths in D_n but there might be some paths that are distinct in $\pi_1(D_n)$ but not in $\pi_1^{\text{tr}}(D)_n$.
- Via direct sums, we have inclusion into any level with contains n as a factor

Let D be an anchored, path connected set such that each D_n is nonempty and choose an anchor $B \subset D$ such that each B_n is also nonempty.

Let $\pi_1^{\text{tr}}(D)_n$ is the subgroup of paths in D_n . Note that $\pi_1^{\text{tr}}(D)_n$ is a quotient of $\pi_1(D_n)$.

Let D be an anchored, path connected set such that each D_n is nonempty and choose an anchor $B \subset D$ such that each B_n is also nonempty.

Let $\pi_1^{\text{tr}}(D)_n$ is the subgroup of paths in D_n . Note that $\pi_1^{\text{tr}}(D)_n$ is a quotient of $\pi_1(D_n)$.

There is a natural inclusion

$$\begin{aligned} \pi_1^{\text{tr}}(D)_n &\hookrightarrow \pi_1^{\text{tr}}(D)_{kn} \\ \gamma &\longmapsto \gamma^{\oplus k} \end{aligned}$$

for all k .

2022-06-03

Searching for Holes in the Matrix Universe

└ Computing $\pi_1^{\text{tr}}(D)$

└ The Direct Limit Approach

- Here are our niceness condition for our computation to work.
- as before, the $_n$ denotes restricting to the $n \times n$ level.
- We are a quotient bc these are all paths in D_n but there might be some paths that are distinct in $\pi_1(D_n)$ but not in $\pi_1^{\text{tr}}(D)_n$.
- Via direct sums, we have inclusion into any level with contains n as a factor

Let D be an anchored, path connected set such that each D_n is nonempty and choose an anchor $B \subset D$ such that each B_n is also nonempty.

Let $\pi_1^{\text{tr}}(D)_n$ is the subgroup of paths in D_n . Note that $\pi_1^{\text{tr}}(D)_n$ is a quotient of $\pi_1(D_n)$.

There is a natural inclusion

$$\begin{aligned} \pi_1^{\text{tr}}(D)_n &\hookrightarrow \pi_1^{\text{tr}}(D)_{kn} \\ \gamma &\longmapsto \gamma^{\oplus k} \end{aligned}$$

for all k .

Now consider the chain of inclusions:

$$\pi_1^{\text{tr}}(D)_1 \hookrightarrow \pi_1^{\text{tr}}(D)_2 \hookrightarrow \pi_1^{\text{tr}}(D)_6 \hookrightarrow \cdots \hookrightarrow \pi_1^{\text{tr}}(D)_{n!} \hookrightarrow \cdots$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Computing $\pi_1^{\text{tr}}(D)$

└ The Direct Limit Approach

Now consider the chain of inclusions:

$$\pi_1^{\text{tr}}(D)_1 \hookrightarrow \pi_1^{\text{tr}}(D)_2 \hookrightarrow \pi_1^{\text{tr}}(D)_6 \hookrightarrow \cdots \hookrightarrow \pi_1^{\text{tr}}(D)_{n!} \hookrightarrow \cdots$$

- starting at the scalar level, lets run through all of the natural inclusions, muliplying by the next positive integer each time.
- If you're familiar with direct limits, its not hard to see that π_1^{tr} is the direct limit of this sequence
- if you have no idea what that means, just think of it as the group that naturally lives at the “end” of this sequence.
- for domains that if within our niceness condition, this is exactly how we compute π_1^{tr} .

Now consider the chain of inclusions:

$$\pi_1^{\text{tr}}(D)_1 \hookrightarrow \pi_1^{\text{tr}}(D)_2 \hookrightarrow \pi_1^{\text{tr}}(D)_6 \hookrightarrow \dots \hookrightarrow \pi_1^{\text{tr}}(D)_{n!} \hookrightarrow \dots$$

The limit of this sequence isomorphic to $\pi_1^{\text{tr}}(D)!$

2022-06-03

Searching for Holes in the Matrix Universe

└ Computing $\pi_1^{\text{tr}}(D)$

└ The Direct Limit Approach

Now consider the chain of inclusions:
 $\pi_1^{\text{tr}}(D)_1 \hookrightarrow \pi_1^{\text{tr}}(D)_2 \hookrightarrow \pi_1^{\text{tr}}(D)_6 \hookrightarrow \dots \hookrightarrow \pi_1^{\text{tr}}(D)_{n!} \hookrightarrow \dots$
 The limit of this sequence isomorphic to $\pi_1^{\text{tr}}(D)!$

- starting at the scalar level, lets run through all of the natural inclusions, muliplying by the next positive integer each time.
- If you're familiar with direct limits, its not hard to see that π_1^{tr} is the direct limit of this sequence
- if you have no idea what that means, just think of it as the group that naturally lives at the “end” of this sequence.
- for domains that if within our niceness condition, this is exactly how we compute π_1^{tr} .

Example: $\pi_1^{\text{tr}}(GL)$

Let $GL = \bigcup_{n \in \mathbb{N}} GL_n(\mathbb{C})$.

2022-06-03

Searching for Holes in the Matrix Universe

└ Computing $\pi_1^{\text{tr}}(D)$

└ An Example

└ Example: $\pi_1^{\text{tr}}(GL)$ Example: $\pi_1^{\text{tr}}(GL)$ Let $GL = \bigcup_{n \in \mathbb{N}} GL_n(\mathbb{C})$.

- Now we can proceed with an actual example! GL is all invertable matrices of all sizes.
- using the direct limit approach, we want look at the 1x1 slice.
- the determinant a scalar is itself, so our set is $\mathbb{C} \setminus \{0\}$, which has fundamental group \mathbb{Z} . The only way to be at torsion free subgroup/quotient of \mathbb{Z} is just to be \mathbb{Z} .
- When we go the second level, we pick up square roots. It is not too difficult to show that we don't get anything else, so we get $\mathbb{Z}[1/2]$

Example: $\pi_1^{\text{tr}}(GL)$

Let $GL = \bigcup_{n \in \mathbb{N}} GL_n(\mathbb{C})$.

$GL_1(\mathbb{C}) = \mathbb{C} \setminus \{0\}$. Since $\pi_1(GL_1) \simeq \mathbb{Z}$, $\pi_1^{\text{tr}}(GL)_1 \simeq \mathbb{Z}$ as well.

2022-06-03

Searching for Holes in the Matrix Universe

└ Computing $\pi_1^{\text{tr}}(D)$

└ An Example

└ Example: $\pi_1^{\text{tr}}(GL)$ Example: $\pi_1^{\text{tr}}(GL)$ Let $GL = \bigcup_{n \in \mathbb{N}} GL_n(\mathbb{C})$. $GL_1(\mathbb{C}) = \mathbb{C} \setminus \{0\}$. Since $\pi_1(GL_1) \simeq \mathbb{Z}$, $\pi_1^{\text{tr}}(GL)_1 \simeq \mathbb{Z}$ as well.

- Now we can proceed with an actual example! GL is all invertable matrices of all sizes.
- using the direct limit approach, we want look at the 1x1 slice.
- the determinant a scalar is itself, so our set is $\mathbb{C} \setminus \{0\}$, which has fundamental group \mathbb{Z} . The only way to be at torsion free subgroup/quotient of \mathbb{Z} is just to be \mathbb{Z} .
- When we go the second level, we pick up square roots. It is not too difficult to show that we don't get anything else, so we get $\mathbb{Z}[1/2]$

Example: $\pi_1^{\text{tr}}(GL)$

Let $GL = \bigcup_{n \in \mathbb{N}} GL_n(\mathbb{C})$.

$GL_1(\mathbb{C}) = \mathbb{C} \setminus \{0\}$. Since $\pi_1(GL_1) \simeq \mathbb{Z}$, $\pi_1^{\text{tr}}(GL)_1 \simeq \mathbb{Z}$ as well.

Inclusion into $\pi_1^{\text{tr}}(GL)_2$ picks up square roots. If $\gamma \in \pi_1^{\text{tr}}(GL)_1$, then

$$\begin{bmatrix} \gamma & \\ & \tau \end{bmatrix} \in \pi_1^{\text{tr}}(GL)_2.$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Computing $\pi_1^{\text{tr}}(D)$

└ An Example

└ Example: $\pi_1^{\text{tr}}(GL)$

- Now we can proceed with an actual example! GL is all invertable matrices of all sizes.
- using the direct limit approach, we want look at the 1x1 slice.
- the determinant a scalar is itself, so our set is $\mathbb{C} \setminus \{0\}$, which has fundamental group \mathbb{Z} . The only way to be at torsion free subgroup/quotient of \mathbb{Z} is just to be \mathbb{Z} .
- When we go the second level, we pick up square roots. It is not too difficult to show that we don't get anything else, so we get $\mathbb{Z}[1/2]$

Let $GL = \bigcup_{n \in \mathbb{N}} GL_n(\mathbb{C})$.

$GL_1(\mathbb{C}) = \mathbb{C} \setminus \{0\}$. Since $\pi_1(GL_1) \simeq \mathbb{Z}$, $\pi_1^{\text{tr}}(GL)_1 \simeq \mathbb{Z}$ as well.

Inclusion into $\pi_1^{\text{tr}}(GL)_2$ picks up square roots. If $\gamma \in \pi_1^{\text{tr}}(GL)_1$, then

$$\begin{bmatrix} \gamma & \\ & \tau \end{bmatrix} \in \pi_1^{\text{tr}}(GL)_2.$$

Example: $\pi_1^{\text{tr}}(GL)$

Let $GL = \bigcup_{n \in \mathbb{N}} GL_n(\mathbb{C})$.

$GL_1(\mathbb{C}) = \mathbb{C} \setminus \{0\}$. Since $\pi_1(GL_1) \simeq \mathbb{Z}$, $\pi_1^{\text{tr}}(GL)_1 \simeq \mathbb{Z}$ as well.

Inclusion into $\pi_1^{\text{tr}}(GL)_2$ picks up square roots. If $\gamma \in \pi_1^{\text{tr}}(GL)_1$, then

$$\begin{bmatrix} \gamma & \\ & \tau \end{bmatrix} \in \pi_1^{\text{tr}}(GL)_2.$$

Thus, $\pi_1^{\text{tr}}(GL)_2 \simeq \mathbb{Z} \left[\frac{1}{2} \right]$.

2022-06-03

Searching for Holes in the Matrix Universe

└ Computing $\pi_1^{\text{tr}}(D)$

└ An Example

└ Example: $\pi_1^{\text{tr}}(GL)$ Example: $\pi_1^{\text{tr}}(GL)$ Let $GL = \bigcup_{n \in \mathbb{N}} GL_n(\mathbb{C})$. $GL_1(\mathbb{C}) = \mathbb{C} \setminus \{0\}$. Since $\pi_1(GL_1) \simeq \mathbb{Z}$, $\pi_1^{\text{tr}}(GL)_1 \simeq \mathbb{Z}$ as well.Inclusion into $\pi_1^{\text{tr}}(GL)_2$ picks up square roots. If $\gamma \in \pi_1^{\text{tr}}(GL)_1$, then

$$\begin{bmatrix} \gamma & \\ & \tau \end{bmatrix} \in \pi_1^{\text{tr}}(GL)_2.$$

Thus, $\pi_1^{\text{tr}}(GL)_2 \simeq \mathbb{Z} \left[\frac{1}{2} \right]$.

- Now we can proceed with an actual example! GL is all invertable matrices of all sizes.
- using the direct limit approach, we want look at the 1x1 slice.
- the determinant a scalar is itself, so our set is $\mathbb{C} \setminus \{0\}$, which has fundamental group \mathbb{Z} . The only way to be at torsion free subgroup/quotient of \mathbb{Z} is just to be \mathbb{Z} .
- When we go the second level, we pick up square roots. It is not too difficult to show that we don't get anything else, so we get $\mathbb{Z}[1/2]$

Similarly, inclusion into $\pi_1^{\text{tr}}(GL)_{3!}$ picks up cube roots:

$$\pi_1^{\text{tr}}(GL)_6 \simeq \mathbb{Z} \left[\frac{1}{2}, \frac{1}{3} \right]$$

2022-06-03

Searching for Holes in the Matrix Universe

└ Computing $\pi_1^{\text{tr}}(D)$

└ An Example

Similarly, inclusion into $\pi_1^{\text{tr}}(GL)_6$ picks up cube roots:

$$\pi_1^{\text{tr}}(GL)_6 \simeq \mathbb{Z} \left[\frac{1}{2}, \frac{1}{3} \right]$$

- For the same reason, when we include into our next one, we get cube roots and adjoin 1/3
- continuing on in the same fashion, we get \mathbb{Q} !

Similarly, inclusion into $\pi_1^{\text{tr}}(GL)_{3!}$ picks up cube roots:

$$\pi_1^{\text{tr}}(GL)_6 \simeq \mathbb{Z} \left[\frac{1}{2}, \frac{1}{3} \right]$$

In the n -th inclusion, we pick up n -th roots and so we adjoin $\frac{1}{n}$ to the preceding group. Therefore,

$$\pi_1^{\text{tr}}(GL) \simeq \mathbb{Z} \left[\frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots \right]$$

- For the same reason, when we include into our next one, we get cube roots and adjoin $1/3$
- continuing on in the same fashion, we get \mathbb{Q} !

Similarly, inclusion into $\pi_1^{\text{tr}}(GL)_{2!}$ picks up cube roots:

$$\pi_1^{\text{tr}}(GL)_6 \simeq \mathbb{Z} \left[\frac{1}{2}, \frac{1}{3} \right]$$

In the n -th inclusion, we pick up n -th roots and so we adjoin $\frac{1}{n}$ to the preceding group. Therefore,

$$\pi_1^{\text{tr}}(GL) \simeq \mathbb{Z} \left[\frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots \right]$$

Similarly, inclusion into $\pi_1^{\text{tr}}(GL)_{3!}$ picks up cube roots:

$$\pi_1^{\text{tr}}(GL)_6 \simeq \mathbb{Z} \left[\frac{1}{2}, \frac{1}{3} \right]$$

In the n -th inclusion, we pick up n -th roots and so we adjoin $\frac{1}{n}$ to the preceding group. Therefore,

$$\pi_1^{\text{tr}}(GL) \simeq \mathbb{Z} \left[\frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots \right] \simeq \mathbb{Q}.$$

- For the same reason, when we include into our next one, we get cube roots and adjoin $1/3$
- continuing on in the same fashion, we get \mathbb{Q} !

Similarly, inclusion into $\pi_1^{\text{tr}}(GL)_{3!}$ picks up cube roots:

$$\pi_1^{\text{tr}}(GL)_6 \simeq \mathbb{Z} \left[\frac{1}{2}, \frac{1}{3} \right]$$

In the n -th inclusion, we pick up n -th roots and so we adjoin $\frac{1}{n}$ to the preceding group. Therefore,

$$\pi_1^{\text{tr}}(GL) \simeq \mathbb{Z} \left[\frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots \right] \simeq \mathbb{Q}.$$

Thank You!

2022-06-03

- That's it! The algebraic topology of free sets is in its infancy but results look promising!
- Thank you!