

# Searching for Holes in the Matrix Universe

Lucas Kerbs

Spring 2022

2022-06-02

## Searching for Holes in the Matrix Universe

Searching for Holes in the Matrix Universe

Lucas Kerbs

Spring 2022

- Eventual goal: lift the tools of algebraic topology to spaces of matrices
- If we only consider  $2 \times 2$  matrices we can use classical theory
- The moment we want more than one size, things the classical theory breaks down
- Today we will develop some *fairly heftly* tools to do just that
- Along the way, hopefully I can convince you that this is an interesting question.
- To do so, we need to go back to our mathematical roots

# Part I: Objects and Maps

A Naive Attempt

2022-06-02

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

Part I: Objects and Maps  
A Naive Attempt

- That’s right—objects and maps.
- Our Naive attempt involves that looking at lifting functions on  $\mathbb{R}$  or  $\mathbb{C}$  to accept matrices as their input.
- An operator theorist would call this a “functional calculus”

# Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ( $A = A^*$ ) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix

# Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.  
 $A$  is diagonalizable as  $A = U \Lambda U^*$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ( $A = A^*$ ) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.

# Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.  
 $A$  is diagonalizable as  $A = U\Lambda U^*$

$$f(A) = a_n A^n + \cdots + a_1 A + a_0 I_k$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

$$f(A) = a_n A^n + \cdots + a_1 A + a_0 I_k$$

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ( $A = A^*$ ) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.
- Watch what happens when we plug this into our polynomial — need to be careful with the constant term.

# Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.

$A$  is diagonalizable as  $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \end{aligned}$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.  
 $A$  is diagonalizable as  $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \end{aligned}$$

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ( $A = A^*$ ) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.
- Watch what happens when we plug this into our polynomial — need to be careful with the constant term.

## Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.

$A$  is diagonalizable as  $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \end{aligned}$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.  
 $A$  is diagonalizable as  $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \end{aligned}$$

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ( $A = A^*$ ) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.
- Watch what happens when we plug this into our polynomial — need to be careful with the constant term.

## Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.

$A$  is diagonalizable as  $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \end{aligned}$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.  
 $A$  is diagonalizable as  $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \end{aligned}$$

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ( $A = A^*$ ) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.
- Watch what happens when we plug this into our polynomial — need to be careful with the constant term.



## Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.

$A$  is diagonalizable as  $A = U\Lambda U^*$

$$\begin{aligned}
 f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\
 &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\
 &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\
 &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \\
 &= U (f(\Lambda)) U^*
 \end{aligned}$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.  
 $A$  is diagonalizable as  $A = U\Lambda U^*$

$$\begin{aligned}
 f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\
 &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\
 &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\
 &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \\
 &= U (f(\Lambda)) U^*
 \end{aligned}$$

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ( $A = A^*$ ) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.
- Watch what happens when we plug this into our polynomial — need to be careful with the constant term.

# Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.

$A$  is diagonalizable as  $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \\ &= U (f(\Lambda)) U^* \end{aligned}$$

$$f \left( \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} \right)$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.  
 $A$  is diagonalizable as  $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \\ &= U (f(\Lambda)) U^* \end{aligned}$$

$$f \left( \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} \right)$$

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ( $A = A^*$ ) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.
- Watch what happens when we plug this into our polynomial — need to be careful with the constant term.

## Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.

$A$  is diagonalizable as  $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \\ &= U (f(\Lambda)) U^* \end{aligned}$$

$$f\left(\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}\right) = \begin{bmatrix} f(\lambda_1) & & \\ & \ddots & \\ & & f(\lambda_n) \end{bmatrix}$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.  
 $A$  is diagonalizable as  $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \\ &= U (f(\Lambda)) U^* \end{aligned}$$

$$f\left(\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}\right) = \begin{bmatrix} f(\lambda_1) & & \\ & \ddots & \\ & & f(\lambda_n) \end{bmatrix}$$

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ( $A = A^*$ ) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.
- Watch what happens when we plug this into our polynomial — need to be careful with the constant term.
- This application along the diagonal is precisely the behavior we want to emulate in the functional calculus.

## Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.

$A$  is diagonalizable as  $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \\ &= U (f(\Lambda)) U^* \end{aligned}$$

$$f\left(\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}\right) = \begin{bmatrix} f(\lambda_1) & & \\ & \ddots & \\ & & f(\lambda_n) \end{bmatrix}$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Functional Calculus

Let  $f \in \mathbb{R}[x]$  and  $A \in M_k(\mathbb{C})$  be self adjoint.  
 $A$  is diagonalizable as  $A = U\Lambda U^*$

$$\begin{aligned} f(A) &= a_n A^n + \cdots + a_1 A + a_0 I_k \\ &= a_n (U\Lambda U^*)^n + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= a_n U\Lambda^n U^* + \cdots + a_1 U\Lambda U^* + a_0 I_k \\ &= U (a_n \Lambda^n + \cdots + a_1 \Lambda + a_0 I_k) U^* \\ &= U (f(\Lambda)) U^* \end{aligned}$$

$$f\left(\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}\right) = \begin{bmatrix} f(\lambda_1) & & \\ & \ddots & \\ & & f(\lambda_n) \end{bmatrix}$$

- Polynomials are the most well behaved functions we have, so lets start with a polynomial and a self adjoint ( $A = A^*$ ) matrix.
- You might say that SA is unnecessary bc we can already evaluate a polynomial on a matrix
- Since we are SA, we can diagonalize with unitary matrices.
- Watch what happens when we plug this into our polynomial — need to be careful with the constant term.
- This application along the diagonal is precisely the behavior we want to emulate in the functional calculus.

Let  $\mathbb{H}_n$  be the set of  $n \times n$  self adjoint matrices, and define

$$\mathbb{H} = \bigcup_{n \in \mathbb{N}} \mathbb{H}_n, \quad \mathcal{M} = \bigcup_{n \in \mathbb{N}} M_n(\mathbb{C})$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

Let  $\mathbb{H}_n$  be the set of  $n \times n$  self adjoint matrices, and define

$$\mathbb{H} = \bigcup_{n \in \mathbb{N}} \mathbb{H}_n, \quad \mathcal{M} = \bigcup_{n \in \mathbb{N}} M_n(\mathbb{C})$$

- With the polynomial case in mind, we can extend a general function. First, a piece of notation
- Lets grab a function on the real line and the self adjoint matrices with their spectrum in that domain
- Then we can lift  $g$  by emulating the behavior of polynomials.
- unwrap a self adjoint matrix, apply  $g$  to the diagonal, then wrap it back up
- Something to notice about this functional calculus—it treats direct sums *very* well
- This is all well and good, but can we do anything with these new functions?

Let  $\mathbb{H}_n$  be the set of  $n \times n$  self adjoint matrices, and define

$$\mathbb{H} = \bigcup_{n \in \mathbb{N}} \mathbb{H}_n, \quad \mathcal{M} = \bigcup_{n \in \mathbb{N}} M_n(\mathbb{C})$$

**Definition:**

Let  $g : [a, b] \rightarrow \mathbb{C}$  and  $D \subset \mathbb{H}$  denote the set of self adjoint matrices with their spectrum in  $[a, b]$ .

2022-06-02

Searching for Holes in the Matrix Universe

- With the polynomial case in mind, we can extend a general function. First, a piece of notation
- Lets grab a function on the real line and the self adjoint matrices with their spectrum in that domain
- Then we can lift  $g$  by emulating the behavior of polynomials.
- unwrap a self adjoint matrix, apply  $g$  to the diagonal, then wrap it back up
- Something to notice about this functional calculus—it treats direct sums *very* well
- This is all well and good, but can we do anything with these new functions?

Let  $\mathbb{H}_n$  be the set of  $n \times n$  self adjoint matrices, and define

$$\mathbb{H} = \bigcup_{n \in \mathbb{N}} \mathbb{H}_n, \quad \mathcal{M} = \bigcup_{n \in \mathbb{N}} M_n(\mathbb{C})$$

**Definition:**  
Let  $g : [a, b] \rightarrow \mathbb{C}$  and  $D \subset \mathbb{H}$  denote the set of self adjoint matrices with their spectrum in  $[a, b]$ .

Let  $\mathbb{H}_n$  be the set of  $n \times n$  self adjoint matrices, and define

$$\mathbb{H} = \bigcup_{n \in \mathbb{N}} \mathbb{H}_n, \quad \mathcal{M} = \bigcup_{n \in \mathbb{N}} M_n(\mathbb{C})$$

### Definition:

Let  $g : [a, b] \rightarrow \mathbb{C}$  and  $D \subset \mathbb{H}$  denote the set of self adjoint matrices with their spectrum in  $[a, b]$ . Then

$$g : D \longrightarrow \mathcal{M}$$

$$X = U \Lambda U^* \longmapsto U \begin{bmatrix} g(\lambda_1) & & \\ & \ddots & \\ & & g(\lambda_n) \end{bmatrix} U^*.$$

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part I: Objects and Maps

#### └ Functional Calculus

- With the polynomial case in mind, we can extend a general function. First, a piece of notation
- Lets grab a function on the real line and the self adjoint matrices with their spectrum in that domain
- Then we can lift  $g$  by emulating the behavior of polynomials.
- unwrap a self adjoint matrix, apply  $g$  to the diagonal, then wrap it back up
- Something to notice about this functional calculus—it treats direct sums *very* well
- This is all well and good, but can we do anything with these new functions?

Let  $\mathbb{H}_n$  be the set of  $n \times n$  self adjoint matrices, and define

$$\mathbb{H} = \bigcup_{n \in \mathbb{N}} \mathbb{H}_n, \quad \mathcal{M} = \bigcup_{n \in \mathbb{N}} M_n(\mathbb{C})$$

**Definition:**

Let  $g : [a, b] \rightarrow \mathbb{C}$  and  $D \subset \mathbb{H}$  denote the set of self adjoint matrices with their spectrum in  $[a, b]$ . Then

$$g : D \longrightarrow \mathcal{M}$$

$$X = U \Lambda U^* \longmapsto U \begin{bmatrix} g(\lambda_1) & & \\ & \ddots & \\ & & g(\lambda_n) \end{bmatrix} U^*.$$

Let  $\mathbb{H}_n$  be the set of  $n \times n$  self adjoint matrices, and define

$$\mathbb{H} = \bigcup_{n \in \mathbb{N}} \mathbb{H}_n, \quad \mathcal{M} = \bigcup_{n \in \mathbb{N}} M_n(\mathbb{C})$$

### Definition:

Let  $g : [a, b] \rightarrow \mathbb{C}$  and  $D \subset \mathbb{H}$  denote the set of self adjoint matrices with their spectrum in  $[a, b]$ . Then

$$g : D \longrightarrow \mathcal{M}$$

$$X = U \Lambda U^* \longmapsto U \begin{bmatrix} g(\lambda_1) & & \\ & \ddots & \\ & & g(\lambda_n) \end{bmatrix} U^*.$$

**Important:** In this functional calculus,

$$g(X \oplus Y) = g(X) \oplus g(Y)$$

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part I: Objects and Maps

#### └ Functional Calculus

- With the polynomial case in mind, we can extend a general function. First, a piece of notation
- Lets grab a function on the real line and the self adjoint matrices with their spectrum in that domain
- Then we can lift  $g$  by emulating the behavior of polynomials.
- unwrap a self adjoint matrix, apply  $g$  to the diagonal, then wrap it back up
- Something to notice about this functional calculus—it treats direct sums *very* well
- This is all well and good, but can we do anything with these new functions?

Let  $\mathbb{H}_n$  be the set of  $n \times n$  self adjoint matrices, and define

$$\mathbb{H} = \bigcup_{n \in \mathbb{N}} \mathbb{H}_n, \quad \mathcal{M} = \bigcup_{n \in \mathbb{N}} M_n(\mathbb{C})$$

**Definition:**

Let  $g : [a, b] \rightarrow \mathbb{C}$  and  $D \subset \mathbb{H}$  denote the set of self adjoint matrices with their spectrum in  $[a, b]$ . Then

$$g : D \longrightarrow \mathcal{M}$$

$$X = U \Lambda U^* \longmapsto U \begin{bmatrix} g(\lambda_1) & & \\ & \ddots & \\ & & g(\lambda_n) \end{bmatrix} U^*.$$

**Important:** In this functional calculus,

$$g(X \oplus Y) = g(X) \oplus g(Y)$$



## Directional Derivative

**Definition:** *Directional Derivative*

Fix some  $X \in \mathbb{H}_n$ . The derivative of  $f$  at  $X$  in the direction  $H \in M_n(\mathbb{C})$  is

$$Df(X)[H] = \lim_{t \rightarrow 0} \frac{f(X + tH) - f(X)}{t}$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Directional Derivative

**Definition:** *Directional Derivative*  
Fix some  $X \in \mathbb{H}_n$ . The derivative of  $f$  at  $X$  in the direction  $H \in M_n(\mathbb{C})$  is

$$Df(X)[H] = \lim_{t \rightarrow 0} \frac{f(X + tH) - f(X)}{t}$$

- We can define a directional derivative—as long as we are careful to have the direction in the same “level-wise” slice.
- Notice that, with some special attention to what operation we are carrying out, this is the exact same definition as classic multivariable calculus.
- There is another formulation that is (generally) more useful for computation

# Directional Derivative

## Definition: *Directional Derivative*

Fix some  $X \in \mathbb{H}_n$ . The derivative of  $f$  at  $X$  in the direction  $H \in M_n(\mathbb{C})$  is

$$Df(X)[H] = \lim_{t \rightarrow 0} \frac{f(X + tH) - f(X)}{t}$$

Alternatively,

$$Df(X)[H] = \left. \frac{df(X + tH)}{dt} \right|_{t=0}$$

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part I: Objects and Maps

#### └ Functional Calculus

#### └ Directional Derivative

- We can define a directional derivative—as long as we are careful to have the direction in the same “level-wise” slice.
- Notice that, with some special attention to what operation we are carrying out, this is the exact same definition as classic multivariable calculus.
- There is another formulation that is (generally) more useful for computation

#### Directional Derivative

**Definition: Directional Derivative**  
Fix some  $X \in \mathbb{H}_n$ . The derivative of  $f$  at  $X$  in the direction  $H \in M_n(\mathbb{C})$  is

$$Df(X)[H] = \lim_{t \rightarrow 0} \frac{f(X + tH) - f(X)}{t}$$

Alternatively,

$$Df(X)[H] = \left. \frac{df(X + tH)}{dt} \right|_{t=0}$$

Example:  $g(x) = x^3$

$$g(X + tH) =$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example:  $g(x) = x^3$ Example:  $g(x) = x^3$ 

$$g(X + tH) =$$

- Now we consider an example. Since  $Df(X)[H]$  is linear, we can just work with a single monomial
- First we expand  $(x + th)^3$ —but we can't use the binomial theorem since  $x$  and  $h$  don't commute
- Once we expand, we take standard derivatives w.r.t  $t$ —treating  $X$  and  $H$  as formal symbols.

Example:  $g(x) = x^3$

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example:  $g(x) = x^3$ Example:  $g(x) = x^3$ 

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

- Now we consider an example. Since  $Df(X)[H]$  is linear, we can just work with a single monomial
- First we expand  $(x + th)^3$ —but we can't use the binomial theorem since  $x$  and  $h$  don't commute
- Once we expand, we take standard derivatives w.r.t  $t$ —treating  $X$  and  $H$  as formal symbols.

Example:  $g(x) = x^3$

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

From here, we can calculate:

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example:  $g(x) = x^3$ Example:  $g(x) = x^3$ 

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

From here, we can calculate:

- Now we consider an example. Since  $Df(X)[H]$  is linear, we can just work with a single monomial
- First we expand  $(x + th)^3$ —but we can't use the binomial theorem since  $x$  and  $h$  don't commute
- Once we expand, we take standard derivatives w.r.t  $t$ —treating  $X$  and  $H$  as formal symbols.

Example:  $g(x) = x^3$

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

From here, we can calculate:

$$\frac{d}{dt}g(X + tH) = X^2H + XHX + 2tXH^2 + HX^2 \\ + 2tHXX + 2tH^2X + 3t^2H^3$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example:  $g(x) = x^3$ Example:  $g(x) = x^3$ 

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

From here, we can calculate:

$$\frac{d}{dt}g(X + tH) = X^2H + XHX + 2tXH^2 + HX^2 \\ + 2tHXX + 2tH^2X + 3t^2H^3$$

- Now we consider an example. Since  $Df(X)[H]$  is linear, we can just work with a single monomial
- First we expand  $(x + th)^3$ —but we can't use the binomial theorem since  $x$  and  $h$  don't commute
- Once we expand, we take standard derivatives w.r.t  $t$ —treating  $X$  and  $H$  as formal symbols.

Example:  $g(x) = x^3$

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

From here, we can calculate:

$$\frac{d}{dt}g(X + tH) = X^2H + XHX + 2tXH^2 + HX^2 \\ + 2tHXX + 2tH^2X + 3t^2H^3$$

$$\frac{d^2}{dt^2}g(X + tH) = 2XH^2 + 2HXX + 2H^2X + 6tH^3$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example:  $g(x) = x^3$ Example:  $g(x) = x^3$ 

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

From here, we can calculate:

$$\frac{d}{dt}g(X + tH) = X^2H + XHX + 2tXH^2 + HX^2 \\ + 2tHXX + 2tH^2X + 3t^2H^3$$

$$\frac{d^2}{dt^2}g(X + tH) = 2XH^2 + 2HXX + 2H^2X + 6tH^3$$

- Now we consider an example. Since  $Df(X)[H]$  is linear, we can just work with a single monomial
- First we expand  $(x + th)^3$ —but we can't use the binomial theorem since  $x$  and  $h$  don't commute
- Once we expand, we take standard derivatives w.r.t  $t$ —treating  $X$  and  $H$  as formal symbols.

Example:  $g(x) = x^3$

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

From here, we can calculate:

$$\frac{d}{dt}g(X + tH) = X^2H + XHX + 2tXH^2 + HX^2 \\ + 2tHXX + 2tH^2X + 3t^2H^3$$

$$\frac{d^2}{dt^2}g(X + tH) = 2XH^2 + 2HXX + 2H^2X + 6tH^3$$

$$\frac{d^3}{dt^3}g(X + tH) = 6H^3.$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example:  $g(x) = x^3$ Example:  $g(x) = x^3$ 

$$g(X + tH) = X^3 + tX^2H + tXHX + t^2XH^2 \\ + tHX^2 + t^2HXX + t^2H^2X + t^3H^3.$$

From here, we can calculate:

$$\frac{d}{dt}g(X + tH) = X^2H + XHX + 2tXH^2 + HX^2 \\ + 2tHXX + 2tH^2X + 3t^2H^3$$

$$\frac{d^2}{dt^2}g(X + tH) = 2XH^2 + 2HXX + 2H^2X + 6tH^3$$

$$\frac{d^3}{dt^3}g(X + tH) = 6H^3.$$

- Now we consider an example. Since  $Df(X)[H]$  is linear, we can just work with a single monomial
- First we expand  $(x + th)^3$ —but we can't use the binomial theorem since  $x$  and  $h$  don't commute
- Once we expand, we take standard derivatives w.r.t  $t$ —treating  $X$  and  $H$  as formal symbols.



Example:  $g(x) = x^3$

And so the first 3 directional derivatives are:

2022-06-02

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example:  $g(x) = x^3$

Example:  $g(x) = x^3$

And so the first 3 directional derivatives are:

- Now all we have to do is set  $t = 0$  in the previous expressions and we get the first three directional derivatives
- Note that they are all in the same direction—mixed derivatives are possible but we won't need them.

Example:  $g(x) = x^3$

And so the first 3 directional derivatives are:

$$Df(X)[H] = X^2H + XHX + HX^2$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example:  $g(x) = x^3$ Example:  $g(x) = x^3$ 

And so the first 3 directional derivatives are:

$$Df(X)[H] = X^2H + XHX + HX^2$$

- Now all we have to do is set  $t = 0$  in the previous expressions and we get the first three directional derivatives
- Note that they are all in the same direction—mixes derivatives are possible but we won't need them.

Example:  $g(x) = x^3$

And so the first 3 directional derivatives are:

$$Df(X)[H] = X^2H + XHX + HX^2$$

$$D^{(2)}f(X)[H] = 2XH^2 + 2HXX + 2H^2X$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example:  $g(x) = x^3$ Example:  $g(x) = x^3$ 

And so the first 3 directional derivatives are:

$$Df(X)[H] = X^2H + XHX + HX^2$$

$$D^{(2)}f(X)[H] = 2XH^2 + 2HXX + 2H^2X$$

- Now all we have to do is set  $t = 0$  in the previous expressions and we get the first three directional derivatives
- Note that they are all in the same direction—mixes derivatives are possible but we won't need them.

Example:  $g(x) = x^3$

And so the first 3 directional derivatives are:

$$Df(X)[H] = X^2H + XHX + HX^2$$

$$D^{(2)}f(X)[H] = 2XH^2 + 2HXX + 2H^2X$$

$$D^{(3)}f(X)[H] = 6H^3$$

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Functional Calculus

└ Example:  $g(x) = x^3$ 

- Now all we have to do is set  $t = 0$  in the previous expressions and we get the first three directional derivatives
- Note that they are all in the same direction—mixes derivatives are possible but we won't need them.

Example:  $g(x) = x^3$ 

And so the first 3 directional derivatives are:

$$Df(X)[H] = X^2H + XHX + HX^2$$

$$D^{(2)}f(X)[H] = 2XH^2 + 2HXX + 2H^2X$$

$$D^{(3)}f(X)[H] = 6H^3$$

# Part I.5: Objects and Maps

## A Second Attempt

2022-06-02

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

Part I.5: Objects and Maps  
*A Second Attempt*

- In seeking a more general theory we need to leave the world of this “SA functional calculus” behind.
- Rather than lifting functions to be matrix valued, we will define *new* objects that behave like those we just looked at.

# Some Definitions

**Definition:** *Matrix Universe*

The  $g$ -dimensional **Matrix Universe** is

$$\mathcal{M}^g = \bigcup_{n \in \mathbb{N}} (M_n(\mathbb{C}))^g$$

2022-06-02

## Searching for Holes in the Matrix Universe

- └ Part I: Objects and Maps
  - └ Matrix Universe
    - └ Some Definitions

**Definition:** *Matrix Universe*  
The  $g$ -dimensional **Matrix Universe** is  
$$\mathcal{M}^g = \bigcup_{n \in \mathbb{N}} (M_n(\mathbb{C}))^g$$

- Definition of the matrix universe —  $g$  tuples of matrices of all sizes
- By convetion, tuples are likes size

# Some Definitions

**Definition:** *Matrix Universe*

The  $g$ -dimensional **Matrix Universe** is

$$\mathcal{M}^g = \bigcup_{n \in \mathbb{N}} (M_n(\mathbb{C}))^g$$

By convention,  $X \in \mathcal{M}^g$  is a tuple of like-size matrices

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

└ Some Definitions

**Definition:** *Matrix Universe*  
The  $g$ -dimensional **Matrix Universe** is

$$\mathcal{M}^g = \bigcup_{n \in \mathbb{N}} (M_n(\mathbb{C}))^g$$

By convention,  $X \in \mathcal{M}^g$  is a tuple of like-size matrices

- Definition of the matrix universe —  $g$  tuples of matrices of all sizes
- By convetion, tuples are likes size

**Definition:** *Free Set*

We say  $D \subset \mathcal{M}^g$  is a **free set** if it is closed with respect to direct sums and unitary conjugation. That is,

2022-06-02

Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

**Definition: Free Set**  
We say  $D \subset \mathcal{M}^g$  is a **free set** if it is closed with respect to direct sums and unitary conjugation. That is,

- In math we often think about substructures that capture the implicit structure our space (subgroup, subspace, etc)
- In the nc setting, this is a *free set*, also called nc set
- direct sums and unitary conjugation are component wise
- If you see a  $D$ , you can assume that it is a free set.
- A subscript denotes a level-wise slice
- Note that this requires a lot of structure on free sets—we want to put a name to these structure.



### Definition: *Free Set*

We say  $D \subset \mathcal{M}^g$  is a **free set** if it is closed with respect to direct sums and unitary conjugation. That is,

$$\textcircled{1} \quad X, Y \in D \text{ means } X \oplus Y = (X_1 \oplus Y_1, \dots, X_g \oplus Y_g) \in D.$$

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part I: Objects and Maps

#### └ Matrix Universe

**Definition: Free Set**  
We say  $D \subset \mathcal{M}^g$  is a **free set** if it is closed with respect to direct sums and unitary conjugation. That is,  
 $\bullet \quad X, Y \in D \text{ means } X \oplus Y = (X_1 \oplus Y_1, \dots, X_g \oplus Y_g) \in D.$

- In math we often think about substructures that capture the implicit structure our space (subgroup, subspace, etc)
- In the nc setting, this is a *free set*, also called nc set
- direct sums and unitary conjugation are component wise
- If you see a  $D$ , you can assume that it is a free set.
- A subscript denotes a level-wise slice
- Note that this requires a lot of structure on free sets—we want to put a name to these structure.

### Definition: *Free Set*

We say  $D \subset \mathcal{M}^g$  is a **free set** if it is closed with respect to direct sums and unitary conjugation. That is,

- ①  $X, Y \in D$  means  $X \oplus Y = (X_1 \oplus Y_1, \dots, X_g \oplus Y_g) \in D$ .
- ② For  $X, U$  like-size matrices with  $U$  unitary and  $X \in D$ , then  $UXU^* = (UX_1 U^*, \dots, UX_g U^*) \in D$ .

.

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part I: Objects and Maps

#### └ Matrix Universe

**Definition: Free Set**  
We say  $D \subset \mathcal{M}^g$  is a **free set** if it is closed with respect to direct sums and unitary conjugation. That is,  
 •  $X, Y \in D$  means  $X \oplus Y = (X_1 \oplus Y_1, \dots, X_g \oplus Y_g) \in D$ .  
 • For  $X, U$  like-size matrices with  $U$  unitary and  $X \in D$ , then  $UXU^* = (UX_1 U^*, \dots, UX_g U^*) \in D$ .

- In math we often think about substructures that capture the implicit structure our space (subgroup, subspace, etc)
- In the nc setting, this is a *free set*, also called nc set
- direct sums and unitary conjugation are component wise
- If you see a  $D$ , you can assume that it is a free set.
- A subscript denotes a level-wise slice
- Note that this requires a lot of structure on free sets—we want to put a name to these structure.

### Definition: *Free Set*

We say  $D \subset \mathcal{M}^g$  is a **free set** if it is closed with respect to direct sums and unitary conjugation. That is,

- ①  $X, Y \in D$  means  $X \oplus Y = (X_1 \oplus Y_1, \dots, X_g \oplus Y_g) \in D$ .
- ② For  $X, U$  like-size matrices with  $U$  unitary and  $X \in D$ , then  $UXU^* = (UX_1U^*, \dots, UX_gU^*) \in D$ .

For  $D$  a free set, define  $D_n = D \cap (M_n(\mathbb{C}))^g$ .

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part I: Objects and Maps

#### └ Matrix Universe

- In math we often think about substructures that capture the implicit structure our space (subgroup, subspace, etc)
- In the nc setting, this is a *free set*, also called nc set
- direct sums and unitary conjugation are component wise
- If you see a  $D$ , you can assume that it is a free set.
- A subscript denotes a level-wise slice
- Note that this requires a lot of structure on free sets—we want to put a name to these structure.

**Definition: Free Set**  
We say  $D \subset \mathcal{M}^g$  is a **free set** if it is closed with respect to direct sums and unitary conjugation. That is,  
 •  $X, Y \in D$  means  $X \oplus Y = (X_1 \oplus Y_1, \dots, X_g \oplus Y_g) \in D$ .  
 • For  $X, U$  like-size matrices with  $U$  unitary and  $X \in D$ , then  $UXU^* = (UX_1U^*, \dots, UX_gU^*) \in D$ .  
 For  $D$  a free set, define  $D_n = D \cap (M_n(\mathbb{C}))^g$ .

### Definition: *Fiber*

Given  $X \in \mathcal{M}^g$ , a tuple of  $n \times n$  matrices, the **fiber** of  $X$  is the set

$$\{X^{\oplus k} \mid k \in \mathbb{N}\}.$$

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part I: Objects and Maps

#### └ Matrix Universe

**Definition: Fiber**  
Given  $X \in \mathcal{M}^g$ , a tuple of  $n \times n$  matrices, the **fiber** of  $X$  is the set

$$\{X^{\oplus k} \mid k \in \mathbb{N}\}.$$

- The fiber is all the points “above” a given point. Conceptually, we imagine identification along the fiber—this will become important when we start doing topology
- The envelope (which will be less important to us) is the unitary smearing of the fiber at each level.

### Definition: *Fiber*

Given  $X \in \mathcal{M}^g$ , a tuple of  $n \times n$  matrices, the **fiber** of  $X$  is the set

$$\{X^{\oplus k} \mid k \in \mathbb{N}\}.$$

### Definition: *Envelope*

Given  $X \in \mathcal{M}^g$ , a tuple of  $n \times n$  matrices, the **envelope** of  $X$  is the set

$$\{U^* X^{\oplus k} U \mid k \in \mathbb{N}, U \text{ Unitary}\}.$$

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part I: Objects and Maps

#### └ Matrix Universe

- The fiber is all the points “above” a given point. Conceptually, we imagine identification along the fiber—this will become important when we start doing topology
- The envelope (which will be less important to us) is the unitary smearing of the fiber at each level.

#### Definition: *Fiber*

Given  $X \in \mathcal{M}^g$ , a tuple of  $n \times n$  matrices, the **fiber** of  $X$  is the set

$$\{X^{\oplus k} \mid k \in \mathbb{N}\}.$$

#### Definition: *Envelope*

Given  $X \in \mathcal{M}^g$ , a tuple of  $n \times n$  matrices, the **envelope** of  $X$  is the set

$$\{U^* X^{\oplus k} U \mid k \in \mathbb{N}, U \text{ Unitary}\}.$$

# A bit of topology

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

└ A bit of topology

- If we are going to look for holes and build up the algebraic topology, we need a point set topology first—so there is a natural question.
- Bad news: there isn't a natural choice
- There are a handful of candidates (fine, fat, free, nc Zariski). I wish we had time to go into detail.
- For us, free sets are open if their level-wise restriction is open
- Other point-set characterizations are similar.

# A bit of topology

What does it mean for  $D \subset \mathcal{M}^g$  to be open?

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

└ A bit of topology

- If we are going to look for holes and build up the algebraic topology, we need a point set topology first—so there is a natural question.
- Bad news: there isn't a natural choice
- There are a handful of candidates (fine, fat, free, nc Zariski). I wish we had time to go into detail.
- For us, free sets are open if their level-wise restriction is open
- Other point-set characterizations are similar.

# A bit of topology

What does it mean for  $D \subset \mathcal{M}^g$  to be open?

- There is not a canonical topology

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

└ A bit of topology

- There is not a canonical topology

- If we are going to look for holes and build up the algebraic topology, we need a point set topology first—so there is a natural question.
- Bad news: there isn't a natural choice
- There are a handful of candidates (fine, fat, free, nc Zariski). I wish we had time to go into detail.
- For us, free sets are open if their level-wise restriction is open
- Other point-set characterizations are similar.



# A bit of topology

What does it mean for  $D \subset \mathcal{M}^g$  to be open?

- There is not a canonical topology
- For us, we will say that  $D$  is open if each  $D_n$  is open.

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

└ A bit of topology

What does it mean for  $D \subset M^g$  to be open?

- There is not a canonical topology
- For us, we will say that  $D$  is open if each  $D_n$  is open.

- If we are going to look for holes and build up the algebraic topology, we need a point set topology first—so there is a natural question.
- Bad news: there isn't a natural choice
- There are a handful of candidates (fine, fat, free, nc Zariski). I wish we had time to go into detail.
- For us, free sets are open if their level-wise restriction is open
- Other point-set characterizations are similar.

## A bit of topology

What does it mean for  $D \subset \mathcal{M}^g$  to be open?

- There is not a canonical topology
- For us, we will say that  $D$  is open if each  $D_n$  is open.
- Simply connected, connected, bounded, etc. are defined similarly.

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part I: Objects and Maps

└ Matrix Universe

└ A bit of topology

What does it mean for  $D \subset M^g$  to be open?

- There is not a canonical topology
- For us, we will say that  $D$  is open if each  $D_n$  is open.
- Simply connected, connected, bounded, etc. are defined similarly.

- If we are going to look for holes and build up the algebraic topology, we need a point set topology first—so there is a natural question.
- Bad news: there isn't a natural choice
- There are a handful of candidates (fine, fat, free, nc Zariski). I wish we had time to go into detail.
- For us, free sets are open if their level-wise restriction is open
- Other point-set characterizations are similar.

# What the natural functions on $\mathcal{M}^g$ ?

2022-06-02

## Searching for Holes in the Matrix Universe

## └ Part I: Objects and Maps

## └ Matrix Universe

└ What the natural functions on  $\mathcal{M}^g$ ?

- We have our objects, but what are the maps?
- Free functions are defined to be anything that behaves like a polynomial
- tracial functions look like traces
- For both of these maps, the directional derivative is defined identically as before—but tracial functions get something extra.

What the natural functions on  $\mathcal{M}^g$ ?**Definition:**

A function  $f : D \rightarrow \mathcal{M}^{\hat{g}}$  is called **free** if

- ①  $f(X \oplus Y) = f(X) \oplus f(Y)$
- ②  $f(UXU^*) = f(U)f(X)f(U^*)$  where  $X$  and  $U$  are like-size and  $U$  is unitary.

2022-06-02

## Searching for Holes in the Matrix Universe

## └ Part I: Objects and Maps

## └ Matrix Universe

└ What the natural functions on  $\mathcal{M}^g$ ?

- We have our objects, but what are the maps?
- Free functions are defined to be anything that behaves like a polynomial
- tracial functions look like traces
- For both of these maps, the directional derivative is defined identically as before—but tracial functions get something extra.

**Definition:**

A function  $f : D \rightarrow \mathcal{M}^{\hat{g}}$  is called **free** if

- $f(X \oplus Y) = f(X) \oplus f(Y)$
- $f(UXU^*) = f(U)f(X)f(U^*)$  where  $X$  and  $U$  are like-size and  $U$  is unitary.

What the natural functions on  $\mathcal{M}^g$ ?**Definition:**

A function  $f : D \rightarrow \mathcal{M}^{\hat{g}}$  is called **free** if

- ①  $f(X \oplus Y) = f(X) \oplus f(Y)$
- ②  $f(UXU^*) = f(U)f(X)f(U^*)$  where  $X$  and  $U$  are like-size and  $U$  is unitary.

**Definition:**

A function  $f : D \rightarrow \mathbb{C}$  is a **tracial function** if

- ①  $f(X \oplus Y) = f(X) + f(Y)$
- ②  $f(UXU^*) = f(X)$  where  $X$  and  $U$  are like-size and  $U$  is unitary.

2022-06-02

## Searching for Holes in the Matrix Universe

## └ Part I: Objects and Maps

## └ Matrix Universe

└ What the natural functions on  $\mathcal{M}^g$ ?

- We have our objects, but what are the maps?
- Free functions are defined to be anything that behaves like a polynomial
- tracial functions look like traces
- For both of these maps, the directional derivative is defined identically as before—but tracial functions get something extra.

**Definition:**

A function  $f : D \rightarrow \mathcal{M}^{\hat{g}}$  is called **free** if

- $f(X \oplus Y) = f(X) \oplus f(Y)$
- $f(UXU^*) = f(U)f(X)f(U^*)$  where  $X$  and  $U$  are like-size and  $U$  is unitary.

**Definition:**

A function  $f : D \rightarrow \mathbb{C}$  is a **tracial function** if

- $f(X \oplus Y) = f(X) + f(Y)$
- $f(UXU^*) = f(X)$  where  $X$  and  $U$  are like-size and  $U$  is unitary.

### Definition: *Free Gradient*

Given a tracial function  $f$ , the free gradient,  $\nabla f$ , is the unique free function satisfying

$$\text{tr}(H \cdot \nabla f(X)) = Df(X)[H],$$

where, if  $A = (A_1, \dots, A_g)$  and  $B = (B_1, \dots, B_g)$  are tuples of like-size matrices then  $A \cdot B = \sum_{i=1}^g A_i B_i$ .

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part I: Objects and Maps

#### └ Uniqueness of the Gradient

**Definition: Free Gradient**  
Given a tracial function  $f$ , the free gradient,  $\nabla f$ , is the unique free function satisfying

$$\text{tr}(H \cdot \nabla f(X)) = Df(X)[H],$$

where, if  $A = (A_1, \dots, A_g)$  and  $B = (B_1, \dots, B_g)$  are tuples of like-size matrices then  $A \cdot B = \sum_{i=1}^g A_i B_i$ .

- The  $\nabla$  of a free function is the unique free function satisfy this equation—where the  $\cdot$  is just like the dot product
- Whenever you see  $\text{tr}(\cdot)$  I want you to think of the inner product—it is slightly distinct but it will make a lot of things make more sense
- Some of you may be hesitant at the fact that I claim  $\nabla$  is unique. Why should this be true?

# Why should $\nabla f$ be unique?

## Theorem (Trace Duality)

*Let  $f, g$  be free functions  $\mathcal{M}^g \rightarrow \mathcal{M}^{\tilde{g}}$ . If  $\text{tr}(H \cdot f) = \text{tr}(H \cdot g)$  for all tuples  $H$ , then  $f = g$ .*

2022-06-02

## Searching for Holes in the Matrix Universe

## └ Part I: Objects and Maps

## └ Uniqueness of the Gradient


└ Why should  $\nabla f$  be unique?

- $f = g$  whenever the domains overlap
- In the vector space setting—with an inner product—this is a fairly immediate result. You would show it by picking vectors of all 0's and a single 1.
- You prove this identically—but with coordinate matrices instead of coordinate vectors.

# Part II: Analytic Continuation and Monodromy



# Analytic Continuation




2022-06-02

## Searching for Holes in the Matrix Universe


└ Part II: Monodromy

└ Analytic Continuation


└ Analytic Continuation




# Analytic Continuation



2022-06-02



# Analytic Continuation




2022-06-02

## Searching for Holes in the Matrix Universe


└ Part II: Monodromy

└ Analytic Continuation

└ Analytic Continuation



# Analytic Continuation




2022-06-02


## Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Analytic Continuation

└ Analytic Continuation




Example: Analytically continuing  $\text{Log } z$ 


2022-06-02

## Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Analytic Continuation

└ Example: Analytically continuing  $\text{Log } z$ Example: Analytically continuing  $\text{Log } z$ 


Example: Analytically continuing  $\text{Log } z$ 


2022-06-02

## Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Analytic Continuation

└ Example: Analytically continuing  $\text{Log } z$ Example: Analytically continuing  $\text{Log } z$ 


Example: Analytically continuing  $\text{Log } z$ 


2022-06-02

## Searching for Holes in the Matrix Universe

## └ Part II: Monodromy

## └ Analytic Continuation

└ Example: Analytically continuing  $\text{Log } z$ Example: Analytically continuing  $\text{Log } z$ 


Example: Analytically continuing  $\text{Log } z$ 

2022-06-02

## Searching for Holes in the Matrix Universe

## └ Part II: Monodromy

## └ Analytic Continuation

└ Example: Analytically continuing  $\text{Log } z$ 



# When are two analytic continuations equal?

2022-06-02

# When are two analytic continuations equal?

## Theorem (Monodromy I)

*Let  $\gamma_1, \gamma_2$  be two paths from  $\alpha$  to  $\beta$  and  $\Gamma_s$  be a fixed-endpoint homotopy between them. If  $f$  can be continued along  $\Gamma_s$  for all  $s \in [0, 1]$ , then the continuations along  $\gamma_1$  and  $\gamma_2$  agree at  $\beta$ .*

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part II: Monodromy

#### └ Monodromy

##### └ When are two analytic continuations equal?


### Theorem (Monodromy I)

*Let  $\gamma_1, \gamma_2$  be two paths from  $\alpha$  to  $\beta$  and  $\Gamma_s$  be a fixed-endpoint homotopy between them. If  $f$  can be continued along  $\Gamma_s$  for all  $s \in [0, 1]$ , then the continuations along  $\gamma_1$  and  $\gamma_2$  agree at  $\beta$ .*

# When are two analytic continuations equal?

## Theorem (Monodromy I)

*Let  $\gamma_1, \gamma_2$  be two paths from  $\alpha$  to  $\beta$  and  $\Gamma_s$  be a fixed-endpoint homotopy between them. If  $f$  can be continued along  $\Gamma_s$  for all  $s \in [0, 1]$ , then the continuations along  $\gamma_1$  and  $\gamma_2$  agree at  $\beta$ .*



2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part II: Monodromy

#### └ Monodromy

#### └ When are two analytic continuations equal?

When are two analytic continuations equal?

Theorem (Monodromy I)

*Let  $\gamma_1, \gamma_2$  be two paths from  $\alpha$  to  $\beta$  and  $\Gamma_s$  be a fixed-endpoint homotopy between them. If  $f$  can be continued along  $\Gamma_s$  for all  $s \in [0, 1]$ , then the continuations along  $\gamma_1$  and  $\gamma_2$  agree at  $\beta$ .*



# When are two analytic continuations equal?

## Theorem (Monodromy II)

*Let  $U \subset \mathbb{C}$  be a disk in  $\mathbb{C}$  centered at  $z_0$  and  $f : U \rightarrow \mathbb{C}$  an analytic function. If  $W$  is an open, simply connected set containing  $U$  and  $f$  continues along any path  $\gamma \subset W$  starting at  $z_0$ , then  $f$  has a unique extension to all of  $W$ .*

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part II: Monodromy

#### └ Monodromy

#### └ When are two analytic continuations equal?

#### Theorem (Monodromy II)

*Let  $U \subset \mathbb{C}$  be a disk in  $\mathbb{C}$  centered at  $z_0$  and  $f : U \rightarrow \mathbb{C}$  an analytic function. If  $W$  is an open, simply connected set containing  $U$  and  $f$  continues along any path  $\gamma \subset W$  starting at  $z_0$ , then  $f$  has a unique extension to all of  $W$ .*

# What about the nc case?

Before we look at a free analogue of the monodromy theorem, we need to ask an important question: What does it mean for a free function to be analytic?

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Free Monodromy

└ What about the nc case?

- be sure to comment on the theorem—it is powerful!

Before we look at a free analogue of the monodromy theorem, we need to ask an important question: What does it mean for a free function to be analytic?

# What about the nc case?

Before we look at a free analogue of the monodromy theorem, we need to ask an important question: What does it mean for a free function to be analytic?

**A:**  $f : D \rightarrow \mathcal{M}^{\hat{g}}$  is analytic if it is analytic as a function of each  $D_n$ .

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Free Monodromy

└ What about the nc case?

- be sure to comment on the theorem—it is powerful!

Before we look at a free analogue of the monodromy theorem, we need to ask an important question: What does it mean for a free function to be analytic?

**A:**  $f : D \rightarrow \mathcal{M}^{\hat{g}}$  is analytic if it is analytic as a function of each  $D_n$ .

# What about the nc case?

Before we look at a free analogue of the monodromy theorem, we need to ask an important question: What does it mean for a free function to be analytic?

**A:**  $f : D \rightarrow \mathcal{M}^{\hat{g}}$  is analytic if it is analytic as a function of each  $D_n$ .

## Theorem (Agler, McCarthy (2016))

*Let  $f : D \rightarrow \mathcal{M}^{\hat{g}}$  be a free function. If  $f$  is locally bounded on each  $D_n$ , then  $f$  is an analytic free function.*

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Free Monodromy

└ What about the nc case?

- be sure to comment on the theorem—it is powerful!

What about the nc case?

Before we look at a free analogue of the monodromy theorem, we need to ask an important question: What does it mean for a free function to be analytic?

**A:**  $f : D \rightarrow \mathcal{M}^{\hat{g}}$  is analytic if it is analytic as a function of each  $D_n$ .

**Theorem (Agler, McCarthy (2016))**  
*Let  $f : D \rightarrow \mathcal{M}^{\hat{g}}$  be a free function. If  $f$  is locally bounded on each  $D_n$ , then  $f$  is an analytic free function.*

Theorem (Free Universal Monodromy, Pascoe 2020)

*Let  $f$  be an analytic free function defined on some ball  $B \subset D$ , for  $D$  an open, connected free set. Then  $f$  analytically continues along every path in  $D$  if and only if  $f$  has a unique analytic continuation to all of  $D$ .*



# Consequences of Free Monodromy

2022-06-02

Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Free Monodromy

└ Consequences of Free Monodromy

# Consequences of Free Monodromy

- ① Free functions can't detect holes!

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part II: Monodromy

└ Free Monodromy

└ Consequences of Free Monodromy

# Consequences of Free Monodromy

- 1 Free functions can't detect holes!
- 2 If we want a fundamental group governed by analytic continuation, we need to look elsewhere.

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part II: Monodromy

#### └ Free Monodromy

##### └ Consequences of Free Monodromy

• Free functions can't detect holes!

• If we want a fundamental group governed by analytic continuation, we need to look elsewhere.

# Part III: Homotopy

2022-06-02

Definition:

A continuous function  $\gamma : [0, 1] \rightarrow D$  **essentially takes**  $X$  to  $Y$  if

$$\gamma(0) = X^{\oplus \ell}, \text{ for some } \ell \in \mathbb{N}$$

$$\gamma(1) = Y^{\oplus k}, \text{ for some } k \in \mathbb{N}.$$

2022-06-02

**Definition:**  
A continuous function  $\gamma : [0, 1] \rightarrow D$  **essentially takes**  $X$  to  $Y$  if

$$\gamma(0) = X^{\oplus \ell}, \text{ for some } \ell \in \mathbb{N}$$


$$\gamma(1) = Y^{\oplus k}, \text{ for some } k \in \mathbb{N}.$$

**Definition:**

A continuous function  $\gamma : [0, 1] \rightarrow D$  **essentially** takes  $X$  to  $Y$  if

$$\gamma(0) = X^{\oplus \ell}, \text{ for some } \ell \in \mathbb{N}$$

$$\gamma(1) = Y^{\oplus k}, \text{ for some } k \in \mathbb{N}.$$



2022-06-02

## Searching for Holes in the Matrix Universe

## └ Part III: Homotopy

## └ A First Fundamental Group

**Definition:**  
A continuous function  $\gamma : [0, 1] \rightarrow D$  **essentially** takes  $X$  to  $Y$  if

$$\gamma(0) = X^{\oplus \ell}, \text{ for some } \ell \in \mathbb{N}$$

$$\gamma(1) = Y^{\oplus k}, \text{ for some } k \in \mathbb{N}.$$



Given  $\gamma$  essentially taking  $X$  to  $Y$  and  $\beta$  taking  $Z$  to  $W$ , define

$$\gamma \oplus \beta(t) := \begin{bmatrix} \gamma(t) & 0 \\ 0 & \beta(t) \end{bmatrix}.$$

2022-06-02

Searching for Holes in the Matrix Universe

└ Part III: Homotopy

└ A First Fundamental Group

Given  $\gamma$  essentially taking  $X$  to  $Y$  and  $\beta$  taking  $Z$  to  $W$ , define

$$\gamma \oplus \beta(t) := \begin{bmatrix} \gamma(t) & 0 \\ 0 & \beta(t) \end{bmatrix}.$$

Given  $\gamma$  essentially taking  $X$  to  $Y$  and  $\beta$  taking  $Z$  to  $W$ , define

$$\gamma \oplus \beta(t) := \begin{bmatrix} \gamma(t) & 0 \\ 0 & \beta(t) \end{bmatrix}.$$

**Definition:**

Let  $\gamma$  and  $\beta$  be paths taking  $X$  to  $Y$  and  $Y$  to  $Z$  respectively. We define their product to be the path essentially taking  $X$  to  $Z$  given by

$$\beta\gamma(t) := \begin{cases} \gamma^{\oplus k}(2t) & t \in [0, 0.5) \\ \beta^{\oplus \ell}(2t - 1) & t \in [0.5, 1] \end{cases}$$

where  $k$  and  $\ell$  are positive integers chosen to make  $\gamma^{\oplus k}$  and  $\beta^{\oplus \ell}$  like size matrices for each  $t \in [0, 1]$ .

2022-06-02

Searching for Holes in the Matrix Universe

Given  $\gamma$  essentially taking  $X$  to  $Y$  and  $\beta$  taking  $Z$  to  $W$ , define

$$\gamma \oplus \beta(t) := \begin{bmatrix} \gamma(t) & 0 \\ 0 & \beta(t) \end{bmatrix}.$$

**Definition:**

Let  $\gamma$  and  $\beta$  be paths taking  $X$  to  $Y$  and  $Y$  to  $Z$  respectively. We define their product to be the path essentially taking  $X$  to  $Z$  given by

$$\beta\gamma(t) := \begin{cases} \gamma^{\oplus k}(2t) & t \in [0, 0.5) \\ \beta^{\oplus \ell}(2t - 1) & t \in [0.5, 1] \end{cases}$$

where  $k$  and  $\ell$  are positive integers chosen to make  $\gamma^{\oplus k}$  and  $\beta^{\oplus \ell}$  like size matrices for each  $t \in [0, 1]$ .



# The Full Fundamental Group

For  $D \subset \mathcal{M}^g$  a connected free set, the **full fundamenal group**,  $\pi_1(D)$ , is the group of paths essentially taking  $X$  to  $X$  up to homotopy equivalence and the relation  $\gamma = \gamma^{\oplus k}$ .

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part III: Homotopy

└ A First Fundamental Group

└ The Full Fundamental Group

- remark about how this misses the link to analytic continuation of functions, but free functions won't do.

Let  $D \subset \mathcal{M}^g$  be a connected, open, free set. If there exists a nonempty, simply-connected, open, free  $B \subset D$ , then we say that  $D$  is **anchored**.

2022-06-02

Let  $D \subset \mathcal{M}^g$  be a connected, open, free set. If there exists a nonempty, simply-connected, open, free  $B \subset D$ , then we say that  $D$  is **anchored**.

For  $D$  an anchored set, and  $B \subset D$  its anchor, we call a tracial function  $f : B \rightarrow \mathbb{C}$  a **global germ** if it analytically continues along every path in  $D$  which starts in  $B$ .

Let  $D \subset \mathcal{M}^g$  be a connected, open, free set. If there exists a nonempty, simply-connected, open, free  $B \subset D$ , then we say that  $D$  is **anchored**.

For  $D$  an anchored set, and  $B \subset D$  its anchor, we call a tracial function  $f : B \rightarrow \mathbb{C}$  a **global germ** if it analytically continues along every path in  $D$  which starts in  $B$ .

For our purposes, we view  $\gamma$  as coupled with its endpoint. Thus, if  $\gamma$  essentially takes  $X$  to  $Y$ , then

$$f(\gamma) = \frac{1}{k}f(Y^{\oplus k}).$$

Let  $D \subset \mathcal{M}^g$  be a connected, open, free set. If there exists a nonempty, simply-connected, open, free  $B \subset D$ , then we say that  $D$  is **anchored**.

For  $D$  an anchored set, and  $B \subset D$  its anchor, we call a tracial function  $f : B \rightarrow \mathbb{C}$  a **global germ** if it analytically continues along every path in  $D$  which starts in  $B$ .

For our purposes, we view  $\gamma$  as coupled with its endpoint. Thus, if  $\gamma$  essentially takes  $X$  to  $Y$ , then

$$f(\gamma) = \frac{1}{k}f(Y^{\oplus k}).$$

# Trace Equivalence

## Definition:

Let  $B \subset D$  be an anchor and fix  $X \in B$ . If  $\gamma$  and  $\beta$  both essentially take  $X$  to  $Y$ , we say they are **trace equivalent** if, for every global germ  $f$  and every path  $\delta$  taking  $Y$  to  $Z$ ,

$$f(\delta\gamma) = f(\delta\beta).$$

2022-06-02

## Searching for Holes in the Matrix Universe

**Definition:**  
Let  $B \subset D$  be an anchor and fix  $X \in B$ . If  $\gamma$  and  $\beta$  both essentially take  $X$  to  $Y$ , we say they are **trace equivalent** if, for every global germ  $f$  and every path  $\delta$  taking  $Y$  to  $Z$ ,

$$f(\delta\gamma) = f(\delta\beta).$$

# Trace Equivalence

## Definition:

Let  $B \subset D$  be an anchor and fix  $X \in B$ . If  $\gamma$  and  $\beta$  both essentially take  $X$  to  $Y$ , we say they are **trace equivalent** if, for every global germ  $f$  and every path  $\delta$  taking  $Y$  to  $Z$ ,

$$f(\delta\gamma) = f(\delta\beta).$$

That is, trace equivalent paths are those which cannot be told apart via analytic continuation of global germ.

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part III: Homotopy

#### └ A Second Fundamental Group

##### └ Trace Equivalence

**Definition:**

Let  $B \subset D$  be an anchor and fix  $X \in B$ . If  $\gamma$  and  $\beta$  both essentially take  $X$  to  $Y$ , we say they are **trace equivalent** if, for every global germ  $f$  and every path  $\delta$  taking  $Y$  to  $Z$ ,

$$f(\delta\gamma) = f(\delta\beta).$$

That is, trace equivalent paths are those which cannot be told apart via analytic continuation of global germ.

# The Tracial Fundamental Group

2022-06-02

- remark that the choice of base point is irrelevant. and that it is a quotient of the full fundamental group

# The Tracial Fundamental Group

Let  $D \subset \mathcal{M}^g$  be an anchored space with  $B$  is anchor. For  $X \in B$  define  $\pi_1^{\text{tr}}(D)$  to be the group of trace equivalent paths essentially taking  $X$  to  $X$ .

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part III: Homotopy

#### └ A Second Fundamental Group

#### └ The Tracial Fundamental Group

- remark that the choice of base point is irrelevant. and that it is a quotient of the full fundamental group

Let  $D \subset \mathcal{M}^g$  be an anchored space with  $B$  is anchor. For  $X \in B$  define  $\pi_1^{\text{tr}}(D)$  to be the group of trace equivalent paths essentially taking  $X$  to  $X$ .



# The Tracial Fundamental Group

Let  $D \subset \mathcal{M}^g$  be an anchored space with  $B$  is anchor. For  $X \in B$  define  $\pi_1^{\text{tr}}(D)$  to be the group of trace equivalent paths essentially taking  $X$  to  $X$ . Computationally, we are still stuck.

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part III: Homotopy

#### └ A Second Fundamental Group

#### └ The Tracial Fundamental Group

- remark that the choice of base point is irrelevant. and that it is a quotient of the full fundamental group

2022-06-02

# Part IV: Cohomology

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ A Short Review of Cohomology

2022-06-02

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ A Short Review of Cohomology

Traditional homology considers a complex of the form

$$\cdots \xleftarrow{\partial_{n-1}} C_{n-1} \xleftarrow{\partial_n} C_n \xleftarrow{\partial_{n+1}} C_{n+1} \xleftarrow{\partial_{n+2}} \cdots$$

2022-06-02

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ A Short Review of Cohomology

Traditional homology considers a complex of the form

$$\cdots \xleftarrow{\partial_{n-1}} C_{n-1} \xleftarrow{\partial_n} C_n \xleftarrow{\partial_{n+1}} C_{n+1} \xleftarrow{\partial_{n+2}} \cdots$$

While cohomology considers a dual complex

$$\cdots \xrightarrow{d_{n-2}} C^{n-1} \xrightarrow{d_{n-1}} C^n \xrightarrow{d_n} C^{n+1} \xrightarrow{d_{n+1}} \cdots$$

2022-06-02

$$\cdots \xleftarrow{\partial_{n-1}} C_{n-1} \xleftarrow{\partial_n} C_n \xleftarrow{\partial_{n+1}} C_{n+1} \xleftarrow{\partial_{n+2}} \cdots$$

$$\cdots \xrightarrow{d_{n-2}} C^{n-1} \xrightarrow{d_{n-1}} C^n \xrightarrow{d_n} C^{n+1} \xrightarrow{d_{n+1}} \cdots$$

Traditional homology considers a complex of the form

$$\cdots \xleftarrow{\partial_{n-1}} C_{n-1} \xleftarrow{\partial_n} C_n \xleftarrow{\partial_{n+1}} C_{n+1} \xleftarrow{\partial_{n+2}} \cdots$$

While cohomology considers a dual complex

$$\cdots \xrightarrow{d_{n-2}} C^{n-1} \xrightarrow{d_{n-1}} C^n \xrightarrow{d_n} C^{n+1} \xrightarrow{d_{n+1}} \cdots$$

In general,  $C^k$  is a group of functions into some abelian group.

2022-06-02

$$\cdots \xleftarrow{\partial_{n-1}} C_{n-1} \xleftarrow{\partial_n} C_n \xleftarrow{\partial_{n+1}} C_{n+1} \xleftarrow{\partial_{n+2}} \cdots$$

$$\cdots \xrightarrow{d_{n-2}} C^{n-1} \xrightarrow{d_{n-1}} C^n \xrightarrow{d_n} C^{n+1} \xrightarrow{d_{n+1}} \cdots$$

Traditional homology considers a complex of the form

$$\cdots \xleftarrow{\partial_{n-1}} C_{n-1} \xleftarrow{\partial_n} C_n \xleftarrow{\partial_{n+1}} C_{n+1} \xleftarrow{\partial_{n+2}} \cdots$$

While cohomology considers a dual complex

$$\cdots \xrightarrow{d_{n-2}} C^{n-1} \xrightarrow{d_{n-1}} C^n \xrightarrow{d_n} C^{n+1} \xrightarrow{d_{n+1}} \cdots$$

In general,  $C^k$  is a group of functions into some abelian group.

The  $k$ -th cohomology group is

$$H^k = \frac{\ker d_i}{\operatorname{Im} d_{i-1}}$$

2022-06-02

# Searching for Holes in the Matrix Universe

## └ Part IV: Cohomology

### └ A Short Review of Cohomology

Traditional homology considers a complex of the form

$$\cdots \xleftarrow{\partial_{n-1}} C_{n-1} \xleftarrow{\partial_n} C_n \xleftarrow{\partial_{n+1}} C_{n+1} \xleftarrow{\partial_{n+2}} \cdots$$

While cohomology considers a dual complex

$$\cdots \xrightarrow{d_{n-2}} C^{n-1} \xrightarrow{d_{n-1}} C^n \xrightarrow{d_n} C^{n+1} \xrightarrow{d_{n+1}} \cdots$$

In general,  $C^k$  is a group of functions into some abelian group.

The  $k$ -th cohomology group is

$$H^k = \frac{\ker d_i}{\operatorname{Im} d_{i-1}}$$

Let  $D$  be an anchored set. Denote the set of (globally defined) tracial functions on  $D$  by  $\mathcal{T}(D)$  and the set of free functions by  $\mathcal{F}(D)$ .

2022-06-02



Let  $D$  be an anchored set. Denote the set of (globally defined) tracial functions on  $D$  by  $\mathcal{T}(D)$  and the set of free functions by  $\mathcal{F}(D)$ . For  $f \in \mathcal{T}(D)$ ,  $\nabla f \in \mathcal{F}(D)$ —so we have the beginnings of a chain complex!

$$0 \rightarrow \mathcal{T}(D) \xrightarrow{\nabla} \mathcal{F}(D) \rightarrow \cdots .$$

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part IV: Cohomology

#### └ Tracial Cohomology

Let  $D$  be an anchored set. Denote the set of (globally defined) tracial functions on  $D$  by  $\mathcal{T}(D)$  and the set of free functions by  $\mathcal{F}(D)$ . For  $f \in \mathcal{T}(D)$ ,  $\nabla f \in \mathcal{F}(D)$ —so we have the beginnings of a chain complex!

$$0 \rightarrow \mathcal{T}(D) \xrightarrow{\nabla} \mathcal{F}(D) \rightarrow \cdots .$$

$$0 \rightarrow \mathcal{T}(D) \overset{\nabla}{\rightarrow} \mathcal{F}(D) \rightarrow \cdots .$$

2022-06-02

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Tracial Cohomology

$$0 \rightarrow \mathcal{T}(D) \overset{\nabla}{\rightarrow} \mathcal{F}(D) \rightarrow \cdots .$$

- test text

$$0 \rightarrow \mathcal{T}(D) \overset{\nabla}{\rightarrow} \mathcal{F}(D) \rightarrow \cdots .$$

A free function  $g : D \rightarrow \mathcal{M}^g$  is **exact** if there exists a tracial function  $f : D \rightarrow \mathbb{C}$  such that  $\nabla f = g$ .

2022-06-02

Searching for Holes in the Matrix Universe

└Part IV: Cohomology

└Tracial Cohomology

- test text

$$0 \rightarrow \mathcal{T}(D) \overset{\nabla}{\rightarrow} \mathcal{F}(D) \rightarrow \cdots .$$

A free function  $g : D \rightarrow \mathcal{M}^g$  is **exact** if there exists a tracial function  $f : D \rightarrow \mathbb{C}$  such that  $\nabla f = g$ .

$$0 \rightarrow \mathcal{T}(D) \overset{\nabla}{\rightarrow} \mathcal{F}(D) \rightarrow \cdots .$$

A free function  $g : D \rightarrow \mathcal{M}^g$  is **exact** if there exists a tracial function  $f : D \rightarrow \mathbb{C}$  such that  $\nabla f = g$ .

A free function  $g : D \rightarrow \mathcal{M}^g$  is **closed** if

$$\mathrm{tr} \left( K \cdot Dg(X)[H] \right) = \mathrm{tr} \left( H \cdot Dg(X)[K] \right)$$

for all directions  $H, K$ .

2022-06-02

Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Tracial Cohomology

- test text

$$0 \rightarrow \mathcal{T}(D) \overset{\nabla}{\rightarrow} \mathcal{F}(D) \rightarrow \cdots .$$

A free function  $g : D \rightarrow \mathcal{M}^g$  is **exact** if there exists a tracial function  $f : D \rightarrow \mathbb{C}$  such that  $\nabla f = g$ .

A free function  $g : D \rightarrow \mathcal{M}^g$  is **closed** if

$$\mathrm{tr} \left( K \cdot Dg(X)[H] \right) = \mathrm{tr} \left( H \cdot Dg(X)[K] \right)$$

for all directions  $H, K$ .

$$0 \rightarrow \mathcal{T}(D) \overset{\nabla}{\rightarrow} \mathcal{F}(D) \rightarrow \cdots .$$

A free function  $g : D \rightarrow \mathcal{M}^{\theta}$  is **exact** if there exists a tracial function  $f : D \rightarrow \mathbb{C}$  such that  $\nabla f = g$ .

A free function  $g : D \rightarrow \mathcal{M}^{\theta}$  is **closed** if

$$\mathrm{tr} \left( K \cdot Dg(X)[H] \right) = \mathrm{tr} \left( H \cdot Dg(X)[K] \right)$$

for all directions  $H, K$ .

**Definition:**

The **first tracial cohomology group** is the vector space of closed free functions moduluo the exact free function. We write  $H^1_{\mathrm{tr}}(D)$ .

2022-06-02

Searching for Holes in the Matrix Universe

- test text

$$0 \rightarrow \mathcal{T}(D) \overset{\nabla}{\rightarrow} \mathcal{F}(D) \rightarrow \cdots .$$

A free function  $g : D \rightarrow \mathcal{M}^{\theta}$  is **exact** if there exists a tracial function  $f : D \rightarrow \mathbb{C}$  such that  $\nabla f = g$ .

A free function  $g : D \rightarrow \mathcal{M}^{\theta}$  is **closed** if

$$\mathrm{tr} \left( K \cdot Dg(X)[H] \right) = \mathrm{tr} \left( H \cdot Dg(X)[K] \right)$$

for all directions  $H, K$ .

**Definition:**  
The **first tracial cohomology group** is the vector space of closed free functions moduluo the exact free function. We write  $H^1_{\mathrm{tr}}(D)$ .

# What out global germs?

- For  $f : B \rightarrow \mathbb{C}$  a global germ, since  $f$  analytically continues along every path, so does  $\nabla f$ .

2022-06-02

## Searching for Holes in the Matrix Universe

## └ Part IV: Cohomology

## └ Tracial Cohomology

## └ What out global germs?

- For  $f : B \rightarrow \mathbb{C}$  a global germ, since  $f$  analytically continues along every path, so does  $\nabla f$ .

# What out global germs?

- For  $f : B \rightarrow \mathbb{C}$  a global germ, since  $f$  analytically continues along every path, so does  $\nabla f$ .
- Free Monodromy means that  $\nabla f$  has a global extension.

2022-06-02

## Searching for Holes in the Matrix Universe

## └ Part IV: Cohomology

## └ Tracial Cohomology

## └ What out global germs?

- For  $f : B \rightarrow \mathbb{C}$  a global germ, since  $f$  analytically continues along every path, so does  $\nabla f$ .
- Free Monodromy means that  $\nabla f$  has a global extension.

# What out global germs?

- For  $f : B \rightarrow \mathbb{C}$  a global germ, since  $f$  analytically continues along every path, so does  $\nabla f$ .
- Free Monodromy means that  $\nabla f$  has a global extension.
- **Important:** If  $f$  is a global germ, then  $\nabla f$  is not necessarily exact since  $\mathcal{T}(D)$  is the set of tracial functions defined on *all of*  $D$ .

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part IV: Cohomology

#### └ Tracial Cohomology

#### └ What out global germs?

- For  $f : B \rightarrow \mathbb{C}$  a global germ, since  $f$  analytically continues along every path, so does  $\nabla f$ .
- Free Monodromy means that  $\nabla f$  has a global extension.
- **Important:** If  $f$  is a global germ, then  $\nabla f$  is not necessarily exact since  $\mathcal{T}(D)$  is the set of tracial functions defined on *all of*  $D$ .



**Goal:** Show that  $\pi_1^{\text{tr}}(D)$  injects into  $\mathbb{C}$ .

2022-06-02

**Goal:** Show that  $\pi_1^{\text{tr}}(D)$  injects into  $\mathbb{C}$ .

### Lemma

*Let  $D$  be an anchored nc domain. For any  $\alpha, \beta \in \pi_1^{\text{tr}}(D)$  and global germ  $f$ ,*

$$f(\alpha\beta) - f(\alpha) = f(\beta) - f(\tau)$$

*where  $\tau$  is the constant path.*

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part IV: Cohomology

#### └ Injecting into $\mathbb{C}$

**Goal:** Show that  $\pi_1^{\text{tr}}(D)$  injects into  $\mathbb{C}$ .

#### Lemma

*Let  $D$  be an anchored nc domain. For any  $\alpha, \beta \in \pi_1^{\text{tr}}(D)$  and global germ  $f$ ,*

$$f(\alpha\beta) - f(\alpha) = f(\beta) - f(\tau)$$

*where  $\tau$  is the constant path.*

For  $D$  an anchored set,  $X \in B_1$  the base point,  $f$  a global germ, and  $\gamma \in \pi_1^{\text{tr}}(D)$ , define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

2022-06-02

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

For  $D$  an anchored set,  $X \in B_1$  the base point,  $f$  a global germ, and  $\gamma \in \pi_1^{\text{tr}}(D)$ , define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

$c^f$  gives us a homomorphism into  $\mathbb{C}$ !

2022-06-02

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

For  $D$  an anchored set,  $X \in B_1$  the base point,  $f$  a global germ, and  $\gamma \in \pi_1^{\text{tr}}(D)$ , define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

$c^f$  gives us a homomorphism into  $\mathbb{C}$ !

$$c^f(\gamma_1\gamma_2) = f(\gamma_1\gamma_2) - f(\tau)$$

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

$$c^f(\gamma_1\gamma_2) = f(\gamma_1\gamma_2) - f(\tau)$$

For  $D$  an anchored set,  $X \in B_1$  the base point,  $f$  a global germ, and  $\gamma \in \pi_1^{\text{tr}}(D)$ , define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

$c^f$  gives us a homomorphism into  $\mathbb{C}$ !

$$\begin{aligned} c^f(\gamma_1\gamma_2) &= f(\gamma_1\gamma_2) - f(\tau) \\ &= f(\gamma_1\gamma_2) - f(\gamma_1) + f(\gamma_1) - f(\tau) \end{aligned}$$

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

$$\begin{aligned} c^f(\gamma_1\gamma_2) &= f(\gamma_1\gamma_2) - f(\tau) \\ &= f(\gamma_1\gamma_2) - f(\gamma_1) + f(\gamma_1) - f(\tau) \end{aligned}$$

For  $D$  an anchored set,  $X \in B_1$  the base point,  $f$  a global germ, and  $\gamma \in \pi_1^{\text{tr}}(D)$ , define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

$c^f$  gives us a homomorphism into  $\mathbb{C}$ !

$$\begin{aligned} c^f(\gamma_1\gamma_2) &= f(\gamma_1\gamma_2) - f(\tau) \\ &= f(\gamma_1\gamma_2) - f(\gamma_1) + f(\gamma_1) - f(\tau) \\ &= f(\gamma_2) - f(\tau) + f(\gamma_1) - f(\tau) \end{aligned}$$

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

$$\begin{aligned} c^f(\gamma_1\gamma_2) &= f(\gamma_1\gamma_2) - f(\tau) \\ &= f(\gamma_1\gamma_2) - f(\gamma_1) + f(\gamma_1) - f(\tau) \\ &= f(\gamma_2) - f(\tau) + f(\gamma_1) - f(\tau) \end{aligned}$$

For  $D$  an anchored set,  $X \in B_1$  the base point,  $f$  a global germ, and  $\gamma \in \pi_1^{\text{tr}}(D)$ , define

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

$c^f$  gives us a homomorphism into  $\mathbb{C}$ !

$$\begin{aligned} c^f(\gamma_1\gamma_2) &= f(\gamma_1\gamma_2) - f(\tau) \\ &= f(\gamma_1\gamma_2) - f(\gamma_1) + f(\gamma_1) - f(\tau) \\ &= f(\gamma_2) - f(\tau) + f(\gamma_1) - f(\tau) \\ &= c^f(\gamma_2) + c^f(\gamma_1). \end{aligned}$$

$$c^f(\gamma) = f(\gamma) - f(\tau)$$

$$\begin{aligned} c^f(\gamma_1\gamma_2) &= f(\gamma_1\gamma_2) - f(\tau) \\ &= f(\gamma_1\gamma_2) - f(\gamma_1) + f(\gamma_1) - f(\tau) \\ &= f(\gamma_2) - f(\tau) + f(\gamma_1) - f(\tau) \\ &= c^f(\gamma_2) + c^f(\gamma_1). \end{aligned}$$



## Lemma

*The map*

$$\Phi : \prod_{\substack{\nabla f \in H_{\text{tr}}^1(D) \\ f \text{ a global germ}}} \pi_1^{\text{tr}}(D) \longrightarrow \prod_{\substack{\nabla f \in H_{\text{tr}}^1(D) \\ f \text{ a global germ}}} \mathbb{C}$$

$$\prod \gamma \longmapsto \prod c^f(\gamma)$$

*is an injective homomorphism.*

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Injecting into  $\mathbb{C}$

**Lemma**

The map

$$\Phi : \prod_{\substack{\nabla f \in H_{\text{tr}}^1(D) \\ f \text{ a global germ}}} \pi_1^{\text{tr}}(D) \longrightarrow \prod_{\substack{\nabla f \in H_{\text{tr}}^1(D) \\ f \text{ a global germ}}} \mathbb{C}$$

$$\prod \gamma \longmapsto \prod c^f(\gamma)$$

is an injective homomorphism.

## Lemma

*The map*

$$\Phi : \prod_{\substack{\nabla f \in H_{\text{tr}}^1(D) \\ f \text{ a global germ}}} \pi_1^{\text{tr}}(D) \longrightarrow \prod_{\substack{\nabla f \in H_{\text{tr}}^1(D) \\ f \text{ a global germ}}} \mathbb{C}$$

$$\prod \gamma \longmapsto \prod c^f(\gamma)$$

*is an injective homomorphism.*

So  $\pi_1^{\text{tr}}(D)$  is commutative and torsion free!

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Injecting into  $\mathbb{C}$

**Lemma**

The map

$$\Phi : \prod_{\substack{\nabla f \in H_{\text{tr}}^1(D) \\ f \text{ a global germ}}} \pi_1^{\text{tr}}(D) \longrightarrow \prod_{\substack{\nabla f \in H_{\text{tr}}^1(D) \\ f \text{ a global germ}}} \mathbb{C}$$

$$\prod \gamma \longmapsto \prod c^f(\gamma)$$

is an injective homomorphism.

So  $\pi_1^{\text{tr}}(D)$  is commutative and torsion free!

$\pi_1^{\text{tr}}(D)$  is divisible

2022-06-02

$\pi_1^{\text{tr}}(D)$  is divisible

For any  $\gamma \in \pi_1^{\text{tr}}(D)$ ,

$$\gamma \oplus \tau = \tau \oplus \gamma.$$

2022-06-02

$$\gamma \oplus \tau = \tau \oplus \gamma.$$

# $\pi_1^{\text{tr}}(D)$ is divisible

For any  $\gamma \in \pi_1^{\text{tr}}(D)$ ,

$$\gamma \oplus \tau = \tau \oplus \gamma.$$

Why?

$$H(t, \theta) = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} (\gamma \oplus \gamma_X) \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}^*$$

is a homotopy between the paths.

2022-06-02

## Searching for Holes in the Matrix Universe

└ Part IV: Cohomology

└ Characterizing  $\pi_1^{\text{tr}}(D)$

└  $\pi_1^{\text{tr}}(D)$  is divisible

$\pi_1^{\text{tr}}(D)$  is divisible

For any  $\gamma \in \pi_1^{\text{tr}}(D)$ ,  
 $\gamma \oplus \tau = \tau \oplus \gamma.$

Why?

$$H(t, \theta) = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} (\gamma \oplus \gamma_X) \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}^*$$

is a homotopy between the paths.

$$\gamma = \underbrace{\begin{bmatrix} \gamma & & \\ & \gamma & \\ & & \ddots \\ & & & \gamma \end{bmatrix}}_{k+1\text{-times}}$$

2022-06-02

$$\gamma = \underbrace{\begin{bmatrix} \gamma & \\ & \gamma \end{bmatrix}}_{k+1\text{-times}}$$

$$\begin{aligned}\gamma &= \underbrace{\begin{bmatrix} \gamma & & \\ & \gamma & \\ & & \ddots \\ & & & \gamma \end{bmatrix}}_{k+1\text{-times}} \\ &= \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix} \begin{bmatrix} \tau & & \\ & \gamma & \\ & & \ddots \\ & & & \tau \end{bmatrix} \cdots \begin{bmatrix} \tau & & \\ & \tau & \\ & & \ddots \\ & & & \gamma \end{bmatrix}\end{aligned}$$

2022-06-02

$$\gamma = \underbrace{\begin{bmatrix} \gamma & \\ & \gamma \end{bmatrix}}_{k+1\text{-times}} = \begin{bmatrix} \gamma & \\ & \gamma \end{bmatrix} \cdots \begin{bmatrix} \gamma & \\ & \gamma \end{bmatrix}$$

$$\begin{aligned}\gamma &= \underbrace{\begin{bmatrix} \gamma & & \\ & \gamma & \\ & & \ddots \\ & & & \gamma \end{bmatrix}}_{k+1\text{-times}} \\ &= \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix} \begin{bmatrix} \tau & & \\ & \gamma & \\ & & \ddots \\ & & & \tau \end{bmatrix} \cdots \begin{bmatrix} \tau & & \\ & \tau & \\ & & \ddots \\ & & & \gamma \end{bmatrix} \\ &= \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \gamma_X \end{bmatrix} \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix} \cdots \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix}\end{aligned}$$

2022-06-02

$$\gamma = \underbrace{\begin{bmatrix} \gamma & \\ & \gamma \end{bmatrix}}_{k+1\text{-times}} = \begin{bmatrix} \gamma & \\ & \gamma \end{bmatrix} \begin{bmatrix} \gamma & \\ & \gamma \end{bmatrix} \cdots \begin{bmatrix} \gamma & \\ & \gamma \end{bmatrix}$$



$$\begin{aligned}\gamma &= \underbrace{\begin{bmatrix} \gamma & & \\ & \gamma & \\ & & \ddots \\ & & & \gamma \end{bmatrix}}_{k+1\text{-times}} \\ &= \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix} \begin{bmatrix} \tau & & \\ & \gamma & \\ & & \ddots \\ & & & \tau \end{bmatrix} \cdots \begin{bmatrix} \tau & & \\ & \tau & \\ & & \ddots \\ & & & \gamma \end{bmatrix} \\ &= \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \gamma_X \end{bmatrix} \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix} \cdots \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix} \\ &= \begin{bmatrix} \gamma & & \\ & \tau & \\ & & \ddots \\ & & & \tau \end{bmatrix}^k\end{aligned}$$

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part IV: Cohomology

#### └ Characterizing $\pi_1^{\text{tr}}(D)$

$$\gamma = \underbrace{\begin{bmatrix} \gamma & \\ & \gamma \end{bmatrix}}_{k+1\text{-times}} = \begin{bmatrix} \gamma & \\ & \gamma \end{bmatrix} \begin{bmatrix} \gamma & \\ & \gamma \end{bmatrix} \cdots \begin{bmatrix} \gamma & \\ & \gamma \end{bmatrix} = \begin{bmatrix} \gamma & \\ & \gamma \end{bmatrix}^k$$

## Theorem

*For  $D$  an anchored free set,  $\pi_1^{\text{tr}}(D)$  is a torsion free, abelian, divisible group. That is,*

$$\pi_1^{\text{tr}}(D) \simeq \bigoplus_{i \in I} \mathbb{Q} = \mathbb{Q}^I$$

*for some set  $I$ .*

2022-06-02

## Searching for Holes in the Matrix Universe

### └ Part IV: Cohomology

#### └ Characterizing $\pi_1^{\text{tr}}(D)$

#### Theorem

*For  $D$  an anchored free set,  $\pi_1^{\text{tr}}(D)$  is a torsion free, abelian, divisible group. That is,*

$$\pi_1^{\text{tr}}(D) \simeq \bigoplus_{i \in I} \mathbb{Q} = \mathbb{Q}^I$$

*for some set  $I$ .*

# Part V: Computing $\pi_1^{\text{tr}}(D)$

- hard bc no VC or MV

Let  $D$  be an anchored, free, path connected set such that each  $D_n$  is nonempty and choose an anchor  $B \subset D$  such that each  $B_n$  is also nonempty.

2022-06-02

Let  $D$  be an anchored, free, path connected set such that each  $D_n$  is nonempty and choose an anchor  $B \subset D$  such that each  $B_n$  is also nonempty.

Let  $D$  be an anchored, free, path connected set such that each  $D_n$  is nonempty and choose an anchor  $B \subset D$  such that each  $B_n$  is also nonempty.

Let  $\pi_1^{\text{tr}}(D)_n$  is the subgroup of paths in  $D_n$ . Note that  $\pi_1^{\text{tr}}(D)_1$  is a quotient of  $\pi_1(D_n)$ .

2022-06-02

Let  $D$  be an anchored, free, path connected set such that each  $D_n$  is nonempty and choose an anchor  $B \subset D$  such that each  $B_n$  is also nonempty.

Let  $\pi_1^{\text{tr}}(D)_n$  is the subgroup of paths in  $D_n$ . Note that  $\pi_1^{\text{tr}}(D)_1$  is a quotient of  $\pi_1(D_n)$ .

Let  $D$  be an anchored, free, path connected set such that each  $D_n$  is nonempty and choose an anchor  $B \subset D$  such that each  $B_n$  is also nonempty.

Let  $\pi_1^{\text{tr}}(D)_n$  is the subgroup of paths in  $D_n$ . Note that  $\pi_1^{\text{tr}}(D)_1$  is a quotient of  $\pi_1(D_n)$ .

There is a natural inclusion

$$\begin{aligned} \pi_1^{\text{tr}}(D)_n &\hookrightarrow \pi_1^{\text{tr}}(D)_{kn} \\ \gamma &\longmapsto \gamma^{\oplus k} \end{aligned}$$

for all  $k$ .

2022-06-02

Let  $D$  be an anchored, free, path connected set such that each  $D_n$  is nonempty and choose an anchor  $B \subset D$  such that each  $B_n$  is also nonempty.

Let  $\pi_1^{\text{tr}}(D)_n$  is the subgroup of paths in  $D_n$ . Note that  $\pi_1^{\text{tr}}(D)_1$  is a quotient of  $\pi_1(D_n)$ .

There is a natural inclusion

$$\begin{aligned} \pi_1^{\text{tr}}(D)_n &\hookrightarrow \pi_1^{\text{tr}}(D)_{kn} \\ \gamma &\longmapsto \gamma^{\oplus k} \end{aligned}$$

for all  $k$ .

Now consider the chain of inclusions:

$$\pi_1^{\text{tr}}(D)_1 \hookrightarrow \pi_1^{\text{tr}}(D)_2 \hookrightarrow \pi_1^{\text{tr}}(D)_6 \hookrightarrow \cdots \hookrightarrow \pi_1^{\text{tr}}(D)_{n!} \hookrightarrow \cdots$$

2022-06-02

Now consider the chain of inclusions:

$$\pi_1^{\text{tr}}(D)_1 \hookrightarrow \pi_1^{\text{tr}}(D)_2 \hookrightarrow \pi_1^{\text{tr}}(D)_6 \hookrightarrow \cdots \hookrightarrow \pi_1^{\text{tr}}(D)_{n!} \hookrightarrow \cdots$$

Now consider the chain of inclusions:

$$\pi_1^{\text{tr}}(D)_1 \hookrightarrow \pi_1^{\text{tr}}(D)_2 \hookrightarrow \pi_1^{\text{tr}}(D)_6 \hookrightarrow \cdots \hookrightarrow \pi_1^{\text{tr}}(D)_{n!} \hookrightarrow \cdots$$

The limit of this sequence isomorphic to  $\pi_1^{\text{tr}}(D)!$

Now consider the chain of inclusions:

$$\pi_1^{\text{tr}}(D)_1 \hookrightarrow \pi_1^{\text{tr}}(D)_2 \hookrightarrow \pi_1^{\text{tr}}(D)_6 \hookrightarrow \cdots \hookrightarrow \pi_1^{\text{tr}}(D)_{n!} \hookrightarrow \cdots$$

The limit of this sequence isomorphic to  $\pi_1^{\text{tr}}(D)!$



Example:  $\pi_1^{\text{tr}}(GL)$

Let  $GL = \bigcup_{n \in \mathbb{N}} GL_n(\mathbb{C})$ .

as well.

2022-06-02

Let  $GL = \bigcup_{n \in \mathbb{N}} GL_n(\mathbb{C})$ .

as well.

# Example: $\pi_1^{\text{tr}}(GL)$

Let  $GL = \bigcup_{n \in \mathbb{N}} GL_n(\mathbb{C})$ .

$GL_1(\mathbb{C}) = \mathbb{C} \setminus \{0\}$ . Since  $\pi_1(GL_1) \simeq \mathbb{Z}$ ,  $\pi_1^{\text{tr}}(GL)_1 \simeq \mathbb{Z}$  as well.

2022-06-02

## Searching for Holes in the Matrix Universe

└ Computing  $\pi_1^{\text{tr}}(D)$

└ An Example

└ Example:  $\pi_1^{\text{tr}}(GL)$

Let  $GL = \bigcup_{n \in \mathbb{N}} GL_n(\mathbb{C})$ .

$GL_1(\mathbb{C}) = \mathbb{C} \setminus \{0\}$ . Since  $\pi_1(GL_1) \simeq \mathbb{Z}$ ,  $\pi_1^{\text{tr}}(GL)_1 \simeq \mathbb{Z}$  as well.

Example:  $\pi_1^{\text{tr}}(GL)$

Let  $GL = \bigcup_{n \in \mathbb{N}} GL_n(\mathbb{C})$ .

$GL_1(\mathbb{C}) = \mathbb{C} \setminus \{0\}$ . Since  $\pi_1(GL_1) \simeq \mathbb{Z}$ ,  $\pi_1^{\text{tr}}(GL)_1 \simeq \mathbb{Z}$  as well.

Inclusion into  $\pi_1^{\text{tr}}(GL)_2$  picks up square roots. If  $\gamma \in \pi_1^{\text{tr}}(GL)_1$ , then

$$\begin{bmatrix} \gamma & \\ & \tau \end{bmatrix} \in \pi_1^{\text{tr}}(GL)_2.$$

2022-06-02

Searching for Holes in the Matrix Universe

└ Computing  $\pi_1^{\text{tr}}(D)$

└ An Example

└ Example:  $\pi_1^{\text{tr}}(GL)$

Let  $GL = \bigcup_{n \in \mathbb{N}} GL_n(\mathbb{C})$ .

$GL_1(\mathbb{C}) = \mathbb{C} \setminus \{0\}$ . Since  $\pi_1(GL_1) \simeq \mathbb{Z}$ ,  $\pi_1^{\text{tr}}(GL)_1 \simeq \mathbb{Z}$  as well.

Inclusion into  $\pi_1^{\text{tr}}(GL)_2$  picks up square roots. If  $\gamma \in \pi_1^{\text{tr}}(GL)_1$ , then

$$\begin{bmatrix} \gamma & \\ & \tau \end{bmatrix} \in \pi_1^{\text{tr}}(GL)_2.$$

# Example: $\pi_1^{\text{tr}}(GL)$

Let  $GL = \bigcup_{n \in \mathbb{N}} GL_n(\mathbb{C})$ .

$GL_1(\mathbb{C}) = \mathbb{C} \setminus \{0\}$ . Since  $\pi_1(GL_1) \simeq \mathbb{Z}$ ,  $\pi_1^{\text{tr}}(GL)_1 \simeq \mathbb{Z}$  as well.

Inclusion into  $\pi_1^{\text{tr}}(GL)_2$  picks up square roots. If  $\gamma \in \pi_1^{\text{tr}}(GL)_1$ , then

$$\begin{bmatrix} \gamma & \\ & \tau \end{bmatrix} \in \pi_1^{\text{tr}}(GL)_2.$$

Thus,  $\pi_1^{\text{tr}}(GL)_2 \simeq \mathbb{Z} \left[ \frac{1}{2} \right]$ .

2022-06-02

## Searching for Holes in the Matrix Universe

└ Computing  $\pi_1^{\text{tr}}(D)$

└ An Example

└ Example:  $\pi_1^{\text{tr}}(GL)$

Let  $GL = \bigcup_{n \in \mathbb{N}} GL_n(\mathbb{C})$ .

$GL_1(\mathbb{C}) = \mathbb{C} \setminus \{0\}$ . Since  $\pi_1(GL_1) \simeq \mathbb{Z}$ ,  $\pi_1^{\text{tr}}(GL)_1 \simeq \mathbb{Z}$  as well.

Inclusion into  $\pi_1^{\text{tr}}(GL)_2$  picks up square roots. If  $\gamma \in \pi_1^{\text{tr}}(GL)_1$ , then

$$\begin{bmatrix} \gamma & \\ & \tau \end{bmatrix} \in \pi_1^{\text{tr}}(GL)_2.$$

Thus,  $\pi_1^{\text{tr}}(GL)_2 \simeq \mathbb{Z} \left[ \frac{1}{2} \right]$ .

Similarly, inclusion into  $\pi_1^{\text{tr}}(GL)_{3!}$  picks up cube roots:

$$\pi_1^{\text{tr}}(GL)_6 \simeq \mathbb{Z} \left[ \frac{1}{2}, \frac{1}{3} \right]$$

$$\pi_1^{\text{tr}}(GL)_6 \simeq \mathbb{Z} \left[ \frac{1}{2}, \frac{1}{3} \right]$$

Similarly, inclusion into  $\pi_1^{\text{tr}}(GL)_{3!}$  picks up cube roots:

$$\pi_1^{\text{tr}}(GL)_6 \simeq \mathbb{Z} \left[ \frac{1}{2}, \frac{1}{3} \right]$$

In the  $n$ -th inclusion, we pick up  $n$ -th roots and so we adjoin  $\frac{1}{n}$  to the preceding group. Therefore,

$$\pi_1^{\text{tr}}(GL) \simeq \mathbb{Z} \left[ \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots \right]$$

Similarly, inclusion into  $\pi_1^{\text{tr}}(GL)_6$  picks up cube roots:

$$\pi_1^{\text{tr}}(GL)_6 \simeq \mathbb{Z} \left[ \frac{1}{2}, \frac{1}{3} \right]$$

In the  $n$ -th inclusion, we pick up  $n$ -th roots and so we adjoin  $\frac{1}{n}$  to the preceding group. Therefore,

$$\pi_1^{\text{tr}}(GL) \simeq \mathbb{Z} \left[ \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots \right]$$

Similarly, inclusion into  $\pi_1^{\text{tr}}(GL)_{3!}$  picks up cube roots:

$$\pi_1^{\text{tr}}(GL)_6 \simeq \mathbb{Z} \left[ \frac{1}{2}, \frac{1}{3} \right]$$

In the  $n$ -th inclusion, we pick up  $n$ -th roots and so we adjoin  $\frac{1}{n}$  to the preceding group. Therefore,

$$\pi_1^{\text{tr}}(GL) \simeq \mathbb{Z} \left[ \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots \right] \simeq \mathbb{Q}.$$

Similarly, inclusion into  $\pi_1^{\text{tr}}(GL)_6$  picks up cube roots:

$$\pi_1^{\text{tr}}(GL)_6 \simeq \mathbb{Z} \left[ \frac{1}{2}, \frac{1}{3} \right]$$

In the  $n$ -th inclusion, we pick up  $n$ -th roots and so we adjoin  $\frac{1}{n}$  to the preceding group. Therefore,

$$\pi_1^{\text{tr}}(GL) \simeq \mathbb{Z} \left[ \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots \right] \simeq \mathbb{Q}.$$

Thank You!

2022-06-02