

# Probability and Random Variables

# Concepts

1. Utility and Probability
2. Random Variables
3. Relationships between Random Variables

# Utility and Probability

Full story: <https://projecteuclid.org/journals/statistical-science/volume-1/issue-3/The-Axioms-of-Subjective-Probability/10.1214/ss/1177013611.full>

# What is a Random Variable?

R.V.  $X$

# Vocabulary/Notation

# Vocabulary/Notation

Term	Definition	Coinflip Example	Uniform Example
------	------------	------------------	-----------------

# Vocabulary/Notation

Bernoulli(0.5)

**Term**

**Definition**

**Coinflip Example**

**Uniform Example**

# Vocabulary/Notation

		Bernoulli(0.5)	$\mathcal{U}(0, 1)$
Term	Definition	Coinflip Example	Uniform Example



# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
support( $X$ )			

# Vocabulary/Notation

Term	Definition	Bernoulli(0.5)	$\mathcal{U}(0, 1)$
Coinflip Example	Uniform Example		
support( $X$ )	All the values that $X$ can take		

# Vocabulary/Notation

Term	Definition	Bernoulli(0.5)	$\mathcal{U}(0, 1)$
Coinflip Example	Uniform Example		
support( $X$ ) $x \in X$	All the values that $X$ can take		

# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
support( $X$ )	All the values that $X$ can take		
$x \in X$			
$X \in [0, 1]$			

# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
support( $X$ ) $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	

# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
support( $X$ ) $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$

# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
support( $X$ ) $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$
Distribution			

# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
support( $X$ ) $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$

## Distribution

- Discrete: PMF
- Continuous: PDF



# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
support( $X$ ) $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$
Distribution <ul style="list-style-type: none"><li>• Discrete: PMF</li><li>• Continuous: PDF</li></ul>	Maps each value in the support to a real number indicating its probability		

# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
$\text{support}(X)$ $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$
Distribution <ul style="list-style-type: none"><li>• Discrete: PMF</li><li>• Continuous: PDF</li></ul>	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$	


# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support( $X$ ) $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution • Discrete: PMF • Continuous: PDF	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <table><tr><td>x</td><td>P(x)</td></tr><tr><td>0</td><td>0.5</td></tr><tr><td>1</td><td>0.5</td></tr></table>	x	P(x)	0	0.5	1	0.5	
x	P(x)								
0	0.5								
1	0.5								


# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support( $X$ ) $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution <ul style="list-style-type: none"><li>• Discrete: PMF</li><li>• Continuous: PDF</li></ul>	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <table><tr><td>x</td><td>P(x)</td></tr><tr><td>0</td><td>0.5</td></tr><tr><td>1</td><td>0.5</td></tr></table>	x	P(x)	0	0.5	1	0.5	$p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$
x	P(x)								
0	0.5								
1	0.5								

# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support( $X$ ) $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution <ul style="list-style-type: none"><li>Discrete: PMF</li><li>Continuous: PDF</li></ul>	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <table><tr><td>x</td><td>P(x)</td></tr><tr><td>0</td><td>0.5</td></tr><tr><td>1</td><td>0.5</td></tr></table>	x	P(x)	0	0.5	1	0.5	$p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ 
x	P(x)								
0	0.5								
1	0.5								


# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
<b>support(<math>X</math>)</b> $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$ 						
<b>Distribution</b> <ul style="list-style-type: none"> <li>Discrete: PMF</li> <li>Continuous: PDF</li> </ul>	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <table border="1" data-bbox="1335 952 1569 1128"> <thead> <tr> <th>x</th> <th>P(x)</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>0.5</td> </tr> <tr> <td>1</td> <td>0.5</td> </tr> </tbody> </table>	x	P(x)	0	0.5	1	0.5	$p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) =$
x	P(x)								
0	0.5								
1	0.5								

# Vocabulary/Notation


Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support( $X$ ) $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution <ul style="list-style-type: none"><li>Discrete: PMF</li><li>Continuous: PDF</li></ul>	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <table><tr><th>x</th><th>P(x)</th></tr><tr><td>0</td><td>0.5</td></tr><tr><td>1</td><td>0.5</td></tr></table>	x	P(x)	0	0.5	1	0.5	$p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ 
x	P(x)								
0	0.5								
1	0.5								

# Vocabulary/Notation


Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support( $X$ ) $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution <ul style="list-style-type: none"><li>Discrete: PMF</li><li>Continuous: PDF</li></ul>	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <table><tr><th>x</th><th>P(x)</th></tr><tr><td>0</td><td>0.5</td></tr><tr><td>1</td><td>0.5</td></tr></table>	x	P(x)	0	0.5	1	0.5	 $p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ $P(X \in [a, b]) = \int_a^b p(x)dx$
x	P(x)								
0	0.5								
1	0.5								




# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
<b>support(<math>X</math>)</b> $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
<b>Distribution</b> <ul style="list-style-type: none"> <li>Discrete: PMF</li> <li>Continuous: PDF</li> </ul>	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <div> <table> <tr> <th>x</th> <th>P(x)</th> </tr> <tr> <td>0</td> <td>0.5</td> </tr> <tr> <td>1</td> <td>0.5</td> </tr> </table> </div>	x	P(x)	0	0.5	1	0.5	 $p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ $P(X \in [a, b]) = \int_a^b p(x) dx$
x	P(x)								
0	0.5								
1	0.5								
<b>Expectation</b>									


# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support( $X$ ) $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$ 						
Distribution <ul style="list-style-type: none"><li>Discrete: PMF</li><li>Continuous: PDF</li></ul>	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <table data-bbox="1335 948 1572 1133"><tr><th>x</th><th>P(x)</th></tr><tr><td>0</td><td>0.5</td></tr><tr><td>1</td><td>0.5</td></tr></table>	x	P(x)	0	0.5	1	0.5	$p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ $P(X \in [a, b]) = \int_a^b p(x)dx$
x	P(x)								
0	0.5								
1	0.5								
Expectation	First moment of the random variable, "mean"								


# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
<b>support(<math>X</math>)</b> $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
<b>Distribution</b> <ul style="list-style-type: none"> <li>Discrete: PMF</li> <li>Continuous: PDF</li> </ul>	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <div> <table> <tr> <th>x</th> <th>P(x)</th> </tr> <tr> <td>0</td> <td>0.5</td> </tr> <tr> <td>1</td> <td>0.5</td> </tr> </table> </div>	x	P(x)	0	0.5	1	0.5	 $p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ $P(X \in [a, b]) = \int_a^b p(x) dx$
x	P(x)								
0	0.5								
1	0.5								
<b>Expectation</b> $E[X]$	First moment of the random variable, "mean"								


# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
<b>support(<math>X</math>)</b> $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
<b>Distribution</b> <ul style="list-style-type: none"> <li>Discrete: PMF</li> <li>Continuous: PDF</li> </ul>	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <div> <table> <tr> <th>x</th> <th>P(x)</th> </tr> <tr> <td>0</td> <td>0.5</td> </tr> <tr> <td>1</td> <td>0.5</td> </tr> </table> </div>	x	P(x)	0	0.5	1	0.5	 $p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ $P(X \in [a, b]) = \int_a^b p(x) dx$
x	P(x)								
0	0.5								
1	0.5								
<b>Expectation</b> $E[X]$	First moment of the random variable, "mean"	$E[X] = \sum_{x \in X} x P(x)$							


# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
<b>support(<math>X</math>)</b> $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
<b>Distribution</b> <ul style="list-style-type: none"> <li>Discrete: PMF</li> <li>Continuous: PDF</li> </ul>	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <div> <table> <tr> <th>x</th> <th>P(x)</th> </tr> <tr> <td>0</td> <td>0.5</td> </tr> <tr> <td>1</td> <td>0.5</td> </tr> </table> </div>	x	P(x)	0	0.5	1	0.5	 $p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ $P(X \in [a, b]) = \int_a^b p(x)dx$
x	P(x)								
0	0.5								
1	0.5								
<b>Expectation</b> $E[X]$	First moment of the random variable, "mean"	$E[X] = \sum_{x \in X} xP(x)$ $= 0.5$							

# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
<b>support(<math>X</math>)</b> $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
<b>Distribution</b> <ul style="list-style-type: none"> <li>Discrete: PMF</li> <li>Continuous: PDF</li> </ul>	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <div> <table> <tr> <th>x</th> <th>P(x)</th> </tr> <tr> <td>0</td> <td>0.5</td> </tr> <tr> <td>1</td> <td>0.5</td> </tr> </table> </div>	x	P(x)	0	0.5	1	0.5	 $p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ $P(X \in [a, b]) = \int_a^b p(x) dx$
x	P(x)								
0	0.5								
1	0.5								
<b>Expectation</b> $E[X]$	First moment of the random variable, "mean"	$E[X] = \sum_{x \in X} xP(x)$ $= 0.5$	$E[X] = \int_{x \in X} xp(x) dx$						

# Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
<b>support(<math>X</math>)</b> $x \in X$ $X \in [0, 1]$	All the values that $X$ can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
<b>Distribution</b> <ul style="list-style-type: none"> <li>Discrete: PMF</li> <li>Continuous: PDF</li> </ul>	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <div> <table> <tr> <th>x</th> <th>P(x)</th> </tr> <tr> <td>0</td> <td>0.5</td> </tr> <tr> <td>1</td> <td>0.5</td> </tr> </table> </div>	x	P(x)	0	0.5	1	0.5	 $p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ $P(X \in [a, b]) = \int_a^b p(x) dx$
x	P(x)								
0	0.5								
1	0.5								
<b>Expectation</b> $E[X]$	First moment of the random variable, "mean"	$E[X] = \sum_{x \in X} xP(x)$ $= 0.5$	$E[X] = \int_{x \in X} xp(x) dx$ $= 0.5$						

# Distributions of related R.V.s



# Distributions of related R.V.s

**Joint Distribution**

# Distributions of related R.V.s

## Joint Distribution

$$P(X, Y, Z)$$

# Distributions of related R.V.s

## Joint Distribution

$$P(X, Y, Z)$$

$X$	$Y$	$Z$	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

# Distributions of related R.V.s

## Joint Distribution

$$P(X, Y, Z)$$

$X$	$Y$	$Z$	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

## Conditional Distribution

# Distributions of related R.V.s

## Joint Distribution

$$P(X, Y, Z)$$

$X$	$Y$	$Z$	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

## Conditional Distribution

$$P(X \mid Y, Z)$$

# Distributions of related R.V.s

## Joint Distribution

$$P(X, Y, Z)$$

$X$	$Y$	$Z$	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

## Conditional Distribution

$$P(X | Y, Z)$$

(Distribution - valued function)

$X$	$P(X   Y=1, Z=1)$
0	0.84
1	0.16

# Distributions of related R.V.s

## Joint Distribution

$$P(X, Y, Z)$$

$X$	$Y$	$Z$	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

## Conditional Distribution

$$P(X | Y, Z)$$

(Distribution - valued function)

$X$	$P(X   Y=1, Z=1)$
0	0.84
1	0.16

## Marginal Distribution

# Distributions of related R.V.s

## Joint Distribution

$$P(X, Y, Z)$$

$X$	$Y$	$Z$	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

## Conditional Distribution

$$P(X \mid Y, Z)$$

(Distribution - valued function)

$X$	$P(X \mid Y=1, Z=1)$
0	0.84
1	0.16

## Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$



# Distributions of related R.V.s

## Joint Distribution

$$P(X, Y, Z)$$

$X$	$Y$	$Z$	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

## Conditional Distribution

$$P(X | Y, Z)$$

(Distribution - valued function)

$X$	$P(X   Y=1, Z=1)$
0	0.84
1	0.16

## Marginal Distribution

$$P(X) \quad P(Y) \quad P(Z)$$

$X$	$P(X)$	$Y$	$P(Y)$
0	0.85	0	0.45
1	0.15	1	0.55

$Z$	$P(Z)$
0	0.20
1	0.80

# Distributions of related R.V.s

**Joint Distribution**

$$P(X, Y, Z)$$

**Conditional Distribution**

$$P(X \mid Y, Z)$$

**Marginal Distribution**

$$P(X) \ P(Y) \ P(Z)$$

# Distributions of related R.V.s

**Joint Distribution**

$$P(X, Y, Z)$$

**Conditional Distribution**

$$P(X \mid Y, Z)$$

**Marginal Distribution**

$$P(X) \ P(Y) \ P(Z)$$

**3 Rules**

# Distributions of related R.V.s

**Joint Distribution**

$$P(X, Y, Z)$$

**Conditional Distribution**

$$P(X \mid Y, Z)$$

**Marginal Distribution**

$$P(X) \ P(Y) \ P(Z)$$

**3 Rules** (Burrito-level)

# Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules

(Burrito-level)

(Blackbelly Level: Axioms of Probability)

**AXIOM 1. STRUCTURE OF UNKNOWN REAL NUMBERS AND PLAUSIBLE VALUE.** We assume a set  $T$  of unknown numbers is a partially ordered commutative algebra over  $\mathbb{R}$  with identity, 1.

We assume in addition a given sub-Boolean algebra  $E$  of  $E(T)$  with  $0, 1 \in E$  and denote by  $E_0$  the set of non-zero members of  $E$ . We assume that the partial ordering in  $E(T)$  as a Boolean algebra coincides with the ordering that  $E(T)$  inherits from the algebra  $T$ . Finally, we assume a function  $PV : T \times E_0 \rightarrow \mathbb{R}$ , called **PLAUSIBLE VALUE**, whose value on the pair  $(x, e)$  is denoted  $PV(x|e)$ .

not on exam

**AXIOM 2. STRONG RESCALING FOR PLAUSIBLE VALUE.** If  $a, b$  belong to  $\mathbb{R}$ , if  $x$  belongs to  $T$ , and if  $e$  belongs to  $E_0$ , then

$$PV(ax + b|e) = aPV(x|e) + b. \quad (2)$$

**AXIOM 3. ORDER CONSISTENCY FOR PLAUSIBLE VALUE.** If  $x, y \in T$  and if  $e \in E_0$ , implies that  $x \leq y$ , then  $PV(x|e) \leq PV(y|e)$ .

Notice that if  $e \in E(T)$ , then  $0 \leq e \leq 1$ , in  $T$ , as it is true in the lattice ordering of  $E(T)$ .

**AXIOM 4. THE COX AXIOM FOR PLAUSIBLE VALUE:** If  $e, c$  are fixed in  $E$ , with  $ec \in E_0$ , if  $x_1, x_2$  are in  $T$ , if  $PV(x_1|ec) = PV(x_2|ec)$ , then  $PV(x_1|c) = PV(x_2|c)$ . That is, we assume that as a function of  $x$ , the plausible value  $PV(x|c)$  depends only on  $PV(x|ec)$ .

**AXIOM 5. RESTRICTED ADDITIVITY OF PLAUSIBLE VALUE.** For each fixed  $y \in T$  and  $e \in E_0$ , the plausible value  $PV(x + y|e)$  as a function of  $x \in T$  depends only on  $PV(x|e)$ , which is to say that if  $x_1, x_2 \in T$  and  $PV(x_1|e) = PV(x_2|e)$ , then  $PV(x_1 + y|e) = PV(x_2 + y|e)$ .

# Distributions of related R.V.s

**Joint Distribution**

$$P(X, Y, Z)$$

**Conditional Distribution**

$$P(X \mid Y, Z)$$

**Marginal Distribution**

$$P(X) \ P(Y) \ P(Z)$$

**3 Rules** (Burrito-level)

1)

# Distributions of related R.V.s

**Joint Distribution**

$$P(X, Y, Z)$$

**Conditional Distribution**

$$P(X \mid Y, Z)$$

**Marginal Distribution**

$$P(X) \ P(Y) \ P(Z)$$

**3 Rules** (Burrito-level)

1) a)  $0 \leq P(X \mid Y) \leq 1$

# Distributions of related R.V.s

**Joint Distribution**

$$P(X, Y, Z)$$

**Conditional Distribution**

$$P(X \mid Y, Z)$$

**Marginal Distribution**

$$P(X) \ P(Y) \ P(Z)$$

**3 Rules** (Burrito-level)

- 1) a)  $0 \leq P(X \mid Y) \leq 1$   
b)  $\sum_{x \in X} P(x \mid Y) = 1$



# Distributions of related R.V.s

**Joint Distribution**

$$P(X, Y, Z)$$

**Conditional Distribution**

$$P(X \mid Y, Z)$$

**Marginal Distribution**

$$P(X) \ P(Y) \ P(Z)$$

**3 Rules** (Burrito-level)

- 1) a)  $0 \leq P(X \mid Y) \leq 1$   
b)  $\sum_{x \in X} P(x \mid Y) = 1$

- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

# Distributions of related R.V.s

**Joint Distribution**

$$P(X, Y, Z)$$

**Conditional Distribution**

$$P(X \mid Y, Z)$$

**Marginal Distribution**

$$P(X) \ P(Y) \ P(Z)$$

**3 Rules** (Burrito-level)

- 1) a)  $0 \leq P(X \mid Y) \leq 1$   
b)  $\sum_{x \in X} P(x \mid Y) = 1$

- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

Joint  $\rightarrow$  Marginal

# Distributions of related R.V.s

**Joint Distribution**

$$P(X, Y, Z)$$

**Conditional Distribution**

$$P(X \mid Y, Z)$$

**Marginal Distribution**

$$P(X) \ P(Y) \ P(Z)$$

## 3 Rules (Burrito-level)

- 1) a)  $0 \leq P(X \mid Y) \leq 1$   
b)  $\sum_{x \in X} P(x \mid Y) = 1$

- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

- 3) Definition of Conditional Probability

$$P(X \mid Y) = \frac{P(X, Y)}{P(Y)}$$

Joint  $\rightarrow$  Marginal

# Distributions of related R.V.s

**Joint Distribution**

$$P(X, Y, Z)$$

**Conditional Distribution**

$$P(X \mid Y, Z)$$

**Marginal Distribution**

$$P(X) P(Y) P(Z)$$

## 3 Rules (Burrito-level)

- 1) a)  $0 \leq P(X \mid Y) \leq 1$   
b)  $\sum_{x \in X} P(x \mid Y) = 1$

- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

- 3) Definition of Conditional Probability

$$P(X \mid Y) = \frac{P(X, Y)}{P(Y)}$$

Joint  $\rightarrow$  Marginal

Joint + Marginal  $\rightarrow$  Conditional

# Distributions of related R.V.s

**Joint Distribution**

$$P(X, Y, Z)$$

**Conditional Distribution**

$$P(X \mid Y, Z)$$

**Marginal Distribution**

$$P(X) \ P(Y) \ P(Z)$$

## 3 Rules (Burrito-level)

- 1) a)  $0 \leq P(X \mid Y) \leq 1$   
b)  $\sum_{x \in X} P(x \mid Y) = 1$

- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

- 3) Definition of Conditional Probability

$$P(X \mid Y) = \frac{P(X, Y)}{P(Y)}$$

Joint  $\rightarrow$  Marginal

Joint + Marginal  $\rightarrow$  Conditional

Marginal + Conditional  $\rightarrow$  Joint

$$P(X, Y) = P(X \mid Y) P(Y)$$

# Distributions of related R.V.s

**Joint Distribution**

$$P(X, Y, Z)$$

**Conditional Distribution**

$$P(X \mid Y, Z)$$

**Marginal Distribution**

$$P(X) \ P(Y) \ P(Z)$$

## 3 Rules

- 1) a)  $0 \leq P(X \mid Y) \leq 1$   
 b)  $\sum_{x \in X} P(x \mid Y) = 1$

X	Y	Z	P(X, Y, Z)
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

- 3) Definition of Conditional Probability

$$P(X \mid Y) = \frac{P(X, Y)}{P(Y)}$$

Joint  $\rightarrow$  Marginal

Joint + Marginal  $\rightarrow$  Conditional

Marginal + Conditional  $\rightarrow$  Joint

$$P(X, Y) = P(X \mid Y) P(Y)$$

# Break

1) a)  $0 \leq P(X | Y) \leq 1$

b)  $\sum_{x \in X} P(x | Y) = 1$

2)  $P(X) = \sum_{y \in Y} P(X, y)$

3)  $P(X | Y) = \frac{P(X, Y)}{P(Y)}$

$$P(X, Y) = P(X|Y) P(Y)$$

- $P \in \{0, 1\}$ : Powder Day
  - $C \in \{0, 1\}$ : Pass Clear
  - 1 in 5 days is a powder day
  - The pass is clear 8 in 10 days
  - If it is a powder day, there is a 50% chance the pass is blocked
- 
- Write out the joint probability distribution for P and C.
  - Suppose it is a non-powder day, what is the probability that the pass is blocked?

# Bayes Rule

- Know:  $P(B \mid A), P(A), P(B)$
- Want:  $P(A \mid B)$



# Independence

# Independence

Definition:  $X$  and  $Y$  are *independent* iff  $P(X, Y) = P(X) P(Y)$

# Independence

Definition:  $X$  and  $Y$  are *independent* iff  $P(X, Y) = P(X) P(Y)$

$$X \perp Y$$

# Independence

Definition:  $X$  and  $Y$  are *independent* iff  $P(X, Y) = P(X) P(Y)$

$$X \perp Y$$

$$P(X|Y) = P(X)$$

# Independence

Definition:  $X$  and  $Y$  are *independent* iff  $P(X, Y) = P(X) P(Y)$

$$X \perp Y$$

$$P(X|Y) = P(X)$$

Definition:  $X$  and  $Y$  are *conditionally independent* given  $Z$  iff

$$P(X, Y | Z) = P(X | Z) P(Y | Z)$$

# Independence

Definition:  $X$  and  $Y$  are *independent* iff  $P(X, Y) = P(X) P(Y)$

$$X \perp Y$$

$$P(X|Y) = P(X)$$

Definition:  $X$  and  $Y$  are *conditionally independent* given  $Z$  iff

$$P(X, Y | Z) = P(X | Z) P(Y | Z)$$

$$X \perp Y | Z$$

# Rules for Continuous RVs

## Discrete

1) a)  $0 \leq P(X | Y) \leq 1$

b)  $\sum_{x \in X} P(x | Y) = 1$

2)  $P(X) = \sum_{y \in Y} P(X, y)$

3)  $P(X | Y) = \frac{P(X, Y)}{P(Y)}$

$$P(X, Y) = P(X | Y) P(Y)$$

## Continuous

1)

# Rules for Continuous RVs

## Discrete

1) a)  $0 \leq P(X | Y) \leq 1$

b)  $\sum_{x \in X} P(x | Y) = 1$

2)  $P(X) = \sum_{y \in Y} P(X, y)$

3)  $P(X | Y) = \frac{P(X, Y)}{P(Y)}$

$$P(X, Y) = P(X | Y) P(Y)$$

## Continuous

1)  $0 \leq p(X | Y)$



# Rules for Continuous RVs

## Discrete

1) a)  $0 \leq P(X | Y) \leq 1$   
b)  $\sum_{x \in X} P(x | Y) = 1$

2)  $P(X) = \sum_{y \in Y} P(X, y)$

3)  $P(X | Y) = \frac{P(X, Y)}{P(Y)}$

$$P(X, Y) = P(X | Y) P(Y)$$

## Continuous

1)  $0 \leq p(X | Y)$   
 $\int_X p(x|Y) dx = 1$

# Rules for Continuous RVs

## Discrete

1) a)  $0 \leq P(X | Y) \leq 1$

b)  $\sum_{x \in X} P(x | Y) = 1$

2)  $P(X) = \sum_{y \in Y} P(X, y)$

3)  $P(X | Y) = \frac{P(X, Y)}{P(Y)}$

$$P(X, Y) = P(X | Y) P(Y)$$

## Continuous

1)  $0 \leq p(X | Y)$

$$\int_X p(x|Y) dx = 1$$

2)

$$p(X) = \int_Y p(X, y) dy$$

# Rules for Continuous RVs

## Discrete

1) a)  $0 \leq P(X | Y) \leq 1$

b)  $\sum_{x \in X} P(x | Y) = 1$

2)  $P(X) = \sum_{y \in Y} P(X, y)$

3)  $P(X | Y) = \frac{P(X, Y)}{P(Y)}$

$$P(X, Y) = P(X | Y) P(Y)$$

## Continuous

1)  $0 \leq p(X | Y)$

$$\int_X p(x|Y) dx = 1$$

2)

$$p(X) = \int_Y p(X, y) dy$$

3)  $p(X | Y) = \frac{p(X, Y)}{p(Y)}$

$$p(X, Y) = p(X | Y) p(Y)$$

# Multivariate Gaussian Distribution

**Joint Distribution**

**Conditional Distribution**

**Marginal Distribution**

# Concepts

1. Utility and Probability
2. Random Variables
3. Relationships between Random Variables