Bayes' rule: update what you believe, using data

$$p(A \mid B) = \frac{p(B \mid A)\, p(A)}{p(B)}$$

$$\underbrace{p(A \mid B)}_{\text{posterior}} = \frac{\overbrace{p(B \mid A)}^{\text{likelihood}}\ \overbrace{p(A)}^{\text{prior}}}{\underbrace{p(B)}_{\text{evidence}}}$$

> Four names for the whole lecture: posterior, likelihood, prior and evidence.

In machine learning, the unknown is the parameters

$$p(\theta \mid \mathcal{D}) = \frac{p(\mathcal{D} \mid \theta)\, p(\theta)}{p(\mathcal{D})}$$

Usual training


keeps one θ: the best fit to the data $\mathcal{D}$.

Bayesian learning

keeps every θ, weighted by $p(\theta \mid \mathcal{D})$.

> Usual training keeps one θ. The Bayesian answer keeps a whole distribution over θ.

A tiny network: one input, two hidden units, one output



$$f_\theta(x) = v_1 \tanh(w_1 x + b_1) + v_2 \tanh(w_2 x + b_2) + c$$

$$\theta = (w_1, w_2, b_1, b_2, v_1, v_2, c) \in \mathbb{R}^7$$


Seven unknown numbers: four weights and three biases.

> Every weight and bias is unknown, so θ is seven numbers.

A prior on every parameter

$$\theta_j \sim \mathcal{N}(0, 1), \quad j = 1, \ldots, 7$$

$$p(\theta) = \prod_{j=1}^{7} \mathcal{N}(\theta_j;\, 0,\, 1)$$




One draw from the prior is seven numbers, one from each bell: a single point θ in $\mathbb{R}^7$.

One draw of θ is one whole function

	w_1	w_2	b_1	b_2	v_1	v_2	c
$\theta^{(1)}$	0.79	1.39	−0.15	−0.02	0.28	−1.10	−1.03
$\theta^{(2)}$	−1.07	−1.40	0.04	0.78	0.45	−1.93	0.77
$\theta^{(3)}$	−1.06	0.20	−0.63	0.22	0.48	−0.56	1.66


Each row, put through the network, is one curve.




> Every point of $\mathbb{R}^7$ is a whole function. Drawing θ from the prior draws functions.

Twenty draws, and the prior predictive

20 draws, and the prior predictive: mean ± 2 sd




draw 1 of 20: $\theta^{(1)} \in \mathbb{R}^7$



Hover over a line, or step through the draws.

Average over every θ and add the 0.1 noise: the prior predictive, $p(y \mid x) = \int p(y \mid x, \theta)\, p(\theta)\, d\theta$. What the model expects before any data.

The data, and how well one θ explains them



$$y_i \mid x_i, \theta \sim \mathcal{N}\big(f_\theta(x_i),\, 0.1^2\big)$$

$$p(\mathcal{D} \mid \theta) = \prod_{i=1}^{17} \mathcal{N}\big(y_i;\, f_\theta(x_i),\, 0.1^2\big)$$


> The likelihood of θ multiplies all seventeen densities: how well this one θ explains the whole dataset.

Meet θ_A: one good setting of the seven numbers

$\theta_A \in \mathbb{R}^7$: the most probable θ




the network with θ_A, and the data



θ_A is the most probable θ: gradient ascent on $\log p(\mathcal{D} \mid \theta) + \log p(\theta)$, the best of ten random starts. For now it is just one candidate.

> Put it through the network: the curve follows the data. How likely does θ_A make the data, and how likely is θ_A under the prior?

The likelihood of θ_A, one data point at a time



Do that for all seventeen points and add: log p($\mathcal{D}$ | θ_A) = 12.17.

The prior of θ_A, then the evidence


$$\log \mathcal{N}(\theta_j; 0, 1) = -\tfrac{1}{2} \log 2\pi - \frac{\theta_j^2}{2} = -0.919 - \frac{\theta_j^2}{2}$$

	w_1	w_2	b_1	b_2	v_1	v_2	c	sum
θ_j	−0.483	−2.004	−0.593	0.453	−0.459	−0.960	−0.158	
$\log \mathcal{N}(\theta_j; 0, 1)$	−1.036	−2.927	−1.095	−1.021	−1.024	−1.380	−0.931	**−9.414**

$$\log p(\mathcal{D}) = \log \int_{\mathbb{R}^7} p(\mathcal{D} \mid \theta)\, p(\theta)\, d\theta = \ \textbf{?}$$

> The third term on the right of Bayes' rule, the evidence, is not about θ_A: it needs every θ at once.

Two settings of θ, side by side



	θ_A	θ_B, a prior draw
$\log p(\mathcal{D} \mid \theta)$	12.2	−2,157.5
$\log p(\theta)$	−9.4	−9.9
sum of the two	2.8	−2,167.4

> Their priors are similar; their likelihoods differ by over 2,000 nats. A ratio of two posteriors does not even need p($\mathcal{D}$).


The evidence needs every θ, and has no formula

$$p(\mathcal{D}) = \int_{\mathbb{R}^7} p(\mathcal{D} \mid \theta)\, p(\theta)\, d\theta$$

1 unknown

100 points

2 unknowns




100 × 100 = 10,000 points

parameters	grid points
1	100
2	10,000
7, our network	10^{14}
10^6, a small real network	$10^{2,000,000}$

> Our seven parameters already need 10^{14} grid points; a real network needs 10^2,000,000.

Sampling from the prior: a few lucky draws carry everything




$$p(\mathcal{D}) \approx \frac{1}{S}\sum_{s=1}^{S} p(\mathcal{D} \mid \theta_s)$$

$$\theta_s \sim p(\theta)$$

Ten draws: off by about 1,000 nats. Ten million: about 103 draws carry the estimate, −1.43.

> Almost every prior draw explains the data terribly. Seven parameters just about work; more parameters or more data make it hopeless.

What we want in the end: predictions with error bars




$$p(y^* \mid x^*, \mathcal{D}) = \int p(y^* \mid x^*, \theta) \times p(\theta \mid \mathcal{D})\, d\theta$$

Average every network's prediction, weighted by the posterior. The band widens where there are no data.

> Here it comes from a long MCMC run, used only as a reference. The band widens beyond the data and in the gap.


Three ways to approximate a posterior

Laplace




A Gaussian at the highest peak: N(−2.0, 1.0²). It never sees the second bump.

MCMC



20,000 random-walk steps. Their mean, −0.80, matches the true −0.80.

Variational inference




The closest Gaussian, found by optimisation: N(−0.78, 1.90²).

> Variational inference: choose the closest member of a simple family, by optimisation. That is today.

Variational inference: choose q, make it close to the posterior


$$q_\psi(\theta) = \prod_{j=1}^{7} \mathcal{N}(\theta_j;\, \mu_j,\, \sigma_j^2), \qquad \psi = (\mu_1, \ldots, \mu_7,\, \sigma_1, \ldots, \sigma_7)$$



$$\psi^* = \arg\min_{\psi} \; \text{distance}\big(q_\psi(\theta),\; p(\theta \mid \mathcal{D})\big)$$

> Tune ψ until q is as close as possible to the posterior. Two questions: what is "close", and how do we tune without the posterior?

How far apart are two distributions?




	p	q
mean	0	0
variance	1	1

Same mean, same variance, different distributions. We need one number that compares the whole shapes.

> But here both match, and the shapes still differ. We want a number that compares the distributions everywhere.

Two surveys: how different are they?




Favourite fruit in a class, last year (q) and this year (p).

Bananas fell from 40% to 10%.
Clementines rose from 10% to 40%.

A fourfold fall and a fourfold rise. Should they count the same?


Ratios say “how many times”; logs make up and down cancel

fruit	p	q	p/q	$\log(p/q)$
apple	0.5	0.5	1	0.00
banana	0.1	0.4	¼	−1.39
clementine	0.4	0.1	4	1.39



Take logs: log 4 = +1.39 and log ¼ = −1.39. Equal and opposite.

Weight by how common each fruit is now: that is KL



$$\mathrm{KL}(p \,\|\, q) = \sum_x p(x) \, \log \frac{p(x)}{q(x)}$$

$$= 0.5(0) + 0.1(-1.39) + 0.4(1.39)$$

$$\approx 0.42 \text{ nats}$$

> The rare banana barely counts; the common clementine dominates. Total 0.42 nats: the Kullback–Leibler divergence.

KL is the extra message length of using the wrong code

weather	p	q	code built for p	code built for q
sunny	0.50	0.25	`0`	`10`
cloudy	0.25	0.50	`10`	`0`
rainy	0.25	0.25	`11`	`11`

$$\text{average length under } p: \quad \text{code for } p = 1.50 \text{ bits}, \qquad \text{code for } q = 1.75 \text{ bits}$$

$$1.75 - 1.50 = 0.25 \text{ bits} = \mathrm{KL}(p \| q) \text{ with } \log_2$$

> A code built for the wrong frequencies q needs 1.75. The extra 0.25 bits a day are exactly KL(pllq), in base 2.

You have minimised KL before: in every cross-entropy loss


$$\underbrace{-\sum_x p(x)\log q(x)}_{\text{cross-entropy } H(p,q)} = \underbrace{-\sum_x p(x)\log p(x)}_{\text{entropy } H(p)} + \mathrm{KL}(p\|q)$$

$$\text{classifier:} \quad \min_\theta H(y,\, q_\theta) \iff \min_\theta \mathrm{KL}(y \,\|\, q_\theta)$$

$$\text{maximum likelihood:} \quad -\tfrac{1}{N}\textstyle\sum_i \log p_\theta(x_i) \;\approx\; H(p_{\text{data}}) + \mathrm{KL}(p_{\text{data}} \,\|\, p_\theta)$$

> Maximum likelihood minimises KL(data ‖ model). Variational inference is the third use.

A KL value is how fast evidence piles up



$$\mathrm{KL}\big(\mathrm{Bern}(0.8) \,\|\, \mathrm{Bern}(0.5)\big) \approx 0.19 \text{ nats}$$

$$\frac{p(\mathcal{D} \mid 0.8)}{p(\mathcal{D} \mid 0.5)} \approx e^{0.19\,n}$$

The reverse direction, KL(0.5‖0.8) = 0.22, is a different number.


About e^(30 × 0.19) ≈ 325. KL is the rate at which evidence piles up: 6.9× after 10 tosses, 325× after 30.

For densities, sums become integrals, and KL ≥ 0

$$\mathrm{KL}(q \,\|\, p) = \int q(u)\, \log \frac{q(u)}{p(u)}\, du$$

$$-\mathrm{KL}(q\|p) = \mathbb{E}_q\left[\log \frac{p}{q}\right] \le \mathbb{E}_q\left[\frac{p}{q} - 1\right]$$

$$= \int p\, du - 1 = 0$$



log t never rises above its tangent t − 1. Inside the average that gives KL ≥ 0, with 0 only when q = p.

Closed forms for simple distributions

distributions	$\mathrm{KL}(p \,\|\, q)$	example
Bernoulli(a), Bernoulli(b)	$a \log \frac{a}{b} + (1-a) \log \frac{1-a}{1-b}$	coin, 0.8 vs 0.5: 0.19
categorical	$\sum_k p_k \log \frac{p_k}{q_k}$	fruit surveys: 0.42
Gaussians	$\log \frac{\sigma_2}{\sigma_1} + \frac{\sigma_1^2 + (\mu_1 - \mu_2)^2}{2\sigma_2^2} - \frac{1}{2}$	derived next
diagonal Gaussians	sum of one-dimensional terms	our q against the prior

A mixture, or a posterior, has no closed form. Then we estimate the average by sampling.

> Anything else, such as a mixture or a posterior, has no formula; then we sample.

Deriving the Gaussian formula


$$q = \mathcal{N}(\mu, \sigma^2), \quad p = \mathcal{N}(m, s^2): \qquad \log \frac{q(u)}{p(u)} = \log \frac{s}{\sigma} + \frac{(u-m)^2}{2s^2} - \frac{(u-\mu)^2}{2\sigma^2}$$

$$\mathbb{E}_q[(u-m)^2] = \sigma^2 + (\mu - m)^2, \qquad \mathbb{E}_q[(u-\mu)^2] = \sigma^2$$

$$\mathrm{KL}(q\|p) = \log \frac{s}{\sigma} + \frac{\sigma^2 + (\mu - m)^2}{2s^2} - \frac{1}{2}$$

The result: a width term, a squared shift of the centres, and −½.

Use the formula: which Gaussian is closer to p?





$$p = \mathcal{N}(1, 4^2)$$

	substitute	KL
$q_1 = \mathcal{N}(0, 2^2)$	$\log 2 + \frac{4+1}{32} - \frac{1}{2}$	0.35
$q_2 = \mathcal{N}(1, 3^2)$	$\log \frac{4}{3} + \frac{9+0}{32} - \frac{1}{2}$	0.07

KL(q_1‖p) = 0.35 nats and KL(q_2‖p) = 0.07. The formula settles the vote.

KL is an area you can watch





Purple area adds and red area subtracts; the net area is the KL. Drag q, or flip the direction.

No formula needed: estimate KL by sampling


$$\mathrm{KL}(q\|p) = \mathbb{E}_{u\sim q}\big[\log q(u) - \log p(u)\big] \approx \frac{1}{S}\sum_{i=1}^{S}\big[\log q(u_i) - \log p(u_i)\big]$$


```
u = q.sample((S,))
log_q = q.log_prob(u)
log_p = p.log_prob(u)
est = (log_q - log_p).mean()
exact = kl_divergence(q, p)
```

	$KL(q_2\|p)$
exact formula	0.069
estimate, S = 10,000	0.067

Checked on a pair that has a formula: 0.067 from 10,000 draws, against the exact 0.069.

Watch the estimate settle







Add draws from q and watch the running average of log q − log p settle on the exact 0.349.

Six candidates: which q is closest to p?


$q = N(0, 2^2)$ KL 0.35



$q = N(1, 3^2)$ **KL 0.07**



$q = N(4, 1^2)$ KL 1.20



$q = N(-3, 4^2)$ KL 0.50



$q = N(1, 8^2)$ KL 0.81




$q = N(2, 5^2)$ KL 0.09



Smallest KL: $N(1, 3^2)$ at 0.07, then $N(2, 5^2)$ at 0.09. Could another q do better?

Search a grid of means and widths




Evaluate $\mathrm{KL}(q\|p)$ at every point of a 51 × 40 grid of means and widths: 2,040 evaluations of the formula.

Best grid point: q = p. But our network's q has 14 numbers; a grid with 40 values each needs $40^{14} \approx 10^{22}$ evaluations.

> The best grid point is N(1, 4²), which is p. For the network's fourteen numbers, a grid is hopeless.

Instead of a grid, walk downhill



$$\psi^* = \arg\min_{\psi} \mathrm{KL}(q_\psi \,\|\, p)$$

$$\psi \leftarrow \psi - \eta \, \nabla_\psi \mathrm{KL}(q_\psi \,\|\, p)$$

Gradient descent from N(4, 0.5²): the teal path. About a hundred steps, and the same recipe works for millions of parameters.

From N(4, 0.5²) the optimiser walks downhill to p. It needs only gradients, which autograd provides.

The same search in code


```
mu  = Parameter(tensor(4.0))
rho = Parameter(tensor(log(0.5)))
opt = Adam([mu, rho], lr=0.05)
for step in range(200):
    q = Normal(mu, rho.exp())
    loss = kl_divergence(q, p)
    opt.zero_grad()
    loss.backward()
    opt.step()
```


$$\sigma = \exp(\rho) > 0 \ \text{ for every } \rho$$

Rebuild q, score it, differentiate, update. The same loop fits our network in Part 5.

Rebuild q, score it, differentiate, update. The same five lines fit our network in Part 5.


Press play: q slides onto p






> The width gets there first and overshoots a little; the centre slides in by step 100. Then KL ≈ 0.

In 2-D, a q that cannot tilt shrinks instead





$$p = \mathcal{N}\left(0, \begin{bmatrix} 1 & 0.9 \\ 0.9 & 1 \end{bmatrix}\right)$$


$$q = \mathcal{N}\left(0, \mathrm{diag}(s^2, s^2)\right)$$

$$s = 0.44, \quad \mathrm{KL} = 0.83$$

$$q = \mathcal{N}(0, LL^\top) : \ \mathrm{KL} = 0$$

> Let q tilt with a full covariance, Σ = LLᵀ, and the KL drops to 0.

PyTorch has no formula for this KL. The definition still works.





```
kl_divergence(q, mixture)
```

```
NotImplementedError:
No KL(p || q) is implemented
for p type Normal and
q type MixtureSameFamily
```

No formula. But KL is an average under q, and averages can be estimated by sampling.


Now optimise it: the code runs, and q does not move

```
u = q.sample((5000,))
log_q = q.log_prob(u)
log_p = p.log_prob(u)
loss = (log_q - log_p).mean()
opt.zero_grad()
loss.backward(); opt.step()
```



No error and no warning, and q stays at N(6, 0.1²). After 400 steps the KL is still about 53.


The gradient that would move q never reaches it



	$\partial \mathrm{KL}/\partial \mu$	$\partial \mathrm{KL}/\partial \sigma$
through sample()	0.0013	−0.0003
true gradient	−0.063	−0.375

> What survives averages to zero: 0.0013 and −0.0003, where the true gradient is −0.063 and −0.375.

Move the randomness outside: u = μ + σε



$$\epsilon \sim \mathcal{N}(0,1), \qquad u = \mu + \sigma\epsilon$$

$$\epsilon = -0.75, \quad \mu = 0, \quad \sigma = 2$$


$$u = -1.5$$

$$\frac{\partial u}{\partial \mu} = 1, \qquad \frac{\partial u}{\partial \sigma} = \epsilon = -0.75$$


u = −1.5, ∂u/∂μ = 1 and ∂u/∂σ = ε = −0.75. The gradient now has a path. In PyTorch this is rsample().

What a fixed ε does when μ or σ changes

move μ from 0 to 1 u: −1.50 → −0.50



widen σ from 2 to 3 u: −1.50 → −2.25




$$\frac{\partial u}{\partial \mu} = 1, \qquad \frac{\partial u}{\partial \sigma} = \epsilon = -0.75$$

> Widen σ by 1: the draw moves by ε = −0.75, further out on its side. These are the derivatives the gradient needs.

Change one word: rsample() fits the mixture


```
# one word changed
u = q.rsample((5000,))
log_q = q.log_prob(u)
log_p = p.log_prob(u)
loss = (log_q - log_p).mean()
opt.zero_grad()
loss.backward(); opt.step()
```




q settles at N(0.43, 0.72²), KL = 0.056: the best single Gaussian for this mixture. No formula was needed.

The price: reverse KL picks one bump


KL(p||q) 0.72



KL(q||p), start left 0.36



KL(q||p), start right 1.20



KL(p||q) spreads over both bumps. KL(q||p) locks onto whichever bump is nearer its start, and reports it confidently.

Back to Bayes: minimise KL to the posterior directly?

$$\mathrm{KL}\big(q_\psi(\theta) \,\|\, p(\theta \mid \mathcal{D})\big) = \mathbb{E}_{\theta\sim q}\big[\log q_\psi(\theta) - \log p(\theta \mid \mathcal{D})\big]$$

the loss needs	available?
draws θ ~ q	yes: rsample, as in Part 3
$\log q_\psi(\theta)$	yes
$\log p(\theta \mid \mathcal{D}) = \log p(\mathcal{D} \mid \theta) + \log p(\theta) - \log p(\mathcal{D})$	no: needs p($\mathcal{D}$)

> Everything is available except log p(θ | $\mathcal{D}$), which needs the evidence p($\mathcal{D}$): the integral we cannot do.

Substitute Bayes' rule and see what survives

$$\mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big) = \mathbb{E}_q\big[\log q(\theta) - \log p(\theta \mid \mathcal{D})\big]$$

$$= \mathbb{E}_q\big[\log q(\theta) - \log p(\mathcal{D} \mid \theta) - \log p(\theta)\big] + \log p(\mathcal{D})$$

$$\log p(\mathcal{D}) = \underbrace{\mathbb{E}_q\big[\log p(\mathcal{D} \mid \theta) + \log p(\theta) - \log q(\theta)\big]}_{\mathrm{ELBO}(q)} + \mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big)$$

Rearrange: log p($\mathcal{D}$) = ELBO + KL. The ELBO needs only the likelihood, the prior and q.

The ELBO for our network: fit the data, stay near the prior


$$\mathrm{ELBO}(q) = \mathbb{E}_q\big[\log p(\mathcal{D} \mid \theta)\big] + \mathbb{E}_q\big[\log p(\theta) - \log q(\theta)\big]$$

$$= \underbrace{\mathbb{E}_q\big[\log p(\mathcal{D} \mid \theta)\big]}_{\text{fit the data}} - \underbrace{\mathrm{KL}\big(q(\theta) \,\|\, p(\theta)\big)}_{\text{stay near the prior}}$$


The second term is the diagonal-Gaussian KL from Part 2: a formula, seven terms. The first is a Monte Carlo average over networks sampled with rsample.

> The KL to the prior has a formula; the data term is estimated by sampling θ from q, with the reparameterisation trick.

Check the identity where p($\mathcal{D}$) is known: a coin



One fixed total



−22.020 + 0.623 = −21.397

> A coin with a Beta(5, 3) prior and 17 heads in 30 tosses has an exact posterior and an exact p($\mathcal{D}$). Drag q: the ELBO and the KL trade off, and their sum never changes.


The ELBO for our network, in code



```
mu  = Parameter(0.5 * randn(7))          # 7 means
rho = Parameter(full((7,), log(0.1)))    # 7 log-widths
q = Normal(mu, rho.exp())                # one bell per parameter
theta = q.rsample((16,))                 # 16 sampled networks
log_lik = log_likelihood(theta).mean()   # forward pass
kl = kl_divergence(q, prior).sum()       # closed form
loss = -(log_lik - kl)                   # minus the ELBO
```

Adam, learning rate 0.02, 4,000 steps; best of five random starts. No p($\mathcal{D}$) anywhere.

Every line is computable: rsample, a forward pass and a formula. Never p($\mathcal{D}$).


Press play: q learns the network





> The ELBO climbs from −1,235 to about −7.2 and the band settles onto the data. It stays below log p(𝒟) ≈ −1.43.


What q learned: one hidden unit switched off



$v_1 \approx 0.01$ with width 0.035: the first hidden unit is switched off, and its w_1 and b_1 fall back to the prior.

> VI switched the first hidden unit off. Mean-field VI often prunes units like this: every parameter kept away from the prior costs KL.

VI against the reference: the right curve, too-narrow bands



predictive sd	VI	reference
at x = −2.5	0.11	0.22
at x = 2.5	0.11	0.17

ELBO −7.18 against log $p(\mathcal{D}) \approx -1.43$: q is about 5.8 nats from the posterior.

> Its bands are too narrow beyond the data, as the diagonal q in 2-D predicted. The ELBO gap says how far q is from the posterior.

Six obstacles, six tools

obstacle	tool	in our network
$p(\mathcal{D})$ is an impossible integral	approximate with q	one Gaussian per parameter
“close” needs a number	$KL(q \Vert p)$	KL to the prior
a grid cannot search 14 numbers	gradient descent	Adam on μ and ρ
no formula for the average	Monte Carlo	16 sampled networks
no gradient through a draw	$\theta = \mu + \sigma\varepsilon$, rsample()	reparameterisation
KL to the posterior needs $p(\mathcal{D})$	maximise the ELBO	the training loss

> Every obstacle today earned a tool, and our network used them all. In Lecture 3 a VAE does the same with one q per image.

Generative models create this inference problem again

A hidden description can generate an observation.

Next: learn representations for images, then ask whether they can generate new ones.

> Lecture 2 builds autoencoders; Lecture 3 returns to VI with an encoder that outputs q for each image.

A sensor has an unknown constant bias


$$u \sim \mathcal{N}(0, 1), \qquad y_i \mid u \sim \mathcal{N}(u, 0.5^2)$$

$$p(\mathcal{D}, u) = p(u) \prod_i p(y_i \mid u)$$

$$\max_{\psi} \mathbb{E}_{q_\psi(u)}[\log p(\mathcal{D}, u) - \log q_\psi(u)]$$


Maximise the ELBO over q's parameters. The same tools apply.

More samples, tighter estimates



Forty repeats at each sample size. The spread shrinks like $1/\sqrt{S}$; the exact value never moves.

A check with an answer key: the Beta coin



After 600 steps: Beta(22.4, 16.2), with KL ≈ 0.0002 to the true posterior.

One literal update of the mean

$$\sigma = 2 \text{ fixed}: \qquad L(\mu) = \text{const} + \frac{(\mu - 1)^2}{32}$$

$$\frac{\partial L}{\partial \mu} = \frac{\mu - 1}{16} = -0.0625 \ \text{ at } \mu = 0$$

$$\mu^{+} = 0 - 4(-0.0625) = 0.25: \quad \text{KL } 0.349 \to 0.336$$

Take one step with η = 4. What is the best KL with σ stuck at 2?

KL ≥ 0 by Jensen's inequality

$$\mathrm{KL}(q\|p) = -\mathbb{E}_q\left[\log \frac{p(u)}{q(u)}\right]$$

$$\geq -\log \mathbb{E}_q\left[\frac{p(u)}{q(u)}\right]$$

$$= -\log 1 = 0$$

What makes the final expectation one?

Five draws can give a negative KL estimate

u_i	$\log q(u_i) - \log p(u_i)$
5.623	−0.232
0.120	0.269
−5.536	−0.751
2.705	0.217
−2.254	0.030

$$\widehat{\mathrm{KL}}_5 \approx -0.093, \qquad \mathrm{KL}(q\|p) \approx 0.069$$

> Does this contradict KL ≥ 0?

The two PML candidates, scored by Monte Carlo

candidate	MC estimate ± 1 s.e.	numerical integral
$N(3, 1^2)$	9.54 ± 0.03	9.49
$N(3, 4.5^2)$	45.02 ± 0.21	45.15

The narrow candidate scores far better. Both sit away from the mixture's mass, so both are poor.

> What role does the numerical integral play?

A fixed-noise finite difference checks the path

```
q = Normal(mu, sigma)
u = q.rsample((S,))
loss = (q.log_prob(u) - p.log_prob(u)).mean()
```

derivative	autograd	finite difference
μ	−0.062466	−0.062466
σ	−0.372223	−0.372223

What must stay fixed on both sides of the finite difference?


Pathwise gradients are one route, not the only one

$$\nabla_\psi \mathbb{E}_{q_\psi}[f_\psi(u)] = \mathbb{E}_{q_\psi}[f_\psi(u)\nabla_\psi \log q_\psi(u) + \nabla_\psi f_\psi(u)]$$

A score-function estimator accounts explicitly for the changing sampling distribution.

> Does this identity require u = μ + σε?

A noisy ELBO estimate can cross the evidence



162 of 600 sampled estimates lie above the known evidence in this run.

> Which object remains a lower bound?


A higher ELBO does not prove a zero gap

$$\mathrm{ELBO}(q_{\mathrm{new}}) > \mathrm{ELBO}(q_{\mathrm{old}})$$

The family may not contain the posterior, and optimisation may stop short.

> Does a stable loss establish q = p(u | $\mathcal{D}$)?

Check the fixed total with three Beta candidates



What happens at the exact posterior?

A mean must say which dimension it averages

object	shape	correct reduction
S scalar Beta draws	[S]	mean over S
S d-dimensional joint log densities	[S]	mean over S
S factorised coordinate log densities	[S, d]	sum over d, then mean over S

MultivariateNormal.log_prob already sums the event dimension. Do not sum the sample axis as well.

> What goes wrong if you sum over S before averaging?

An ordered sequence and a count have different evidence

$$p(H = 17 \mid r) = \binom{30}{17} r^{17}(1-r)^{13}$$

observed object	log evidence
one ordered sequence	−21.397
only the head count	−2.796

> Does this constant change the posterior or its KL gap?

Check the sampled objective with analytic Beta moments

$$\mathbb{E}_q[\log r] = \mathrm{digamma}(a) - \mathrm{digamma}(a+b)$$

$$\mathbb{E}_q[\log(1-r)] = \mathrm{digamma}(b) - \mathrm{digamma}(a+b)$$

Insert these into expected log likelihood + expected log prior − expected log q.

> How was the identity checked non-circularly?