▶ Notebook

Lecture 1 at a glance: five parts, twenty-three sections

Part 1 · Bayes' rule for a neural network

1.1 Bayes' rule ▶
1.2 A tiny network and its prior ▶
1.3 Likelihood and training ▶
1.4 The evidence is intractable ▶
1.5 Approximating the posterior ▶

Part 2 · How far apart are two distributions?

2.1 Surprise ▶
2.2 Entropy: average surprise ▶
2.3 Codes: entropy in bits ▶
2.4 Cross-entropy and KL ▶
2.5 Classification loss ▶
2.6 Maximum likelihood ▶
2.7 Closed forms and PyTorch ▶
2.8 KL by sampling ▶

Part 3 · Find the closest q

3.1 Search a grid ▶
3.2 Gradient descent ▶
3.3 Covariance in 2-D ▶
3.4 Mixtures: no formula ▶
3.5 The reparameterisation trick ▶
3.6 Which direction of KL? ▶

Part 4 · Back to Bayes: the ELBO

4.1 Deriving the ELBO ▶
4.2 Checking it on a coin ▶

Part 5 · Implement it

5.1 VI for the tiny network ▶
5.2 Wrap-up ▶

> Five parts, twenty-three numbered sections. The notebook uses the same numbers: the ▶ Notebook link at the top right of every slide opens its section in Colab.

▶ Notebook 1.1

Bayes' rule: update what you believe, using data

$$p(A \mid B) = \frac{p(B \mid A)\, p(A)}{p(B)}$$

> Bayes' rule: what to believe after seeing B, from what we believed before and how well each belief explains B.

▶ Notebook 1.1

Bayes' rule: update what you believe, using data

$$p(A \mid B) = \frac{p(B \mid A)\, p(A)}{p(B)}$$

$$\underbrace{p(A \mid B)}_{\text{posterior}} = \frac{\overbrace{p(B \mid A)}^{\text{likelihood}}\ \overbrace{p(A)}^{\text{prior}}}{\underbrace{p(B)}_{\text{evidence}}}$$

> Four names for the whole lecture: posterior, likelihood, prior and evidence.

▶ Notebook 1.1

In machine learning, the unknown is the parameters

$$p(\theta \mid \mathcal{D}) = \frac{p(\mathcal{D} \mid \theta)\, p(\theta)}{p(\mathcal{D})}$$

A becomes θ, every weight and bias of the model; B becomes $\mathcal{D}$, the training data.

▶ Notebook 1.1

In machine learning, the unknown is the parameters

$$p(\theta \mid \mathcal{D}) = \frac{p(\mathcal{D} \mid \theta)\, p(\theta)}{p(\mathcal{D})}$$

Usual training

keeps one θ: the best fit to the data $\mathcal{D}$.


Bayesian learning

keeps every θ, weighted by $p(\theta \mid \mathcal{D})$.

> Usual training keeps one θ. The Bayesian answer keeps a whole distribution over θ.

▶ Notebook 1.2

A tiny network: one input, two hidden units, one output




$$f_\theta(x) = v_1 \tanh(w_1 x + b_1) + v_2 \tanh(w_2 x + b_2) + c$$

Our running example: one input x, two tanh hidden units and one output f.

▶ Notebook 1.2

A tiny network: one input, two hidden units, one output



$$f_\theta(x) = v_1 \tanh(w_1 x + b_1) + v_2 \tanh(w_2 x + b_2) + c$$

$$\theta = (w_1, w_2, b_1, b_2, v_1, v_2, c) \in \mathbb{R}^7$$

Seven unknown numbers: four weights and three biases.

> Every weight and bias is unknown, so θ is seven numbers.

▶ Notebook 1.2

A prior on every parameter

$$\theta_j \sim \mathcal{N}(0, 1), \quad j = 1, \ldots, 7$$

$$p(\theta) = \prod_{j=1}^{7} \mathcal{N}(\theta_j;\, 0,\, 1)$$


Before seeing data, every parameter gets its own prior: $\theta_j \sim N(0, 1)$, independently.

▶ Notebook 1.2

A prior on every parameter

$$\theta_j \sim \mathcal{N}(0, 1), \quad j = 1, \ldots, 7$$

$$p(\theta) = \prod_{j=1}^{7} \mathcal{N}(\theta_j;\, 0,\, 1)$$




One draw from the prior is seven numbers, one from each bell: a single point θ in $\mathbb{R}^7$.

▶ Notebook 1.2

One draw of θ is one whole function

	w_1	w_2	b_1	b_2	v_1	v_2	c
$\theta^{(1)}$	0.79	1.39	−0.15	−0.02	0.28	−1.10	−1.03

Each row, put through the network, is one curve.




> Draw $\theta^{(1)}$ from the prior, put its seven numbers into the network, and evaluate f at every x: one curve.

▶ Notebook 1.2

One draw of θ is one whole function

	w_1	w_2	b_1	b_2	v_1	v_2	c
$\theta^{(1)}$	0.79	1.39	−0.15	−0.02	0.28	−1.10	−1.03
$\theta^{(2)}$	−1.07	−1.40	0.04	0.78	0.45	−1.93	0.77

Each row, put through the network, is one curve.




A second draw, $\theta^{(2)}$, gives a different curve.

▶ Notebook 1.2

One draw of θ is one whole function

	w_1	w_2	b_1	b_2	v_1	v_2	c
$\theta^{(1)}$	0.79	1.39	−0.15	−0.02	0.28	−1.10	−1.03
$\theta^{(2)}$	−1.07	−1.40	0.04	0.78	0.45	−1.93	0.77
$\theta^{(3)}$	−1.06	0.20	−0.63	0.22	0.48	−0.56	1.66


Each row, put through the network, is one curve.




> Every point of $\mathbb{R}^7$ is a whole function. Drawing θ from the prior draws functions.

▶ Notebook 1.2

Twenty draws, and the prior predictive






Twenty draws of θ, twenty functions. Hover over a line, or step through, to see its point in $\mathbb{R}^7$.


▶ Notebook 1.2

Twenty draws, and the prior predictive

20 draws, and the prior predictive: mean ± 2 sd



draw 1 of 20: $\theta^{(1)} \in \mathbb{R}^7$




Hover over a line, or step through the draws.

> Average over every θ and add the 0.1 noise: the prior predictive, $p(y \mid x) = \int p(y \mid x, \theta)\, p(\theta)\, d\theta$. What the model expects before any data.

▶ Notebook 1.3

The data, and how well one θ explains them




$$y_i \mid x_i, \theta \sim \mathcal{N}\left(f_\theta(x_i),\, 0.1^2\right)$$

> Seventeen training points with a gap in the middle. Each y is the network's output plus Gaussian noise of size 0.1.

▶ Notebook 1.3

The data, and how well one θ explains them



$$y_i \mid x_i, \theta \sim \mathcal{N}\big(f_\theta(x_i),\, 0.1^2\big)$$

$$p(\mathcal{D} \mid \theta) = \prod_{i=1}^{17} \mathcal{N}\big(y_i;\, f_\theta(x_i),\, 0.1^2\big)$$


> The likelihood of θ multiplies all seventeen densities: how well this one θ explains the whole dataset.

▶ Notebook 1.3

Finding a good θ: train the network

$$\theta_A = \arg\max_{\theta} \left[\log p(\mathcal{D} \mid \theta) + \log p(\theta) \right]$$

the network during training



loss 559 · grey: the other nine starts




To pick one good θ, train the network the usual way: maximise log p($\mathcal{D}$ | θ) + log p(θ). That is squared error plus weight decay.

▶ Notebook 1.3


Finding a good θ: train the network

$$\theta_A = \arg\max_{\theta} \left[\log p(\mathcal{D} \mid \theta) + \log p(\theta) \right]$$

the network during training




loss 14.3 · grey: the other nine starts



From a random start, gradient descent bends the curve onto the data. Ten starts finish at slightly different losses; the best finish is θ_A.

▶ Notebook 1.3


Meet θ_A: the network with its trained numbers




θ_A is one point in $\mathbb{R}^7$: these seven numbers, written on the network.

▶ Notebook 1.3

Meet θ_A: the network with its trained numbers





With these numbers the network's curve follows the data. Next, score θ_A with the likelihood and the prior.

▶ Notebook 1.3

The likelihood of θ_A: formula, then value

$$p(\mathcal{D} \mid \theta_A) = \prod_{i=1}^{17} \mathcal{N}\left(y_i;\ f_{\theta_A}(x_i),\ 0.1^2\right)$$

Fix $\theta = \theta_A$. The likelihood multiplies one Gaussian density per data point: how well the curve explains each y.

▶ Notebook 1.3

The likelihood of θ_A: formula, then value

$$p(\mathcal{D} \mid \theta_A) = \prod_{i=1}^{17} \mathcal{N}\left(y_i;\, f_{\theta_A}(x_i),\, 0.1^2\right)$$

$$\log p(\mathcal{D} \mid \theta_A) = -\frac{1}{2 \cdot 0.1^2} \sum_{i=1}^{17} \left(y_i - f_{\theta_A}(x_i)\right)^2 - \frac{17}{2} \log\left(2\pi \cdot 0.1^2\right)$$

Take logs: the product becomes a sum of squared errors, scaled by the noise, plus a constant.

▶ Notebook 1.3

The likelihood of θ_A: formula, then value

$$p(\mathcal{D} \mid \theta_A) = \prod_{i=1}^{17} \mathcal{N}\left(y_i;\ f_{\theta_A}(x_i),\ 0.1^2\right)$$

$$\log p(\mathcal{D} \mid \theta_A) = -\frac{1}{2 \cdot 0.1^2} \sum_{i=1}^{17} \left(y_i - f_{\theta_A}(x_i)\right)^2 - \frac{17}{2} \log\left(2\pi \cdot 0.1^2\right)$$

$$\log p(\mathcal{D} \mid \theta_A) = 12.17$$

Put in θ_A's curve and the seventeen points: $\log p(\mathcal{D} \mid \theta_A) = 12.17$.

▶ Notebook 1.3

The prior of θ_A, then the evidence

$$p(\theta_A) = \prod_{j=1}^{7} \mathcal{N}(\theta_j;\, 0,\, 1)$$

The prior multiplies one N(0, 1) density per parameter.

▶ Notebook 1.3

The prior of θ_A, then the evidence

$$p(\theta_A) = \prod_{j=1}^{7} \mathcal{N}(\theta_j;\, 0,\, 1)$$

$$\log p(\theta_A) = -\frac{1}{2}\sum_{j=1}^{7} \theta_j^2 - \frac{7}{2}\log 2\pi = -9.41$$

> Take logs: minus half the sum of squared parameters, plus a constant. For θ_A it is −9.41; the large w_2 costs the most.

▶ Notebook 1.3

The prior of θ_A, then the evidence

$$p(\theta_A) = \prod_{j=1}^{7} \mathcal{N}(\theta_j;\, 0,\, 1)$$


$$\log p(\theta_A) = -\frac{1}{2}\sum_{j=1}^{7} \theta_j^2 - \frac{7}{2}\log 2\pi = -9.41$$

$$p(\mathcal{D}) = \int p(\mathcal{D} \mid \theta)\, p(\theta)\, d\theta = \ \mathbf{?}$$

> The third term, the evidence, is not about θ_A: it needs every θ at once.

▶ Notebook 1.3


Two settings of θ, side by side



θ_A follows the data. θ_B, one random draw from the prior, does not.

▶ Notebook 1.3

Two settings of θ, side by side




	θ_A	θ_B, a prior draw
$\log p(\mathcal{D} \mid \theta)$	12.2	−2,157.5
$\log p(\theta)$	−9.4	−9.9
sum of the two	2.8	−2,167.4

> Their priors are similar; their likelihoods differ by over 2,000 nats. A ratio of two posteriors does not even need p($\mathcal{D}$).

▶ Notebook 1.4

The evidence is an area: add up thin rectangles



Suppose only w_2 were unknown, with the other six numbers fixed at θ_A:


$$p(\mathcal{D}) = \int p(\mathcal{D} \mid w_2)\, p(w_2)\, dw_2$$

With one unknown, the evidence is the area under one curve: likelihood times prior.

▶ Notebook 1.4

The evidence is an area: add up thin rectangles

p($\mathcal{D}$ | w_2) p(w_2), the other six fixed at θ_A



Suppose only w_2 were unknown, with the other six numbers fixed at θ_A:

$$p(\mathcal{D}) = \int p(\mathcal{D} \mid w_2)\, p(w_2)\, dw_2$$

$$\approx \sum_{k=1}^{20} p(\mathcal{D} \mid w_{2,k})\, p(w_{2,k})\, \Delta w$$

> Chop the area into thin rectangles and add them up. Twenty rectangles already match a much finer grid to three decimals.

▶ Notebook 1.4


Intractable: why the rectangles do not scale

Intractable: no formula, and a cost that grows exponentially with the number of unknowns.

1 unknown

100 points

2 unknowns



100 × 100 = 10,000 points

> With 100 rectangles per unknown, two unknowns already need 10,000 boxes.


▶ Notebook 1.4

Intractable: why the rectangles do not scale

1 unknown

100 points

2 unknowns



100 × 100 = 10,000 points


Intractable: no formula, and a cost that grows exponentially with the number of unknowns.

unknowns	grid points
1	100
2	10,000
7, our network	10^{14}: a day
10^6	$10^{2,000,000}$: never

> Our seven need 10^{14}; a real network, $10^{2,000,000}$. That is what intractable means.

▶ Notebook 1.5

What we want in the end: predictions with error bars




$$p(y^* \mid x^*, \mathcal{D}) = \int p(y^* \mid x^*, \theta) \times p(\theta \mid \mathcal{D})\, d\theta$$

> The posterior is a means to an end. The posterior predictive averages every network's prediction at a new x*.

▶ Notebook 1.5

What we want in the end: predictions with error bars



$$p(y^* \mid x^*, \mathcal{D}) = \int p(y^* \mid x^*, \theta) \times p(\theta \mid \mathcal{D})\, d\theta$$

Average every network's prediction, weighted by the posterior. The band widens where there are no data.

> Here it comes from a long MCMC run, used only as a reference. The band widens beyond the data and in the gap.

▶ Notebook 1.5

The core problem of probabilistic machine learning

$$p(\theta \mid \mathcal{D}) = \frac{\overbrace{p(\mathcal{D} \mid \theta)}^{\text{easy}} \; \overbrace{p(\theta)}^{\text{easy}}}{\underbrace{p(\mathcal{D})}_{\text{intractable}}}$$

> The likelihood and the prior are easy to evaluate for any θ. The evidence, which turns them into the posterior, is not.

▶ Notebook 1.5

The core problem of probabilistic machine learning

$$p(\theta \mid \mathcal{D}) = \frac{\overbrace{p(\mathcal{D} \mid \theta)}^{\text{easy}} \overbrace{p(\theta)}^{\text{easy}}}{\underbrace{p(\mathcal{D})}_{\text{intractable}}}$$

How do we compute, or approximate, the posterior $p(\theta \mid \mathcal{D})$?


Bayesian neural networks, VAEs, mixture models and topic models all meet this same question.

> So the central question of probabilistic machine learning: how do we compute, or approximate, the posterior?

▶ Notebook 1.5

Three ways to approximate the posterior; we take the third

Laplace




A Gaussian at the highest peak: $N(-2.0, 1.0^2)$. It never sees the second bump.

> Laplace: fit a Gaussian to the curvature at the highest peak.

▶ Notebook 1.5


Three ways to approximate the posterior; we take the third

Laplace



A Gaussian at the highest peak: $N(-2.0, 1.0^2)$. It never sees the second bump.

MCMC




20,000 random-walk steps. Their mean, −0.80, matches the true −0.80.

> MCMC: a random walk whose visits follow the posterior. Right in the long run, but slow.

▶ Notebook 1.5


Three ways to approximate the posterior; we take the third

Laplace




A Gaussian at the highest peak: N(−2.0, 1.0²). It never sees the second bump.

MCMC



20,000 random-walk steps. Their mean, −0.80, matches the true −0.80.

Variational inference



The closest Gaussian, found by optimisation: N(−0.78, 1.90²).

> Variational inference: the closest member of a simple family, found by optimisation. The rest of the lecture develops only this one.

▶ Notebook 1.5

Variational inference: choose q, make it close to the posterior


$$q_\psi(\theta) = \prod_{j=1}^{7} \mathcal{N}(\theta_j;\, \mu_j,\, \sigma_j^2), \qquad \psi = (\mu_1, \ldots, \mu_7,\, \sigma_1, \ldots, \sigma_7)$$

Pick a simple family: one Gaussian per parameter, each with its own mean and width. ψ is fourteen numbers.

▶ Notebook 1.5

Variational inference: choose q, make it close to the posterior

$$q_\psi(\theta) = \prod_{j=1}^{7} \mathcal{N}(\theta_j;\, \mu_j,\, \sigma_j^2), \qquad \psi = (\mu_1, \ldots, \mu_7,\, \sigma_1, \ldots, \sigma_7)$$




One bell per parameter. Together they define q(θ), and drawing θ from q is easy.

▶ Notebook 1.5

Variational inference: choose q, make it close to the posterior

$$q_\psi(\theta) = \prod_{j=1}^{7} \mathcal{N}(\theta_j;\, \mu_j,\, \sigma_j^2), \qquad \psi = (\mu_1, \ldots, \mu_7,\, \sigma_1, \ldots, \sigma_7)$$




$$\psi^* = \arg\min_{\psi} \text{ distance}\big(q_\psi(\theta),\, p(\theta \mid \mathcal{D})\big)$$

> Tune ψ until q is as close as possible to the posterior. Two questions: what is "close", and how do we tune without the posterior?

▶ Notebook 2.1

For now, pretend we know the posterior

— p(θ | $\mathcal{D}$), pretend




$$p(\theta \mid \mathcal{D}) = \mathcal{N}(1, 4^2)$$

A stand-in for the real posterior: known, one-dimensional, one bump.

To learn the tools, pretend for a while that we know the posterior: one Gaussian, in one dimension.

▶ Notebook 2.1

For now, pretend we know the posterior



$$p(\theta \mid \mathcal{D}) = \mathcal{N}(1, 4^2)$$

A stand-in for the real posterior: known, one-dimensional, one bump.


$$q(\theta) = \mathcal{N}(\mu, \sigma^2)$$

Our approximation: we choose μ and σ. Shown: μ = 0, σ = 2.

> Our approximation q is another Gaussian; we choose its mean and its width.

▶ Notebook 2.1

For now, pretend we know the posterior



$$p(\theta \mid \mathcal{D}) = \mathcal{N}(1, 4^2)$$

A stand-in for the real posterior: known, one-dimensional, one bump.

$$q(\theta) = \mathcal{N}(\mu, \sigma^2)$$


Our approximation: we choose μ and σ. Shown: μ = 0, σ = 2.

How far apart are p and q? We need one number, so that we can make it small.

> How far apart are p and q? We need one number, and we will build it from scratch, starting from surprise.

▶ Notebook 2.1

How surprised would you be? A forecast of 90% sun




The forecast for tomorrow: sun with probability 0.9, rain with probability 0.1.

> Start with one outcome. The forecast says 90% sun, 10% rain.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.1

How surprised would you be? A forecast of 90% sun



The forecast for tomorrow: sun with probability 0.9, rain with probability 0.1.

If it rains you are surprised; if the sun shines, hardly at all. **Rare outcomes carry more surprise.**

Surprise is information: the outcome you did not expect tells you the most.

> Rain would surprise you; sun would not. We want a number for surprise: big for rare outcomes, small for common ones.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.1

What should surprise be? Three requirements

$$I(1) = 0$$

a certain outcome is no surprise

$$p \downarrow \Rightarrow \; I(p) \uparrow$$

rarer means more surprising

$$I(p_1\, p_2) = I(p_1) + I(p_2)$$

two independent days: probabilities multiply, surprises add

candidate	I(1) = 0	rarer, more	days add: I(½ · 0.1) vs I(½) + I(0.1)

> Surprise I(p) should be zero for a certain outcome, larger for rarer ones, and for two independent days, whose probabilities multiply, the surprises should add.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.1

What should surprise be? Three requirements

$$I(1) = 0$$

a certain outcome is no surprise

$$p \downarrow \Rightarrow \; I(p) \uparrow$$

rarer means more surprising

$$I(p_1\, p_2) = I(p_1) + I(p_2)$$

two independent days: probabilities multiply, surprises add

candidate	I(1) = 0	rarer, more	days add: I(½ · 0.1) vs I(½) + I(0.1)
$1 - p$	✓	✓	✗ 0.95 vs 1.4
$1/p$	✗ 1	✓	✗ 20 vs 12

> Test 1 − p and 1/p on two days with probabilities ½ and 0.1. Neither adds up, and 1/p even gives a certain outcome a surprise of 1.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.1

What should surprise be? Three requirements

$$I(1) = 0$$

a certain outcome is no surprise

$$p \downarrow \Rightarrow \; I(p) \uparrow$$

rarer means more surprising

$$I(p_1\, p_2) = I(p_1) + I(p_2)$$

two independent days: probabilities multiply, surprises add

candidate	I(1) = 0	rarer, more	days add: I(½ · 0.1) vs I(½) + I(0.1)
$1 - p$	✓	✓	✗ 0.95 vs 1.4
$1/p$	✗ 1	✓	✗ 20 vs 12
$\log_2 \frac{1}{p}$	✓	✓	✓ 4.32 = 1 + 3.32

> The logarithm passes all three: log $1/(p_1p_2)$ = log $1/p_1$ + log $1/p_2$. Only the log turns products into sums. So surprise is I(x) = log 1/p(x).

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.1

Five days: probabilities multiply, surprises add



p = 0.9	×	p = 0.9	×	p = 0.9	×	p = 0.9	×	p = 0.9	**= 0.590**	probabilities multiply
0.152 bits	+	0.152 bits	+	0.152 bits	+	0.152 bits	+	0.152 bits	**= 0.76 bits**	surprises add: $\log_2 1/0.590 = 0.76$

> Five sunny days: the probability multiplies to $0.9^5 = 0.59$, and the surprise adds: $5 \times 0.152 = 0.76$ bits.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.1

Five days: probabilities multiply, surprises add



p = 0.9	×	p = 0.9	×	p = 0.9	×	p = 0.9	×	p = 0.9	**= 0.590**	probabilities multiply
0.152 bits	+	0.152 bits	+	0.152 bits	+	0.152 bits	+	0.152 bits	**= 0.76 bits**	surprises add: $\log_2 1/0.590 = 0.76$




p = 0.9	×	p = 0.9	×	p = 0.1	×	p = 0.9	×	p = 0.9	**= 0.0656**	probabilities multiply
0.152 bits	+	0.152 bits	+	3.32 bits	+	0.152 bits	+	0.152 bits	**= 3.93 bits**	surprises add: $\log_2 1/0.0656 = 3.93$

> Make one day rainy: the probability drops to 0.066 and the surprise jumps to 3.93 bits. The single rainy day carries most of it.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.1

Surprise in bits: $I(x) = \log_2 1/p(x)$



p(x) = 0.10

surprise = $\log_2 1/p$ = 3.32 bits

= ln 1/p = 2.30 nats

a rare outcome: a real surprise


one bit = the surprise of a fair coin flip

A certain outcome gives 0 bits, and halving the probability adds exactly one bit. Drag p, or try the buttons.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.1

Surprise in bits: $I(x) = \log_2 1/p(x)$



p(x) = 0.10

surprise = $\log_2 1/p$ = 3.32 bits

= ln $1/p$ = 2.30 nats

a rare outcome: a real surprise

one bit = the surprise of a fair coin flip


Base 2 gives bits; the natural log gives nats; 1 nat ≈ 1.44 bits.

> One bit is the surprise of a fair coin flip: one yes/no answer you could not guess. Information theory counts in bits, PyTorch in nats.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.2

Entropy: how surprised are you, on average?



$$\text{sunny: } \log_2 \frac{1}{0.9} = 0.15 \text{ bits}$$


$$\text{rainy: } \log_2 \frac{1}{0.1} = 3.32 \text{ bits}$$

> Each kind of weather has its surprise: 0.15 bits for sun, 3.32 for rain.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.2

Entropy: how surprised are you, on average?



$$\text{sunny: } \log_2 \frac{1}{0.9} = 0.15 \text{ bits}$$

$$\text{rainy: } \log_2 \frac{1}{0.1} = 3.32 \text{ bits}$$


$$H = 0.9\,(0.15) + 0.1\,(3.32) = 0.47 \text{ bits}$$

> How surprised are you on a typical day? 90% of days give 0.15 bits and 10% give 3.32: on average 0.47 bits.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.2

Entropy: how surprised are you, on average?



$$\text{sunny: } \log_2 \frac{1}{0.9} = 0.15 \text{ bits}$$

$$\text{rainy: } \log_2 \frac{1}{0.1} = 3.32 \text{ bits}$$

$$H = 0.9\,(0.15) + 0.1\,(3.32) = 0.47 \text{ bits}$$

$$H(p) = \sum_x p(x) \log_2 \frac{1}{p(x)} = \mathbb{E}_{x \sim p}\left[\log_2 \frac{1}{p(x)}\right]$$


> That weighted average of surprise is the entropy: the surprise you expect from one draw.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)


▶ Notebook 2.2

Why an expectation? It is the average over many days

days drawn from the forecast (latest 30), with their surprise




1,000 days



> Let days go by, each with its surprise: 0.15 bits if sunny, 3.32 if rainy. The running average settles on 0.47 bits. An expectation is a long-run average.

▶ Notebook 2.2


From certain to fully uncertain: between 0 and 1 bit



H = average surprise

$= 0.900 \times 0.15 + 0.100 \times 3.32$

$= 0.469$ bits




> Drag the chance of sun. A 50/50 forecast carries 1 bit, the most two outcomes can; 87.5/12.5 carries 0.54; a certain forecast, 0.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.2

Three kinds of weather: 1.5 bits




weather	p	$\log_2 \frac{1}{p}$
sunny	½	1 bit
cloudy	¼	2 bits
rainy	¼	2 bits

> Our town: sunny half the time, cloudy a quarter, rainy a quarter. Their surprises are 1, 2 and 2 bits.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.2

Three kinds of weather: 1.5 bits



weather	p	$\log_2 \frac{1}{p}$
sunny	½	1 bit
cloudy	¼	2 bits
rainy	¼	2 bits


$$H = \frac{1}{2}(1) + \frac{1}{4}(2) + \frac{1}{4}(2) = 1.5 \text{ bits}$$

> Average surprise: ½(1) + ¼(2) + ¼(2) = 1.5 bits per day.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.2

Entropy counts yes/no questions




Guess today's weather with yes/no questions, asking about the likeliest first.

> Find today's weather with yes/no questions. Ask "Is it sunny?" first: half the time you are done.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.2

Entropy counts yes/no questions



Guess today's weather with yes/no questions, asking about the likeliest first.


$$\tfrac{1}{2}(1) + \tfrac{1}{4}(2) + \tfrac{1}{4}(2) = 1.5 \text{ questions}$$

> Otherwise ask "Is it cloudy?". On average 1.5 questions: exactly the entropy.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.2

Entropy counts yes/no questions



Guess today's weather with yes/no questions, asking about the likeliest first.

$$\tfrac{1}{2}(1) + \tfrac{1}{4}(2) + \tfrac{1}{4}(2) = 1.5 \text{ questions}$$


Write each answer as a bit, yes = 0 and no = 1: sunny is **0**, cloudy **10**, rainy **11**. That is a code.

> Each answer is one bit. Writing the answers down gives a code for the weather: Section 2.3.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.3

Way 2: send the weather, in bits




> Turn the question game into messages: a weather station sends each day's weather to a receiver as 0s and 1s. Every bit costs, so we want as few as possible.

Coding story after Aurélien Géron, "A Short Introduction to Entropy, Cross-Entropy and KL-Divergence" (YouTube)

▶ Notebook 2.3

Way 2: send the weather, in bits



	sunny	cloudy	rainy
how often, p	½	¼	¼

> Sunny on half the days, cloudy on a quarter, rainy on a quarter: p = (½, ¼, ¼).

Coding story after Aurélien Géron, "A Short Introduction to Entropy, Cross-Entropy and KL-Divergence" (YouTube)

▶ Notebook 2.3

A code: a string of bits for each kind of day

weather	p	uniform code	bits
sunny	½	`00`	2
cloudy	¼	`01`	2
rainy	¼	`10`	2

The simplest code gives every kind of day the same length: 2 bits.

A code assigns each kind of day a string of bits. The uniform code spends 2 bits on every day.

Coding story after Aurélien Géron, "A Short Introduction to Entropy, Cross-Entropy and KL-Divergence" (YouTube)

▶ Notebook 2.3

A code: a string of bits for each kind of day

weather	p	uniform code	bits
sunny	½	`00`	2
cloudy	¼	`01`	2
rainy	¼	`10`	2

The simplest code gives every kind of day the same length: 2 bits.

Fair, but wasteful: half the days are sunny, yet a sunny day costs as much as a rare one.


> Wasteful: common days cost as much as rare ones. Could common days get shorter strings?

Coding story after Aurélien Géron, "A Short Introduction to Entropy, Cross-Entropy and KL-Divergence" (YouTube)

▶ Notebook 2.3

Prefix codes: the bits split up without separators

prefix code: sunny 0 · cloudy 10 · rainy 11



→ the receiver reads left to right; each day is unambiguous


> Give sunny 0, cloudy 10, rainy 11. No codeword is the start of another, so the receiver reads left to right and each day pops out.

Coding story after Aurélien Géron, "A Short Introduction to Entropy, Cross-Entropy and KL-Divergence" (YouTube)

▶ Notebook 2.3

Prefix codes: the bits split up without separators

prefix code: sunny 0 · cloudy 10 · rainy 11



→ the receiver reads left to right; each day is unambiguous

not a prefix code: sunny 0 · cloudy 01 · rainy 1



= sunny, then rainy?

or cloudy?

0 is the start of 01, so the receiver cannot tell.

> Break that rule, say 0, 01 and 1, and "01" could mean cloudy, or sunny then rainy. A prefix code avoids the confusion.

Coding story after Aurélien Géron, "A Short Introduction to Entropy, Cross-Entropy and KL-Divergence" (YouTube)

▶ Notebook 2.3

Huffman's recipe: merge the two rarest, repeat


Huffman (1952) builds the best prefix code from the probabilities:



Start with the three outcomes and their probabilities.

▶ Notebook 2.3

Huffman's recipe: merge the two rarest, repeat




Huffman (1952) builds the best prefix code from the probabilities:

1. Merge the two least likely outcomes into one node.

Merge the two rarest, cloudy and rainy, into one node: ¼ + ¼ = ½.

▶ Notebook 2.3

Huffman's recipe: merge the two rarest, repeat



Huffman (1952) builds the best prefix code from the probabilities:


1. Merge the two least likely outcomes into one node.

2. Repeat until one node is left.

Merge again: sunny and the new node make the root.

▶ Notebook 2.3

Huffman's recipe: merge the two rarest, repeat



Huffman (1952) builds the best prefix code from the probabilities:

1. Merge the two least likely outcomes into one node.

2. Repeat until one node is left.

3. Read each codeword off its path: left 0, right 1.

Read the codes off the tree: sunny 0, cloudy 10, rainy 11. The question game's code.

▶ Notebook 2.3

Each codeword is exactly as long as its surprise

weather	p	code	length	surprise
sunny	½	`0`	1	1 bit
cloudy	¼	`10`	2	2 bits
rainy	¼	`11`	2	2 bits

> Compare the code lengths with the surprises: 1, 2, 2 and 1, 2, 2. The best code spends exactly the surprise on each outcome.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.3

Each codeword is exactly as long as its surprise

weather	p	code	length	surprise
sunny	½	`0`	1	1 bit
cloudy	¼	`10`	2	2 bits
rainy	¼	`11`	2	2 bits

$$\text{average length} = \tfrac{1}{2}(1) + \tfrac{1}{4}(2) + \tfrac{1}{4}(2) = 1.5 \text{ bits} = H(p)$$


Shannon (1948): no code averages fewer than H(p) bits. Huffman reaches it when every probability is a power of ½, and comes within 1 bit otherwise.

> So the average message length is the average surprise, the entropy: 1.5 bits per day instead of 2.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.4

A code built on a wrong belief




The station was designed for a town where rain is the common weather: q = (¼, ¼, ½).

> Suppose the station's code was built for another town, where rain is common: q = (¼, ¼, ½). Our weather is still p = (½, ¼, ¼).

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.4

A code built on a wrong belief



The station was designed for a town where rain is the common weather: q = (¼, ¼, ½).

weather	q	their code	$\log_2 \frac{1}{q}$
sunny	¼	`10`	2 bits
cloudy	¼	`11`	2 bits
rainy	½	`0`	1 bit

> Huffman on q gives rainy the short codeword: sunny 10, cloudy 11, rainy 0. Each length is $\log_2 1/q$: the surprise under the wrong belief.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.4

Cross-entropy: the average cost when the code expects q

$$H(p, q) = \sum_{x} p(x) \log_2 \frac{1}{q(x)}$$

> Cross-entropy: average, over our real weather p, of the code lengths built from q.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.4

Cross-entropy: the average cost when the code expects q

$$H(p, q) = \sum_x p(x) \log_2 \frac{1}{q(x)}$$

$$= \tfrac{1}{2}(2) + \tfrac{1}{4}(2) + \tfrac{1}{4}(1) = 1.75 \text{ bits per day}$$

> ½ × 2 + ¼ × 2 + ¼ × 1 = 1.75 bits per day, against 1.5 with the right code.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.4

Cross-entropy: the average cost when the code expects q

$$H(p, q) = \sum_x p(x) \log_2 \frac{1}{q(x)}$$

$$= \tfrac{1}{2}(2) + \tfrac{1}{4}(2) + \tfrac{1}{4}(1) = 1.75 \text{ bits per day}$$

weights from the truth p: how often each day happens

lengths from the belief q: what each day costs

> The two play different roles: p decides how often, q decides how much. Entropy is the special case q = p.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.4

Watch it happen: our weather through both codes

the latest days, and the bits each code sends



code for p	11	11	11	0	0	0	11	10
code for q	0	0	0	10	10	10	0	11


1,000 days sent

code for p: 1,540 bits, 1.540 per day

code for q: 1,733 bits, 1.733 per day

wasted: 0.193 bits per day

bits per day so far



> Send days of our weather through both codes. With 10, 100, 1,000 days the right code settles at 1.5 bits per day and the wrong one at 1.75: 0.25 bits per day wasted.

▶ Notebook 2.4

KL divergence: the extra bits we waste

$$\mathrm{KL}(p \,\|\, q) = H(p, q) - H(p) = 1.75 - 1.5 = 0.25 \text{ bits}$$

> The KL divergence is the waste: cross-entropy minus entropy, here 0.25 bits per day.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.4

KL divergence: the extra bits we waste

$$\mathrm{KL}(p \,\|\, q) = H(p, q) - H(p) = 1.75 - 1.5 = 0.25 \text{ bits}$$

$$= \sum_x p(x) \left(\log_2 \frac{1}{q(x)} - \log_2 \frac{1}{p(x)} \right)$$

> Both are sums over the same outcomes with the same weights p(x), so subtract term by term.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.4

KL divergence: the extra bits we waste

$$\mathrm{KL}(p \,\|\, q) = H(p, q) - H(p) = 1.75 - 1.5 = 0.25 \text{ bits}$$

$$= \sum_x p(x) \left(\log_2 \frac{1}{q(x)} - \log_2 \frac{1}{p(x)} \right)$$

$$= \sum_x p(x) \log_2 \frac{p(x)}{q(x)} = \tfrac{1}{2}(1) + \tfrac{1}{4}(0) + \tfrac{1}{4}(-1) = 0.25$$

A difference of logs is the log of a ratio: KL(p‖q) = Σ p log p/q. For our weather ½(1) + ¼(0) + ¼(−1) = 0.25.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.4

Never negative, and zero only for the right code

$$H(p, q) \geq H(p) \quad \Longrightarrow \quad \mathrm{KL}(p \,\|\, q) \geq 0$$

No code averages fewer than H(p) bits, so a code built for q can only waste bits.

> The right code is the best possible, so any other code can only cost more: KL is never negative.

▶ Notebook 2.4

Never negative, and zero only for the right code

$$H(p, q) \geq H(p) \quad \Longrightarrow \quad \mathrm{KL}(p \,\|\, q) \geq 0$$

No code averages fewer than H(p) bits, so a code built for q can only waste bits.

$$\mathrm{KL}(p \,\|\, q) = 0 \quad \Longleftrightarrow \quad q = p$$

Single terms can be negative, as rainy's was; the weighted total cannot. Part 4 proves KL ≥ 0 for any distributions in one line.

> And it is zero only when the belief matches the truth.

▶ Notebook 2.4

Which one is the truth? KL is not symmetric

“Oliver is 10 cm taller than Oscar” and “Oscar is 10 cm shorter than Oliver”: the description depends on who you measure from. With KL even the size changes.


> Which distribution is the reference? For heights, only the wording changes. For KL, the number does.

Story after Jia-Bin Huang, “Fantastic KL Divergence and How to (Actually) Compute It” (YouTube)

▶ Notebook 2.4

Which one is the truth? KL is not symmetric

"Oliver is 10 cm taller than Oscar" and "Oscar is 10 cm shorter than Oliver": the description depends on who you measure from. With KL even the size changes.



	bits wasted per day
our weather, their code	1.84


> A sunny town expects sun 98% of the time, so its code gives cloudy and rainy 6.6-bit codewords. On our weather, where they come half the time, that wastes 1.84 bits a day: KL(p‖q).

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.4

Which one is the truth? KL is not symmetric

"Oliver is 10 cm taller than Oscar" and "Oscar is 10 cm shorter than Oliver": the description depends on who you measure from. With KL even the size changes.




	bits wasted per day
our weather, their code	1.84
their weather, our code	0.86

> Swap the roles: our code on their weather wastes only 0.86 bits a day, KL(q‖p). The truth supplies the weights, so swapping changes the answer.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.5

A classifier outputs a distribution




> An image goes into a classifier; out comes a distribution over four labels: ape 0.05, bear 0.05, cat 0.60, dog 0.30. Call it q.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.5

A classifier outputs a distribution



> The true label is a distribution too: all of its probability on cat. Call it p.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.5

The loss is the model's surprise at the true label

$$H(p, q) = \sum_{y} p(y) \log \frac{1}{q(y)}$$

> The cross-entropy between the label p and the prediction q: the label supplies the weights.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.5

The loss is the model's surprise at the true label

$$H(p, q) = \sum_{y} p(y) \log \frac{1}{q(y)}$$

$$= 0 \cdot \log \frac{1}{0.05} + 0 \cdot \log \frac{1}{0.05} + 1 \cdot \log \frac{1}{0.6} + 0 \cdot \log \frac{1}{0.3}$$

> Write out all four terms. Three are multiplied by zero.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.5

The loss is the model's surprise at the true label

$$H(p, q) = \sum_{y} p(y) \log \frac{1}{q(y)}$$

$$= 0 \cdot \log \tfrac{1}{0.05} + 0 \cdot \log \tfrac{1}{0.05} + 1 \cdot \log \tfrac{1}{0.6} + 0 \cdot \log \tfrac{1}{0.3}$$

$$= -\log q(\text{cat}) = 0.51 \text{ nats} = 0.74 \text{ bits}$$

> What is left is −log q(cat): the model's surprise at the true label. That is the loss you have been minimising all along.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.5

With a one-hot label, cross-entropy is exactly KL

$$\mathrm{KL}(p \,\|\, q) = H(p, q) - H(p)$$

> From Section 2.4: KL is cross-entropy minus entropy.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.5

With a one-hot label, cross-entropy is exactly KL

$$\mathrm{KL}(p \,\|\, q) = H(p, q) - H(p)$$

$$H(p) = 1 \cdot \log \tfrac{1}{1} + 0 + 0 + 0 = 0$$

A one-hot label is certain, so it has no surprise (0 · log 1/0 counts as 0).

> The entropy of a one-hot label is 0: a certain outcome carries no surprise.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.5

With a one-hot label, cross-entropy is exactly KL

$$\mathrm{KL}(p \,\|\, q) = H(p, q) - H(p)$$

$$H(p) = 1 \cdot \log \tfrac{1}{1} + 0 + 0 + 0 = 0$$

A one-hot label is certain, so it has no surprise (0 · log 1/0 counts as 0).


```
F.cross_entropy(logits, target)   # 0.511 nats: here equal to KL(p ‖ q)
```

> So the classification loss is exactly KL(label ‖ prediction). PyTorch computes it in nats: 0.511 for our cat.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.5

The loss curve is the surprise curve




The loss is −ln q(true class): the surprise curve of Section 2.1, in nats.

Plot the loss against the probability the model gives to the true class: it is the surprise curve again.

▶ Notebook 2.5

The loss curve is the surprise curve




The loss is −ln q(true class): the surprise curve of Section 2.1, in nats.

Confident and wrong is expensive: q(true) = 0.01 costs 4.6 nats; q(true) = 0.6 costs 0.51.

Training pushes q(true) up the curve towards 1. Being confidently wrong is punished hardest.

▶ Notebook 2.5

Soft labels: cross-entropy = entropy + KL




$$H(p,q) = \underbrace{H(p)}_{0.72} + \underbrace{\mathrm{KL}(p \,\|\, q)}_{0.05} = 0.77$$

> A soft label: a teacher network says cat 0.75, dog 0.20, bear 0.03, ape 0.02. Now H(p) is not 0.

▶ Notebook 2.5

Soft labels: cross-entropy = entropy + KL




$$H(p, q) = \underbrace{H(p)}_{0.72} + \underbrace{\mathrm{KL}(p \,\|\, q)}_{0.05} = 0.77$$

In knowledge distillation a small student learns from a big teacher's predictions. H(p) is fixed by the teacher, so lowering the cross-entropy lowers the KL by exactly as much.

> Training changes only the student q, so minimising cross-entropy is still minimising KL(teacher ‖ student).

▶ Notebook 2.6


The data: twenty days of weather



> Now real data: twenty recorded days. We want a model q for the weather. Which q explains these days best?

▶ Notebook 2.6

The data: twenty days of weather



Counts: 10 sunny, 5 cloudy, 5 rainy. Frequencies $\hat{p} = (½, ¼, ¼)$.

> Count them: 10 sunny, 5 cloudy, 5 rainy. Their frequencies are $\hat{p} = (½, ¼, ¼)$.

▶ Notebook 2.6

How probable are these twenty days under q?

$$p(\mathcal{D} \mid q) = \prod_{i=1}^{20} q(x_i) = q(\text{sunny})^{10}\, q(\text{cloudy})^{5}\, q(\text{rainy})^{5}$$

> Under the model the days are independent, so the probability of the record multiplies: one factor per day. Equal days give powers.

▶ Notebook 2.6

How probable are these twenty days under q?

$$p(\mathcal{D} \mid q) = \prod_{i=1}^{20} q(x_i) = q(\text{sunny})^{10}\, q(\text{cloudy})^{5}\, q(\text{rainy})^{5}$$

$$q = (\tfrac{1}{4}, \tfrac{1}{4}, \tfrac{1}{2}): \quad (\tfrac{1}{4})^{10}(\tfrac{1}{4})^{5}(\tfrac{1}{2})^{5} = 2^{-35} \approx 2.9 \times 10^{-11}$$

Take the wrong belief q = (¼, ¼, ½): the record has probability 2^{-35}, about 3 in a hundred billion.

▶ Notebook 2.6

How probable are these twenty days under q?

$$p(\mathcal{D} \mid q) = \prod_{i=1}^{20} q(x_i) = q(\text{sunny})^{10}\, q(\text{cloudy})^{5}\, q(\text{rainy})^{5}$$

$$q = (\tfrac{1}{4}, \tfrac{1}{4}, \tfrac{1}{2}): \quad (\tfrac{1}{4})^{10}(\tfrac{1}{4})^{5}(\tfrac{1}{2})^{5} = 2^{-35} \approx 2.9 \times 10^{-11}$$

$$\log_2 p(\mathcal{D} \mid q) = -35$$

Tiny numbers are why we take logs: $\log_2 p(\mathcal{D} \mid q) = -35$. The whole record is 35 bits of surprise.

▶ Notebook 2.6

Per day: the data's average surprise under q

$$-\frac{1}{20}\log_2 p(\mathcal{D} \mid q) = \frac{1}{20}\sum_{i=1}^{20}\log_2 \frac{1}{q(x_i)} = \frac{35}{20} = 1.75 \text{ bits}$$

Divide by the number of days: minus the log-likelihood per day is the average surprise of a recorded day under q. Here 1.75 bits.

▶ Notebook 2.6

Per day: the data's average surprise under q

$$-\frac{1}{20}\log_2 p(\mathcal{D} \mid q) = \frac{1}{20}\sum_{i=1}^{20}\log_2 \frac{1}{q(x_i)} = \frac{35}{20} = 1.75 \text{ bits}$$

Have we seen 1.75 before?

> Section 2.4 had a 1.75: the average code length when the code expects q but the weather is p.

▶ Notebook 2.6

Per day: the data's average surprise under q

$$-\frac{1}{20}\log_2 p(\mathcal{D} \mid q) = \frac{1}{20}\sum_{i=1}^{20}\log_2 \frac{1}{q(x_i)} = \frac{35}{20} = 1.75 \text{ bits}$$

Have we seen 1.75 before?

It is the cross-entropy H(p, q) of Section 2.4.

> Same number: the data's average surprise under q is a cross-entropy. Why?

▶ Notebook 2.6

Why: group the days by weather

$$\frac{1}{20}\sum_{i=1}^{20}\log_2\frac{1}{q(x_i)} = \frac{1}{20}\left[10\log_2\frac{1}{q(\text{sunny})} + 5\log_2\frac{1}{q(\text{cloudy})} + 5\log_2\frac{1}{q(\text{rainy})}\right]$$

> Sort the twenty terms by weather: ten equal sunny terms, five cloudy, five rainy. Equal terms add up to count × term.

▶ Notebook 2.6

Why: group the days by weather

$$\frac{1}{20}\sum_{i=1}^{20}\log_2\frac{1}{q(x_i)} = \frac{1}{20}\left[10\log_2\frac{1}{q(\text{sunny})} + 5\log_2\frac{1}{q(\text{cloudy})} + 5\log_2\frac{1}{q(\text{rainy})}\right]$$

$$= \frac{10}{20}\log_2\frac{1}{q(\text{sunny})} + \frac{5}{20}\log_2\frac{1}{q(\text{cloudy})} + \frac{5}{20}\log_2\frac{1}{q(\text{rainy})}$$

Divide by 20: the counts become the frequencies, 10/20, 5/20, 5/20.

▶ Notebook 2.6

Why: group the days by weather

$$\frac{1}{20}\sum_{i=1}^{20}\log_2\frac{1}{q(x_i)} = \frac{1}{20}\left[10\log_2\frac{1}{q(\text{sunny})} + 5\log_2\frac{1}{q(\text{cloudy})} + 5\log_2\frac{1}{q(\text{rainy})}\right]$$

$$= \frac{10}{20}\log_2\frac{1}{q(\text{sunny})} + \frac{5}{20}\log_2\frac{1}{q(\text{cloudy})} + \frac{5}{20}\log_2\frac{1}{q(\text{rainy})}$$

$$= \sum_x \hat{p}(x)\log_2\frac{1}{q(x)} = H(\hat{p},\, q)$$

> Those frequencies are p̂. The data's average surprise is the cross-entropy H(p̂, q): the data became a distribution.

▶ Notebook 2.6

The data's surprise = entropy of the data + KL

$$H(\hat{p}, q) = \underbrace{H(\hat{p})}_{\text{fixed by the data}} + \underbrace{\mathrm{KL}(\hat{p} \,\|\, q)}_{\text{depends on } q}$$

Section 2.4's split: cross-entropy = entropy + KL. The entropy of the data is fixed; only the KL depends on the model.

▶ Notebook 2.6

The data's surprise = entropy of the data + KL

$$H(\hat{p}, q) = \underbrace{H(\hat{p})}_{\text{fixed by the data}} + \underbrace{\mathrm{KL}(\hat{p} \,\|\, q)}_{\text{depends on } q}$$


$$1.75 = 1.5 + 0.25 \qquad \text{for } q = (\tfrac{1}{4}, \tfrac{1}{4}, \tfrac{1}{2})$$

For q = (¼, ¼, ½): 1.75 = 1.5 + 0.25. The model can only remove the KL part.

▶ Notebook 2.6


Find the q that makes the data least surprising

data frequencies $\hat{p}$ and the model q



the data's average surprise under q

= 1.750 bits per day



$KL(\hat{p} \,\|\, q) = 0.250$ bits

$\log_2 p(\mathcal{D} \mid q) = -35.0$: 32× less probable than under $\hat{p}$

> Move q. The teal part, $H(\hat{p})$, never changes; the red part, the KL, shrinks to zero exactly at $q = \hat{p} = (½, ¼, ¼)$. Maximum likelihood matches the frequencies.

▶ Notebook 2.6

Maximum likelihood minimises KL(data ‖ model)

$$\max_\theta \ \frac{1}{N}\sum_{i=1}^{N} \log q_\theta(x_i) \iff \min_\theta \ H(\hat{p}, q_\theta) \iff \min_\theta \ \mathrm{KL}(\hat{p} \,\|\, q_\theta)$$

> Nothing in the derivation was special to weather: maximising the likelihood minimises KL(data ‖ model), with the data first.

▶ Notebook 2.6

Maximum likelihood minimises KL(data ‖ model)

$$\max_\theta \ \frac{1}{N}\sum_{i=1}^{N} \log q_\theta(x_i) \iff \min_\theta \ H(\hat{p}, q_\theta) \iff \min_\theta \ \mathrm{KL}(\hat{p} \,\|\, q_\theta)$$

Any model, any data: the average log-likelihood is minus the data's cross-entropy, and the data's entropy is fixed.

> With more data $\hat{p}$ approaches the true distribution of the data, so the model is pulled towards it in this KL.

▶ Notebook 2.6

Maximum likelihood minimises KL(data ‖ model)

$$\max_\theta \ \frac{1}{N}\sum_{i=1}^{N} \log q_\theta(x_i) \iff \min_\theta \ H(\hat{p}, q_\theta) \iff \min_\theta \ \mathrm{KL}(\hat{p} \,\|\, q_\theta)$$

Any model, any data: the average log-likelihood is minus the data's cross-entropy, and the data's entropy is fixed.

A classifier's training set does the same, label by label: each example costs $-\log q_\theta(y_i \mid x_i)$, the surprise at its label.

> The classification loss of 2.5 is the same story, averaged over a training set: training a classifier is maximum likelihood.

▶ Notebook 2.6

Takeaway: KL is everywhere in machine learning

training …	minimises	first argument
a classifier with cross-entropy	KL(labels ‖ predictions)	the labels
a model by maximum likelihood	KL(data ‖ model)	the data
a student by distillation	KL(teacher ‖ student)	the teacher
a chatbot with RLHF	reward − KL(policy ‖ reference)	the policy
an approximation by VI	KL(q ‖ posterior)	our approximation q

> The same quantity hides in many losses. Variational inference minimises one with q first; Section 3.6 shows why the order matters.

Story after Jia-Bin Huang, "Fantastic KL Divergence and How to (Actually) Compute It" (YouTube)

▶ Notebook 2.7

From sums to integrals

$$\text{discrete:} \qquad \mathrm{KL}(p\|q) = \sum_x p(x) \log \frac{p(x)}{q(x)}$$

$$\text{continuous:} \qquad \mathrm{KL}(p\|q) = \int p(\theta) \log \frac{p(\theta)}{q(\theta)}\, d\theta$$

For a continuous unknown like θ, the sum over grades becomes an integral over θ.

▶ Notebook 2.7

From sums to integrals

$$\text{discrete:} \qquad \mathrm{KL}(p\|q) = \sum_x p(x) \log \frac{p(x)}{q(x)}$$

$$\text{continuous:} \qquad \mathrm{KL}(p\|q) = \int p(\theta) \log \frac{p(\theta)}{q(\theta)}\, d\theta$$

Probabilities become densities and the sum becomes an integral; the recipe is the same weighted average of log ratios.

> Nothing else changes. KL is never negative either; Part 4 proves it, where we need it.


▶ Notebook 2.7

Bernoulli: a coin with P(x = 1) = a

p = Bernoulli(0.8)



q = Bernoulli(0.5)



$$p(x) = a^x(1-a)^{1-x}, \qquad x \in \{0, 1\}$$

$$p(1) = a, \qquad p(0) = 1 - a$$

Bernoulli(a): x is 1 with probability a and 0 otherwise. Here p = Bernoulli(0.8), q = Bernoulli(0.5).


▶ Notebook 2.7

Bernoulli: a coin with P(x = 1) = a

p = Bernoulli(0.8)



q = Bernoulli(0.5)



$$p(x) = a^x(1-a)^{1-x}, \qquad x \in \{0, 1\}$$

$$p(1) = a, \qquad p(0) = 1 - a$$

Your turn: write $\mathrm{KL}(p \,\|\, q)$ for p = Bernoulli(a) and q = Bernoulli(b).

Your turn: put the PMF into Σ_x p(x) log p(x)/q(x). There are only two values of x.

▶ Notebook 2.7

KL between two Bernoullis, step by step

$$\mathrm{KL}(p \,\|\, q) = \sum_{x \in \{0,1\}} p(x) \log \frac{p(x)}{q(x)}$$

Start from the definition: a sum over the values of x.

▶ Notebook 2.7

KL between two Bernoullis, step by step

$$\mathrm{KL}(p \,\|\, q) = \sum_{x \in \{0,1\}} p(x) \log \frac{p(x)}{q(x)}$$

$$= p(1) \log \frac{p(1)}{q(1)} + p(0) \log \frac{p(0)}{q(0)}$$

There are two: write out the term for x = 1 and the term for x = 0.

▶ Notebook 2.7

KL between two Bernoullis, step by step

$$\mathrm{KL}(p \,\|\, q) = \sum_{x \in \{0,1\}} p(x) \log \frac{p(x)}{q(x)}$$

$$= p(1) \log \frac{p(1)}{q(1)} + p(0) \log \frac{p(0)}{q(0)}$$

$$= a \log \frac{a}{b} + (1 - a) \log \frac{1 - a}{1 - b}$$

Substitute p(1) = a, p(0) = 1 − a, and b for q.

▶ Notebook 2.7

KL between two Bernoullis, step by step

$$\mathrm{KL}(p \,\|\, q) = \sum_{x \in \{0,1\}} p(x) \log \frac{p(x)}{q(x)}$$

$$= p(1) \log \frac{p(1)}{q(1)} + p(0) \log \frac{p(0)}{q(0)}$$


$$= a \log \frac{a}{b} + (1-a) \log \frac{1-a}{1-b}$$

$$a = 0.8,\ b = 0.5: \quad 0.8 \log 1.6 + 0.2 \log 0.4 = 0.376 - 0.183 = 0.19$$

For a = 0.8 and b = 0.5: 0.38 − 0.18 = 0.19 nats.

▶ Notebook 2.7

Categorical: K outcomes with probabilities $\pi_1, \ldots, \pi_K$



$$p(x = k) = \pi_k, \quad k = 1, \ldots, K$$


$$\pi_k \geq 0, \qquad \sum_k \pi_k = 1$$

Our grades: K = 3, with π = (0.4, 0.1, 0.5) this year and ρ = (0.1, 0.4, 0.5) last year.

> Categorical(π): one of K outcomes, outcome k with probability π_k. Our grade distributions are categorical, with K = 3.

▶ Notebook 2.7

Categorical: K outcomes with probabilities $\pi_1, \ldots, \pi_K$



$$p(x = k) = \pi_k, \quad k = 1, \ldots, K$$

$$\pi_k \geq 0, \qquad \sum_k \pi_k = 1$$

Our grades: K = 3, with π = (0.4, 0.1, 0.5) this year and ρ = (0.1, 0.4, 0.5) last year.

Your turn: write $\mathrm{KL}(p \,\|\, q)$ for p = Categorical(π) and q = Categorical(ρ).

> Your turn: the same recipe, now with K terms.

▶ Notebook 2.7

KL between two categoricals, step by step

$$\mathrm{KL}(p \,\|\, q) = \sum_{k=1}^{K} p(k) \log \frac{p(k)}{q(k)} = \sum_{k=1}^{K} \pi_k \log \frac{\pi_k}{\rho_k}$$

Substitute p(k) = π_k and q(k) = ρ_k: one term per outcome.

▶ Notebook 2.7

KL between two categoricals, step by step

$$\mathrm{KL}(p \,\|\, q) = \sum_{k=1}^{K} p(k) \log \frac{p(k)}{q(k)} = \sum_{k=1}^{K} \pi_k \log \frac{\pi_k}{\rho_k}$$

$$K = 2: \quad \pi = (a,\, 1-a),\ \rho = (b,\, 1-b) \ \Rightarrow\ a \log \frac{a}{b} + (1-a) \log \frac{1-a}{1-b}$$

With K = 2 this is exactly the Bernoulli formula: a Bernoulli is a categorical with two outcomes.

▶ Notebook 2.7

KL between two categoricals, step by step

$$\mathrm{KL}(p \,\|\, q) = \sum_{k=1}^{K} p(k) \log \frac{p(k)}{q(k)} = \sum_{k=1}^{K} \pi_k \log \frac{\pi_k}{\rho_k}$$


$$K = 2: \quad \pi = (a,\, 1-a),\ \rho = (b,\, 1-b) \ \Rightarrow\ a \log \frac{a}{b} + (1-a) \log \frac{1-a}{1-b}$$

$$\text{grades: } 0.4 \log 4 + 0.1 \log \tfrac{1}{4} + 0.5 \log 1 = 0.42$$

For the grades: 0.42 nats, the number we built by hand in Way 1.

▶ Notebook 2.7

The Gaussian: a bell with mean μ and width σ




$$\mathcal{N}(\theta; \mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}}\, e^{-(\theta-\mu)^2/(2\sigma^2)}$$

q = N(μ, σ²) is our approximation and p = N(m, s²) the pretend posterior. Shown: q = N(0, 2²), p = N(1, 4²).

> The Gaussian density: a normalising constant times e to the minus squared distance from the mean, scaled by the variance.

▶ Notebook 2.7

The Gaussian: a bell with mean μ and width σ



$$\mathcal{N}(\theta; \mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}}\, e^{-(\theta-\mu)^2/(2\sigma^2)}$$

q = N(μ, σ²) is our approximation and p = N(m, s²) the pretend posterior. Shown: q = N(0, 2²), p = N(1, 4²).

Your turn: work out $\mathrm{KL}(q \,\|\, p)$. Start with log q(θ) – log p(θ).

> Your turn: work out KL(q ‖ p). The log makes the exponentials easy.

▶ Notebook 2.7

Gaussian KL, step 1: the log ratio

$$\log q(\theta) = -\tfrac{1}{2}\log(2\pi\sigma^2) - \frac{(\theta-\mu)^2}{2\sigma^2}$$

Take the log of q's density: the exponential disappears.

▶ Notebook 2.7

Gaussian KL, step 1: the log ratio

$$\log q(\theta) = -\tfrac{1}{2}\log(2\pi\sigma^2) - \frac{(\theta-\mu)^2}{2\sigma^2}$$

$$\log p(\theta) = -\tfrac{1}{2}\log(2\pi s^2) - \frac{(\theta-m)^2}{2s^2}$$

The same for p, with its own mean m and width s.

▶ Notebook 2.7

Gaussian KL, step 1: the log ratio

$$\log q(\theta) = -\tfrac{1}{2}\log(2\pi\sigma^2) - \frac{(\theta-\mu)^2}{2\sigma^2}$$

$$\log p(\theta) = -\tfrac{1}{2}\log(2\pi s^2) - \frac{(\theta-m)^2}{2s^2}$$

$$\log q(\theta) - \log p(\theta) = \log\frac{s}{\sigma} - \frac{(\theta-\mu)^2}{2\sigma^2} + \frac{(\theta-m)^2}{2s^2}$$

Subtract. The 2π terms cancel, and the two log widths combine into log(s/σ).

▶ Notebook 2.7

Gaussian KL, step 2: average over q

$$\mathrm{KL}(q\|p) = \mathbb{E}_q\left[\log\frac{s}{\sigma} - \frac{(\theta-\mu)^2}{2\sigma^2} + \frac{(\theta-m)^2}{2s^2}\right]$$

Average the log ratio over θ drawn from q. The first term is a constant, so it stays as it is.

▶ Notebook 2.7

Gaussian KL, step 2: average over q

$$\mathrm{KL}(q\|p) = \mathbb{E}_q\left[\log\frac{s}{\sigma} - \frac{(\theta-\mu)^2}{2\sigma^2} + \frac{(\theta-m)^2}{2s^2}\right]$$

$$\mathbb{E}_q\left[(\theta-\mu)^2\right] = \sigma^2 \quad \Rightarrow \quad -\frac{\sigma^2}{2\sigma^2} = -\frac{1}{2}$$

The second: the average squared distance from q's own mean is its variance, σ². That term becomes −½.

▶ Notebook 2.7

Gaussian KL, step 2: average over q

$$\mathrm{KL}(q\|p) = \mathbb{E}_q\Big[\log\frac{s}{\sigma} - \frac{(\theta-\mu)^2}{2\sigma^2} + \frac{(\theta-m)^2}{2s^2}\Big]$$

$$\mathbb{E}_q\big[(\theta-\mu)^2\big] = \sigma^2 \quad \Rightarrow \quad -\frac{\sigma^2}{2\sigma^2} = -\frac{1}{2}$$

$$\mathbb{E}_q\big[(\theta-m)^2\big] = \mathbb{E}_q\big[\big((\theta-\mu)+(\mu-m)\big)^2\big] = \sigma^2 + (\mu-m)^2$$

The third: split θ − m into (θ − μ) + (μ − m). The cross term averages to zero, leaving σ² + (μ − m)².

▶ Notebook 2.7

Gaussian KL, step 2: average over q

$$\mathrm{KL}(q\|p) = \mathbb{E}_q\Big[\log\frac{s}{\sigma} - \frac{(\theta-\mu)^2}{2\sigma^2} + \frac{(\theta-m)^2}{2s^2}\Big]$$

$$\mathbb{E}_q\big[(\theta-\mu)^2\big] = \sigma^2 \quad \Rightarrow \quad -\frac{\sigma^2}{2\sigma^2} = -\frac{1}{2}$$


$$\mathbb{E}_q\big[(\theta-m)^2\big] = \mathbb{E}_q\big[\big((\theta-\mu)+(\mu-m)\big)^2\big] = \sigma^2 + (\mu-m)^2$$

$$\mathrm{KL}(q\|p) = \log\frac{s}{\sigma} + \frac{\sigma^2+(\mu-m)^2}{2s^2} - \frac{1}{2}$$

Together: a width term, the squared shift between the means, and −½. That is the Gaussian KL.

▶ Notebook 2.7

A diagonal Gaussian: one bell per coordinate



$$q(\boldsymbol{\theta}) = \prod_{j=1}^{d} \mathcal{N}(\theta_j;\, \mu_j,\, \sigma_j^2)$$


$$p(\boldsymbol{\theta}) = \prod_{j=1}^{d} \mathcal{N}(\theta_j;\, m_j,\, s_j^2)$$

Independent coordinates, so the contours are axis-aligned. Our network's q is exactly this, with d = 7.

A diagonal Gaussian is a product of one-dimensional Gaussians: each coordinate has its own mean and width.

▶ Notebook 2.7

A diagonal Gaussian: one bell per coordinate



$$q(\boldsymbol{\theta}) = \prod_{j=1}^{d} \mathcal{N}(\theta_j;\, \mu_j,\, \sigma_j^2)$$

$$p(\boldsymbol{\theta}) = \prod_{j=1}^{d} \mathcal{N}(\theta_j;\, m_j,\, s_j^2)$$

Independent coordinates, so the contours are axis-aligned. Our network's q is exactly this, with d = 7.

Your turn: what is KL(q ‖ p) in d dimensions?

> Your turn: what happens to the log of a product?

▶ Notebook 2.7

KL between diagonal Gaussians, step by step

$$\log \frac{q(\boldsymbol{\theta})}{p(\boldsymbol{\theta})} = \sum_{j=1}^{d} \log \frac{\mathcal{N}(\theta_j; \mu_j, \sigma_j^2)}{\mathcal{N}(\theta_j; m_j, s_j^2)}$$

The log of a product is a sum of logs: the log ratio splits into one term per coordinate.

▶ Notebook 2.7

KL between diagonal Gaussians, step by step

$$\log \frac{q(\boldsymbol{\theta})}{p(\boldsymbol{\theta})} = \sum_{j=1}^{d} \log \frac{\mathcal{N}(\theta_j; \mu_j, \sigma_j^2)}{\mathcal{N}(\theta_j; m_j, s_j^2)}$$

$$\mathrm{KL}(q \| p) = \sum_{j=1}^{d} \mathrm{KL}\big(\mathcal{N}(\mu_j, \sigma_j^2) \,\|\, \mathcal{N}(m_j, s_j^2)\big)$$

Each term involves only θ_j, so its average under q is a one-dimensional Gaussian KL. The total is their sum.

▶ Notebook 2.7

KL between diagonal Gaussians, step by step

$$\log \frac{q(\boldsymbol{\theta})}{p(\boldsymbol{\theta})} = \sum_{j=1}^{d} \log \frac{\mathcal{N}(\theta_j; \mu_j, \sigma_j^2)}{\mathcal{N}(\theta_j; m_j, s_j^2)}$$

$$\mathrm{KL}(q\|p) = \sum_{j=1}^{d} \mathrm{KL}\big(\mathcal{N}(\mu_j, \sigma_j^2) \,\|\, \mathcal{N}(m_j, s_j^2)\big)$$

$$p = \mathcal{N}(0, I): \qquad \mathrm{KL}(q\|p) = \sum_{j=1}^{d} \Big[-\log \sigma_j + \frac{\sigma_j^2 + \mu_j^2}{2} - \frac{1}{2} \Big]$$

> Against a N(0, 1) prior on every coordinate: exactly the KL term in our network's ELBO, and in every VAE.

▶ Notebook 2.7

The same formulas in PyTorch: Bernoulli and categorical

```python
@register_kl(Bernoulli, Bernoulli)
def _kl_bernoulli_bernoulli(p, q):
    t1 = p.probs * (softplus(-q.logits) - softplus(-p.logits))
    t2 = (1 - p.probs) * (softplus(q.logits) - softplus(p.logits))
    return t1 + t2

@register_kl(Categorical, Categorical)
def _kl_categorical_categorical(p, q):
    t = p.probs * (p.logits - q.logits)
    return t.sum(-1)
```

From PyTorch's source, torch/distributions/kl.py: the Bernoulli and categorical KLs.

PyTorch source: torch/distributions/kl.py

▶ Notebook 2.7

The same formulas in PyTorch: Bernoulli and categorical

```python
@register_kl(Bernoulli, Bernoulli)
def _kl_bernoulli_bernoulli(p, q):
    t1 = p.probs * (softplus(-q.logits) - softplus(-p.logits))
    t2 = (1 - p.probs) * (softplus(q.logits) - softplus(p.logits))
    return t1 + t2

@register_kl(Categorical, Categorical)
def _kl_categorical_categorical(p, q):
    t = p.probs * (p.logits - q.logits)
    return t.sum(-1)
```

softplus(−logit) = −log(probability), so t1 = a log(a/b) and t2 = (1 − a) log((1 − a)/(1 − b)). For a categorical, logits are log-probabilities: $\Sigma\ \pi_k\ (\log \pi_k - \log \rho_k)$.

> Read them against our formulas: a log(a/b) + (1 − a) log((1 − a)/(1 − b)), and $\Sigma_k\ \pi_k \log(\pi_k/\rho_k)$. The same.

PyTorch source: torch/distributions/kl.py

▶ Notebook 2.7

The same formulas in PyTorch: Gaussian and diagonal

```python
@register_kl(Normal, Normal)
def _kl_normal_normal(p, q):
    var_ratio = (p.scale / q.scale).pow(2)
    t1 = ((p.loc - q.loc) / q.scale).pow(2)
    return 0.5 * (var_ratio + t1 - 1 - var_ratio.log())

@register_kl(Independent, Independent)
def _kl_independent_independent(p, q):
    result = kl_divergence(p.base_dist, q.base_dist)
    return _sum_rightmost(result, p.reinterpreted_batch_ndims)
```

> PyTorch's Gaussian KL, and the rule for independent coordinates.

PyTorch source: torch/distributions/kl.py

▶ Notebook 2.7

The same formulas in PyTorch: Gaussian and diagonal

```python
@register_kl(Normal, Normal)
def _kl_normal_normal(p, q):
    var_ratio = (p.scale / q.scale).pow(2)
    t1 = ((p.loc - q.loc) / q.scale).pow(2)
    return 0.5 * (var_ratio + t1 - 1 - var_ratio.log())

@register_kl(Independent, Independent)
def _kl_independent_independent(p, q):
    result = kl_divergence(p.base_dist, q.base_dist)
    return _sum_rightmost(result, p.reinterpreted_batch_ndims)
```


$\frac{1}{2}(\sigma^2/s^2 + (\mu - m)^2/s^2 - 1 - \log \sigma^2/s^2) = \log(s/\sigma) + (\sigma^2 + (\mu - m)^2)/(2s^2) - \frac{1}{2}$. Independent: the one-dimensional KLs, summed over coordinates.

> Expand the Normal line: it is our formula. Independent sums the per-coordinate KLs: the diagonal-Gaussian rule.

PyTorch source: torch/distributions/kl.py

▶ Notebook 2.7

Use the formula: which Gaussian is closer to p?




$$p = \mathcal{N}(1, 4^2)$$

	substitute	KL
$q_1 = \mathcal{N}(0, 2^2)$	$\log 2 + \frac{4+1}{32} - \frac{1}{2}$	
$q_2 = \mathcal{N}(1, 3^2)$	$\log \frac{4}{3} + \frac{9+0}{32} - \frac{1}{2}$	

The PML vote: which is closer to p, q_1 or q_2? Substitute into the formula.

▶ Notebook 2.7

Use the formula: which Gaussian is closer to p?




$$p = \mathcal{N}(1, 4^2)$$


	substitute	KL
$q_1 = \mathcal{N}(0, 2^2)$	$\log 2 + \frac{4+1}{32} - \frac{1}{2}$	0.35
$q_2 = \mathcal{N}(1, 3^2)$	$\log \frac{4}{3} + \frac{9+0}{32} - \frac{1}{2}$	0.07

KL(q_1‖p) = 0.35 nats and KL(q_2‖p) = 0.07. The formula settles the vote.

▶ Notebook 2.7

KL is an area you can watch





Purple area adds and red area subtracts; the net area is the KL. Drag q, or flip the direction.

▶ Notebook 2.8

KL is an average: an expectation under p

$$\mathrm{KL}(p \,\|\, q) = \sum_x p(x) \log \frac{p(x)}{q(x)}$$

Look again at the formula: each log ratio is weighted by p(x), and the weights add up to 1.

▶ Notebook 2.8

KL is an average: an expectation under p

$$\mathrm{KL}(p \,\|\, q) = \sum_x p(x) \log \frac{p(x)}{q(x)}$$

$$= \mathbb{E}_{x \sim p}\left[\log \frac{p(x)}{q(x)} \right]$$

E with x ~ p means: the average value when x is drawn from p.

So KL is an expectation: the average log ratio when x is drawn from p.

▶ Notebook 2.8

Two ways to compute an expectation

$$\mathbb{E}_{x\sim p}[f(x)] = \sum_x p(x)\, f(x)$$

For a discrete x, an expectation is a weighted sum: each value f(x) times its probability.

▶ Notebook 2.8

Two ways to compute an expectation

$$\mathbb{E}_{x\sim p}[f(x)] = \sum_{x} p(x)\, f(x)$$

$$\mathbb{E}_{\theta\sim p}[f(\theta)] = \int p(\theta)\, f(\theta)\, d\theta$$

For a continuous θ, the sum becomes an integral.

▶ Notebook 2.8

Two ways to compute an expectation

$$\mathbb{E}_{x\sim p}[f(x)] = \sum_{x} p(x)\, f(x)$$

$$\mathbb{E}_{\theta\sim p}[f(\theta)] = \int p(\theta)\, f(\theta)\, d\theta$$


$$\mathbb{E}_{x\sim p}[f(x)] \approx \frac{1}{S}\sum_{s=1}^{S} f(x_s), \qquad x_s \sim p$$


Draws land on each x in proportion to p(x), so a plain average over draws weights each f(x) by p(x) automatically.

> Or draw samples from p and take a plain average. Values that are more probable show up more often, so the weighting happens by itself.

▶ Notebook 2.8

Sample days and watch the average find the KL





Draw days from our weather p and average $\log_2$ p/q: +1 bit on sunny days, 0 on cloudy, −1 on rainy. The average settles on the KL, 0.25 bits, with no formula needed.

▶ Notebook 2.8

No formula needed: estimate KL by sampling

$$\mathrm{KL}(q\|p) = \mathbb{E}_{\theta\sim q}\big[\log q(\theta) - \log p(\theta)\big] \approx \frac{1}{S}\sum_{i=1}^{S}\big[\log q(\theta_i) - \log p(\theta_i)\big]$$

```
theta = q.sample((S,))
log_q = q.log_prob(theta)
log_p = p.log_prob(theta)
est = (log_q - log_p).mean()
exact = kl_divergence(q, p)
```

> Draw θ from q, score each draw with both log densities, and average. No formula needed.

▶ Notebook 2.8

No formula needed: estimate KL by sampling

$$\mathrm{KL}(q\|p) = \mathbb{E}_{\theta\sim q}\big[\log q(\theta) - \log p(\theta)\big] \approx \frac{1}{S}\sum_{i=1}^{S}\big[\log q(\theta_i) - \log p(\theta_i)\big]$$


```
theta = q.sample((S,))
log_q = q.log_prob(theta)
log_p = p.log_prob(theta)
est = (log_q - log_p).mean()
exact = kl_divergence(q, p)
```


	KL(q_2‖p)
exact formula	0.069
estimate, S = 10,000	0.067

Checked on a pair that has a formula: 0.067 from 10,000 draws, against the exact 0.069.

▶ Notebook 2.8

Watch the estimate settle






Add draws from q and watch the running average of log q − log p settle on the exact 0.349.


▶ Notebook 3.1

Six candidates: which q is closest to p?


$q = N(0, 2^2)$



$q = N(1, 3^2)$



$q = N(4, 1^2)$



$q = N(-3, 4^2)$



$q = N(1, 8^2)$



$q = N(2, 5^2)$




Six Gaussians against the same target $p = N(1, 4^2)$. Vote: which is closest?


▶ Notebook 3.1

Six candidates: which q is closest to p?


q = N(0, 2²) KL 0.35



q = N(1, 3²) **KL 0.07**



q = N(4, 1²) KL 1.20



q = N(−3, 4²) KL 0.50



q = N(1, 8²) KL 0.81




q = N(2, 5²) KL 0.09




Smallest KL: N(1, 3²) at 0.07, then N(2, 5²) at 0.09. Could another q do better?

▶ Notebook 3.1

Search a grid of means and widths: hover over it







Every point of the grid is one q = N(μ, σ²), coloured by its KL to p. Hover over it: the right panel shows that q against p, and its score.

▶ Notebook 3.1

Search a grid of means and widths: hover over it






The lightest point is q = p: 2,040 evaluations. For our network's fourteen numbers a grid needs $40^{14} \approx 3 \times 10^{22}$. We need to walk downhill instead.

▶ Notebook 3.2

Instead of a grid, walk downhill




$$\psi^* = \arg\min_{\psi} \mathrm{KL}(q_\psi \,\|\, p)$$

$$\psi \leftarrow \psi - \eta \, \nabla_\psi \mathrm{KL}(q_\psi \,\|\, p)$$

Cast the search as optimisation: follow the gradient of KL with respect to q's parameters.

▶ Notebook 3.2

Instead of a grid, walk downhill



$$\psi^* = \arg\min_{\psi} \mathrm{KL}(q_\psi \,\|\, p)$$

$$\psi \leftarrow \psi - \eta \, \nabla_\psi \mathrm{KL}(q_\psi \,\|\, p)$$

Gradient descent from N(4, 0.5²): the teal path. About a hundred steps, and the same recipe works for millions of parameters.

From N(4, 0.5²) the optimiser walks downhill to p. It needs only gradients, which autograd provides.

▶ Notebook 3.2

The same search in code

```
mu  = Parameter(tensor(4.0))
rho = Parameter(tensor(log(0.5)))
opt = Adam([mu, rho], lr=0.05)
for step in range(200):
    q = Normal(mu, rho.exp())
    loss = kl_divergence(q, p)
    opt.zero_grad()
    loss.backward()
    opt.step()
```

$$\sigma = \exp(\rho) > 0 \text{ for every } \rho$$

Store ρ = log σ, so that σ = exp(ρ) stays positive whatever an update does.

▶ Notebook 3.2

The same search in code

```
mu  = Parameter(tensor(4.0))
rho = Parameter(tensor(log(0.5)))
opt = Adam([mu, rho], lr=0.05)
for step in range(200):
    q = Normal(mu, rho.exp())
    loss = kl_divergence(q, p)
    opt.zero_grad()
    loss.backward()
    opt.step()
```


$$\sigma = \exp(\rho) > 0 \ \text{ for every } \rho$$


Rebuild q, score it, differentiate, update. The notebook runs exactly this loop and plots the loss, μ, σ and the learned curve.

> Rebuild q, score it, differentiate, update. The same five lines fit our network in Part 5.

▶ Notebook 3.2

Press play: the loss falls as q slides onto p







Start q far away, at N(4, 0.5²). Predict: which reaches the target first, the centre or the width?

▶ Notebook 3.2

Press play: the loss falls as q slides onto p






> The width gets there first and overshoots a little; the centre slides in by step 100. Then the loss, KL(q‖p), is about 0. Drag the slider to replay any step.

▶ Notebook 3.3

In two dimensions, q needs a covariance matrix: three families

isotropic: $s^2 I$



$$\Sigma = \begin{bmatrix} 1.00 & 0.00 \\ 0.00 & 1.00 \end{bmatrix}$$


1 number, whatever the dimension

Isotropic: one width for every direction, so the contours are circles. $\Sigma = s^2 I$ is a single number.

▶ Notebook 3.3

In two dimensions, q needs a covariance matrix: three families


isotropic: $s^2 I$



$$\Sigma = \begin{bmatrix} 1.00 & 0.00 \\ 0.00 & 1.00 \end{bmatrix}$$

1 number, whatever the dimension

diagonal: diag(s_1^2, s_2^2)



$$\Sigma = \begin{bmatrix} 1.70 & 0.00 \\ 0.00 & 0.36 \end{bmatrix}$$


2 numbers; d in d dimensions

> Diagonal: its own width per coordinate but no correlation, so the ellipses are axis-aligned. This is mean-field: $q(\theta) = q(\theta_1)\, q(\theta_2)$.

▶ Notebook 3.3

In two dimensions, q needs a covariance matrix: three families


isotropic: $s^2 I$



$$\Sigma = \begin{bmatrix} 1.00 & 0.00 \\ 0.00 & 1.00 \end{bmatrix}$$

1 number, whatever the dimension


diagonal: $\mathrm{diag}(s_1^2, s_2^2)$



$$\Sigma = \begin{bmatrix} 1.70 & 0.00 \\ 0.00 & 0.36 \end{bmatrix}$$

2 numbers; d in d dimensions

full: LL^T



$$\Sigma = \begin{bmatrix} 1.50 & 0.95 \\ 0.95 & 0.90 \end{bmatrix}$$

3 numbers; d(d + 1)/2 in general

Our network has d = 7 parameters: 1, 7 or 28 numbers. A network with a million weights: 1, a million, or 5×10^{11}.


> Full: the ellipse can tilt. Writing $\Sigma = LL^T$ with L lower-triangular keeps Σ a valid covariance for any L. It costs d(d + 1)/2 numbers.

▶ Notebook 3.3

Fit all three to the same tilted target

isotropic KL 7.01


$s^2 I$ · 1 number



$$\Sigma_q = \begin{bmatrix} 0.80 & 0.00 \\ 0.00 & 0.80 \end{bmatrix}$$

diagonal KL 7.05


$\mathrm{diag}(s_1^2, s_2^2)$ · 2 numbers



$$\Sigma_q = \begin{bmatrix} 0.78 & 0.00 \\ 0.00 & 0.82 \end{bmatrix}$$

full KL 14

LL^T · 3 numbers



$$\Sigma_q = \begin{bmatrix} 0.78 & -1.23 \\ -1.23 & 2.74 \end{bmatrix}$$


The target p (teal) has widths 1.5 and 0.75 and correlation 0.8. All three q start from the same random Gaussian, tilted the wrong way. Predict where each ends.

▶ Notebook 3.3

Fit all three to the same tilted target

isotropic KL 0.73


$s^2 I$ · 1 number



$$\Sigma_q = \begin{bmatrix} 0.32 & 0.00 \\ 0.00 & 0.32 \end{bmatrix}$$

diagonal KL 0.51


$\mathrm{diag}(s_1^2, s_2^2)$ · 2 numbers



$$\Sigma_q = \begin{bmatrix} 0.81 & 0.00 \\ 0.00 & 0.20 \end{bmatrix}$$

full KL ≈ 0

LL^T · 3 numbers



$$\Sigma_q = \begin{bmatrix} 2.25 & 0.90 \\ 0.90 & 0.56 \end{bmatrix}$$

Press play. The full q matches p exactly: KL → 0. The diagonal q cannot tilt; it shrinks to widths 0.90 and 0.45. The isotropic q shrinks to a circle of radius 0.57.

▶ Notebook 3.3

A q that cannot tilt shrinks instead

q family	widths it finds	KL(q‖p)
target p	1.50 and 0.75, tilted	—
isotropic	0.57 and 0.57	0.73
diagonal (mean-field)	0.90 and 0.45	0.51
full	1.50 and 0.75, tilted	0

> The full q is exact. The diagonal q is narrower than p in both coordinates; the isotropic q is narrower still in the wide direction.

▶ Notebook 3.3

A q that cannot tilt shrinks instead

q family	widths it finds	KL(q‖p)
target p	1.50 and 0.75, tilted	–
isotropic	0.57 and 0.57	0.73
diagonal (mean-field)	0.90 and 0.45	0.51
full	1.50 and 0.75, tilted	0

KL(q‖p) charges heavily for q-mass where p is small, off the tilted ridge. So a q that cannot tilt shrinks: it underestimates the uncertainty. Our network's q is diagonal; Part 5 shows what that does to its error bars.

> Rather than spill mass off the ridge, a q that cannot tilt shrinks. Mean-field VI underestimates uncertainty.

▶ Notebook 3.4

A mixture of Gaussians: first pick a bump, then draw from it

step 1: pick a bump

bump 1
probability ½

bump 2
probability ½

z


z ~ Categorical(½, ½)

step 2: draw θ from bump z

bump 1: N(−0.2, 0.4²)
bump 2: N(1.0, 0.5²)

θ

θ | z ~ N(μ_z, σ_z^2)



We see θ. We never see which bump made it.

> A mixture of Gaussians makes θ in two steps. First pick a bump, like tossing a fair coin: z = 1 or 2. Then draw θ from that bump's Gaussian.

▶ Notebook 3.4

A mixture of Gaussians: first pick a bump, then draw from it

step 1: pick a bump

bump 1
probability ½

bump 2
probability ½

z


z ~ Categorical(½, ½)

step 2: draw θ from bump z

bump 1: N(−0.2, 0.4²)
bump 2: N(1.0, 0.5²)

θ

θ | z ~ N(μ_z, σ_z^2)



We see θ. We never see which bump made it.

> Each bump contributes its own bell, scaled by how often it is picked: ½ N(−0.2, 0.4²) and ½ N(1, 0.5²).

▶ Notebook 3.4

A mixture of Gaussians: first pick a bump, then draw from it

step 1: pick a bump

bump 1
probability ½

bump 2
probability ½

z

z ~ Categorical(½, ½)


step 2: draw θ from bump z

bump 1: N(−0.2, 0.4²)
bump 2: N(1.0, 0.5²)

θ

θ | z ~ N(μ_z, σ_z^2)

We see θ. We never see which bump made it.



$$p(\theta) = \sum_k \pi_k \mathcal{N}(\theta; \mu_k, \sigma_k^2)$$

We see θ but not the bump that made it, so the density adds both bells: $p(\theta) = \Sigma_k \pi_k N(\theta; \mu_k, \sigma_k^2)$. Two bumps, one distribution.

▶ Notebook 3.4

Draw from the mixture yourself

step 1: pick a bump


bump 1: probability ½
N(−0.2, 0.4²) · picked 539×

bump 2: probability ½
N(1.0, 0.5²) · picked 461×

step 2: draw θ from that bump

latest: bump 1, θ = −0.30


1,000 draws so far



> Each draw first picks a bump (its colour), then draws θ from it. Add 10, 100, then 1,000 draws and watch the histogram fill in the teal density.

▶ Notebook 3.4

PyTorch has no formula for this KL. The definition still works.





```
kl_divergence(q, mixture)
```

How far is a Gaussian q from this mixture? How would you compute KL(q ‖ p)? Then ask PyTorch.

▶ Notebook 3.4

PyTorch has no formula for this KL. The definition still works.




```
kl_divergence(q, mixture)
```


```
NotImplementedError:
No KL(p || q) is implemented
for p type Normal and
q type MixtureSameFamily
```

No formula exists for a Gaussian against a mixture. But KL(q‖p) is an average under q, and averages can be estimated by sampling, as in 2.5.

▶ Notebook 3.4

Now optimise it: the code runs, and q does not move

```
theta = q.sample((5000,))
log_q = q.log_prob(theta)
log_p = p.log_prob(theta)
loss = (log_q - log_p).mean()
opt.zero_grad()
loss.backward(); opt.step()
```




Use the sampled KL as the loss and run the same loop on the mixture. Predict: does q move?

▶ Notebook 3.4

Now optimise it: the code runs, and q does not move


```
theta = q.sample((5000,))
log_q = q.log_prob(theta)
log_p = p.log_prob(theta)
loss = (log_q - log_p).mean()
opt.zero_grad()
loss.backward(); opt.step()
```



> No error, no warning, and q stays at N(6, 0.1²): after 400 steps the KL is still about 53. Is this a bug? Hold that question: Section 3.5 answers it.

▶ Notebook 3.5

Why q does not move: the gradient stops at a random node




> Boxes are deterministic nodes, computed from their inputs; circles are stochastic, drawn at random. With sample(), θ is stochastic, and the backward pass (red) cannot get from θ back to μ and σ.

Graph conventions after Lilian Weng, "From Autoencoder to Beta-VAE", which follows Kingma's NIPS 2015 talk

▶ Notebook 3.5

Why q does not move: the gradient stops at a random node




	$\partial \mathrm{KL}/\partial \mu$	$\partial \mathrm{KL}/\partial \sigma$
what autograd gets through sample()	0.0013	−0.0003
the true gradient	−0.063	−0.375

> Only the path through log q(θ)'s formula survives, and it averages to zero: 0.0013 and −0.0003 against −0.063 and −0.375. Not a bug: a missing path in the graph.

Graph conventions after Lilian Weng, "From Autoencoder to Beta-VAE", which follows Kingma's NIPS 2015 talk

▶ Notebook 3.5

The fix: move the randomness into ε, and θ = μ + σε




$$\epsilon \sim \mathcal{N}(0,1), \qquad \theta = \mu + \sigma\epsilon$$

> Draw the noise ε from a fixed N(0, 1), then shift and scale it. θ has the same distribution, N(μ, σ²), but it is now a deterministic node: a function of μ, σ and ε.

Graph conventions after Lilian Weng, "From Autoencoder to Beta-VAE"

▶ Notebook 3.5

The fix: move the randomness into ε, and θ = μ + σε




$$\epsilon \sim \mathcal{N}(0,1), \qquad \theta = \mu + \sigma\epsilon$$

Pause: with ε = −0.75, μ = 0 and σ = 2, find θ, ∂θ/∂μ and ∂θ/∂σ.

Graph conventions after Lilian Weng, "From Autoencoder to Beta-VAE"

▶ Notebook 3.5

The fix: move the randomness into ε, and θ = μ + σε



$$\epsilon \sim \mathcal{N}(0,1), \qquad \theta = \mu + \sigma\epsilon \qquad\qquad \theta = -1.5, \quad \frac{\partial\theta}{\partial\mu} = 1, \quad \frac{\partial\theta}{\partial\sigma} = \epsilon = -0.75$$


> θ = −1.5, ∂θ/∂μ = 1 and ∂θ/∂σ = ε = −0.75. The only random node, ε, has no parameters, so nothing blocks the path. In PyTorch this is rsample().

Graph conventions after Lilian Weng, "From Autoencoder to Beta-VAE"


▶ Notebook 3.5

What a fixed ε does when μ or σ changes

move μ from 0 to 1 θ: −1.50 → −0.50



widen σ from 2 to 3 θ: −1.50 → −2.25




Keep the same noise draw, ε = −0.75. Shift μ by 1: the draw moves right by exactly 1.


▶ Notebook 3.5

What a fixed ε does when μ or σ changes

move μ from 0 to 1 θ: −1.50 → −0.50



widen σ from 2 to 3 θ: −1.50 → −2.25




$$\frac{\partial \theta}{\partial \mu} = 1, \qquad \frac{\partial \theta}{\partial \sigma} = \epsilon = -0.75$$

> Widen σ by 1: the draw moves by ε = −0.75, further out on its side. These are the derivatives the gradient needs.

▶ Notebook 3.5

Side by side: where the randomness lives




deterministic node · stochastic node · forward: compute · backward: gradient · blocked

> With sample(), the random node sits between the parameters and the loss, and the gradient cannot cross it.

Graph conventions after Lilian Weng, "From Autoencoder to Beta-VAE"

▶ Notebook 3.5

Side by side: where the randomness lives




With rsample(), the randomness enters as a parameter-free input ε and θ = μ + σε is deterministic: the gradient flows from the loss to μ and σ. Same distribution, different graph.

Graph conventions after Lilian Weng, "From Autoencoder to Beta-VAE"

▶ Notebook 3.5

Change one word: rsample() fits the mixture

```
# one word changed
theta = q.rsample((5000,))
log_q = q.log_prob(theta)
log_p = p.log_prob(theta)
loss = (log_q - log_p).mean()
opt.zero_grad()
loss.backward(); opt.step()
```




Same loop, same start, same seed, with q.rsample. Press play.

▶ Notebook 3.5

Change one word: rsample() fits the mixture

```
# one word changed
theta = q.rsample((5000,))
log_q = q.log_prob(theta)
log_p = p.log_prob(theta)
loss = (log_q - log_p).mean()
opt.zero_grad()
loss.backward(); opt.step()
```




q settles at N(0.43, 0.72²), KL = 0.056: the best single Gaussian for this mixture. No formula was needed.


▶ Notebook 3.6

The price: reverse KL picks one bump


KL(pllq) 6.43



KL(qllp), start left 3.92



KL(qllp), start right 4.42




Fit one Gaussian to two narrow bumps, three times. Predict where each q settles.


▶ Notebook 3.6

The price: reverse KL picks one bump


KL(p||q) 0.72



KL(q||p), start left 0.36



KL(q||p), start right 1.20



KL(p||q) spreads over both bumps. KL(q||p) locks onto whichever bump is nearer its start, and reports it confidently.

▶ Notebook 4.1

Back to Bayes: minimise KL to the posterior directly?

$$\mathrm{KL}\big(q_\psi(\theta) \,\|\, p(\theta \mid \mathcal{D})\big) = \mathbb{E}_{\theta\sim q}\big[\log q_\psi(\theta) - \log p(\theta \mid \mathcal{D})\big]$$

> The plan from Part 1: make q close to the posterior. Use KL(q ‖ posterior): an average under q, which we can sample.

A careful written derivation: Gregory Gundersen, "The ELBO in Variational Inference"

▶ Notebook 4.1

Back to Bayes: minimise KL to the posterior directly?

$$\mathrm{KL}\big(q_\psi(\theta) \,\|\, p(\theta \mid \mathcal{D})\big) = \mathbb{E}_{\theta\sim q}\big[\log q_\psi(\theta) - \log p(\theta \mid \mathcal{D})\big]$$

the loss needs	available?
draws θ ~ q	yes: rsample, as in 3.5
$\log q_\psi(\theta)$	yes
$\log p(\theta \mid \mathcal{D})$	no: it needs p($\mathcal{D}$), the integral of 1.4

> Everything is available except log p(θ | $\mathcal{D}$), which needs the evidence p($\mathcal{D}$): the integral we cannot do. Four small steps get rid of it.

A careful written derivation: Gregory Gundersen, "The ELBO in Variational Inference"

▶ Notebook 4.1

Step 1: Bayes' rule, in logs

$$p(\theta \mid \mathcal{D}) = \frac{p(\mathcal{D} \mid \theta)\, p(\theta)}{p(\mathcal{D})}$$

Bayes' rule from Part 1: the posterior is likelihood times prior, divided by the evidence.

▶ Notebook 4.1

Step 1: Bayes' rule, in logs

$$p(\theta \mid \mathcal{D}) = \frac{p(\mathcal{D} \mid \theta)\, p(\theta)}{p(\mathcal{D})}$$

$$\log p(\theta \mid \mathcal{D}) = \log p(\mathcal{D} \mid \theta) + \log p(\theta) - \log p(\mathcal{D})$$

Rule: log(a · b / c) = log a + log b − log c.

> Take logs. A product becomes a sum, and a division a subtraction.

▶ Notebook 4.1

Step 1: Bayes' rule, in logs

$$p(\theta \mid \mathcal{D}) = \frac{p(\mathcal{D} \mid \theta)\, p(\theta)}{p(\mathcal{D})}$$

$$\log p(\theta \mid \mathcal{D}) = \log p(\mathcal{D} \mid \theta) + \log p(\theta) - \log p(\mathcal{D})$$

Rule: log(a · b / c) = log a + log b − log c.

Look at the last term: $\log p(\mathcal{D})$ has no θ in it. Whatever θ is, it is the same number, fixed by our model and our data.

> The last term does not contain θ: it is one fixed number. Remember that; it does the work in step 3.

▶ Notebook 4.1

Step 2: substitute into the KL

$$\mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big) = \mathbb{E}_q\big[\log q(\theta) - \log p(\theta \mid \mathcal{D})\big]$$

The KL we want: an average, under q, of log q minus log posterior.

▶ Notebook 4.1

Step 2: substitute into the KL

$$\mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big) = \mathbb{E}_q\big[\log q(\theta) - \log p(\theta \mid \mathcal{D})\big]$$

$$= \mathbb{E}_q\Big[\log q(\theta) - \big(\log p(\mathcal{D} \mid \theta) + \log p(\theta) - \log p(\mathcal{D})\big)\Big]$$

Replace the log posterior by its three terms from step 1, in brackets.

▶ Notebook 4.1

Step 2: substitute into the KL

$$\mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big) = \mathbb{E}_q\big[\log q(\theta) - \log p(\theta \mid \mathcal{D})\big]$$

$$= \mathbb{E}_q\Big[\log q(\theta) - \big(\log p(\mathcal{D} \mid \theta) + \log p(\theta) - \log p(\mathcal{D})\big)\Big]$$

$$= \mathbb{E}_q\big[\log q(\theta) - \log p(\mathcal{D} \mid \theta) - \log p(\theta) + \log p(\mathcal{D})\big]$$

$\mathrm{E}_q[\,\cdot\,]$ means: draw θ from q, evaluate the bracket, average. Rule used: −(a + b − c) = −a − b + c.

Open the brackets: every sign flips, and log p($\mathcal{D}$) now enters with a plus sign.

▶ Notebook 4.1

Step 3: a constant comes out of an average

$$\mathbb{E}_q\big[f(\theta) + c\big] = \mathbb{E}_q\big[f(\theta)\big] + c \qquad \text{when } c \text{ does not depend on } \theta$$

> A rule about averages: add the same number c to every value, and the average goes up by exactly c.

▶ Notebook 4.1

Step 3: a constant comes out of an average

$$\mathbb{E}_q\big[f(\theta) + c\big] = \mathbb{E}_q\big[f(\theta)\big] + c \qquad \text{when } c \text{ does not depend on } \theta$$

	three draws	their average
$f(\theta)$	2, 4, 6	4
$f(\theta) + 10$	12, 14, 16	14 = 4 + 10

> Check it: the average of 2, 4 and 6 is 4. Add 10 to each and the average is 14.

▶ Notebook 4.1

Step 3: a constant comes out of an average

$$\mathbb{E}_q\big[f(\theta) + c\big] = \mathbb{E}_q\big[f(\theta)\big] + c \qquad \text{when } c \text{ does not depend on } \theta$$

	three draws	their average
$f(\theta)$	2, 4, 6	4
$f(\theta) + 10$	12, 14, 16	14 = 4 + 10

$$\mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big) = \mathbb{E}_q\big[\log q(\theta) - \log p(\mathcal{D} \mid \theta) - \log p(\theta)\big] + \log p(\mathcal{D})$$

> log p($\mathcal{D}$) is such a constant: step 1 showed it has no θ. So it leaves the average.

▶ Notebook 4.1

Step 4: rearrange, and name the part we can compute

$$\mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big) = -\mathbb{E}_q\big[\log p(\mathcal{D} \mid \theta) + \log p(\theta) - \log q(\theta)\big] + \log p(\mathcal{D})$$

> Take a minus sign outside the average: inside, the likelihood and the prior now have plus signs, and log q a minus.

▶ Notebook 4.1

Step 4: rearrange, and name the part we can compute

$$\mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big) = -\mathbb{E}_q\big[\log p(\mathcal{D} \mid \theta) + \log p(\theta) - \log q(\theta)\big] + \log p(\mathcal{D})$$

$$\log p(\mathcal{D}) = \mathbb{E}_q\big[\log p(\mathcal{D} \mid \theta) + \log p(\theta) - \log q(\theta)\big] + \mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big)$$

Move that average to the other side: log p($\mathcal{D}$) = average + KL.

▶ Notebook 4.1

Step 4: rearrange, and name the part we can compute

$$\mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big) = -\mathbb{E}_q\big[\log p(\mathcal{D} \mid \theta) + \log p(\theta) - \log q(\theta)\big] + \log p(\mathcal{D})$$

$$\log p(\mathcal{D}) = \mathbb{E}_q\big[\log p(\mathcal{D} \mid \theta) + \log p(\theta) - \log q(\theta)\big] + \mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big)$$

$$\log p(\mathcal{D}) = \underbrace{\mathbb{E}_q\big[\log p(\mathcal{D} \mid \theta) + \log p(\theta) - \log q(\theta)\big]}_{\mathrm{ELBO}(q)} + \mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big)$$

> Give the average a name: the ELBO, the evidence lower bound. Every term in it is computable: draws from q, the likelihood, the prior and log q. No p($\mathcal{D}$) anywhere.

▶ Notebook 4.1

One fixed number = what we can compute + what we cannot

$$\underbrace{\log p(\mathcal{D})}_{\text{fixed}} = \underbrace{\mathrm{ELBO}(q)}_{\text{computable}} + \underbrace{\mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big)}_{\text{unknown, but } \geq 0}$$

> Read the identity as a budget: one fixed number, split into a part we can compute and a part we cannot.

▶ Notebook 4.1

One fixed number = what we can compute + what we cannot

$$\underbrace{\log p(\mathcal{D})}_{\text{fixed}} = \underbrace{\mathrm{ELBO}(q)}_{\text{computable}} + \underbrace{\mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big)}_{\text{unknown, but } \geq 0}$$


piece	what it is	changes with q?
$\log p(\mathcal{D})$	the log evidence: the data's probability, averaged over every θ	no: fixed
$\mathrm{ELBO}(q)$	average under q of log likelihood + log prior − log q	yes; computable
the KL	how far q is from the posterior	yes; ≥ 0, and 0 only at the posterior

> The KL is the part we want small. We cannot compute it, but we know one thing about it: it is never negative.

▶ Notebook 4.1

Why KL is never negative

$$\log t \leq t - 1 \quad \text{for every } t > 0$$




A fact about the logarithm: its curve never rises above its tangent line at t = 1, so log t ≤ t − 1, with equality only at t = 1.

▶ Notebook 4.1

Why KL is never negative

$$\log t \leq t - 1 \quad \text{for every } t > 0$$

$$-\mathrm{KL}(q\|p) = \mathbb{E}_q\left[\log \frac{p(\theta)}{q(\theta)}\right] \leq \mathbb{E}_q\left[\frac{p(\theta)}{q(\theta)} - 1\right]$$



Apply it to the ratio t = p(θ)/q(θ), inside the average under q. Minus the KL is the average of log(p/q).


▶ Notebook 4.1

Why KL is never negative

$$\log t \le t - 1 \quad \text{for every } t > 0$$

$$-\mathrm{KL}(q\|p) = \mathbb{E}_q\left[\log \frac{p(\theta)}{q(\theta)}\right] \le \mathbb{E}_q\left[\frac{p(\theta)}{q(\theta)} - 1\right]$$

$$\mathbb{E}_q\left[\frac{p}{q}\right] = \int q(\theta)\,\frac{p(\theta)}{q(\theta)}\,d\theta = \int p(\theta)\,d\theta = 1$$



So $-\mathrm{KL} \le 1 - 1 = 0$: KL ≥ 0, and KL = 0 only when q = p.

> The average of p/q under q is the integral of p, which is 1. So minus the KL is at most 1 − 1 = 0.

▶ Notebook 4.1

The ELBO is a lower bound on the log evidence

In plain words: whatever q we choose, the **ELBO** never rises above the **log evidence log p($\mathcal{D}$)**. It is a lower bound **on log p($\mathcal{D}$)**: an **E**vidence **L**ower **BO**und.

$$\mathrm{ELBO}(q) = \log p(\mathcal{D}) - \underbrace{\mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big)}_{\geq 0} \leq \log p(\mathcal{D})$$


> Lower bound on what? On the log evidence, log p($\mathcal{D}$). The red KL is never negative, so subtracting it can only lower the number: the purple ELBO always sits below the blue log p($\mathcal{D}$).

▶ Notebook 4.1

The ELBO is a lower bound on the log evidence

In plain words: whatever q we choose, the **ELBO** never rises above the **log evidence log p($\mathcal{D}$)**. It is a lower bound **on log p($\mathcal{D}$)**: an **E**vidence **L**ower **BO**und.

$$\mathrm{ELBO}(q) = \log p(\mathcal{D}) - \underbrace{\mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big)}_{\geq 0} \leq \log p(\mathcal{D})$$



A coin with q = Beta(10, 10): the ELBO sits 0.62 nats below log p($\mathcal{D}$), and the gap is exactly the KL to the posterior.

▶ Notebook 4.1

The ELBO is a lower bound on the log evidence

In plain words: whatever q we choose, the **ELBO** never rises above the **log evidence log p($\mathcal{D}$)**. It is a lower bound **on log p($\mathcal{D}$)**: an **E**vidence **L**ower **BO**und.


$$\mathrm{ELBO}(q) = \log p(\mathcal{D}) - \underbrace{\mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big)}_{\geq 0} \leq \log p(\mathcal{D})$$

	log p($\mathcal{D}$)	it is the log of …
the coin: discrete data	−21.40	a probability: never above 0
our network: continuous data	−1.43	a density: either sign

> "Lower bound" says nothing about signs: a probability is at most 1, but a density can exceed it (log p($\mathcal{D}$ | θ_A) = +12.2). Only the order is guaranteed.

▶ Notebook 4.1


Raise the ELBO, and the KL falls by exactly as much



Three q's for the same coin. log p(𝒟) does not move. As q improves, the ELBO climbs and the gap, the KL, shrinks by exactly the same amount.

▶ Notebook 4.1

Raise the ELBO, and the KL falls by exactly as much



$$\max_q \ \mathrm{ELBO}(q) \quad \Longleftrightarrow \quad \min_q \ \mathrm{KL}\big(q \,\|\, p(\theta \mid \mathcal{D})\big)$$

> So maximising the ELBO, which we can compute, minimises the KL to the posterior, which we cannot. At the best q the gap is 0.

▶ Notebook 4.1

The ELBO in two terms: fit the data, stay near the prior

$$\mathrm{ELBO}(q) = \mathbb{E}_q\big[\log p(\mathcal{D} \mid \theta) + \log p(\theta) - \log q(\theta)\big]$$

Take the ELBO apart.

▶ Notebook 4.1

The ELBO in two terms: fit the data, stay near the prior

$$\mathrm{ELBO}(q) = \mathbb{E}_q\big[\log p(\mathcal{D} \mid \theta) + \log p(\theta) - \log q(\theta)\big]$$

$$= \mathbb{E}_q\big[\log p(\mathcal{D} \mid \theta)\big] + \mathbb{E}_q\big[\log p(\theta) - \log q(\theta)\big]$$

> The average of a sum is the sum of the averages: a likelihood term and the rest.

▶ Notebook 4.1


The ELBO in two terms: fit the data, stay near the prior

$$\begin{aligned}\mathrm{ELBO}(q) &= \mathbb{E}_q\big[\log p(\mathcal{D} \mid \theta) + \log p(\theta) - \log q(\theta)\big] \\ &= \mathbb{E}_q\big[\log p(\mathcal{D} \mid \theta)\big] + \mathbb{E}_q\big[\log p(\theta) - \log q(\theta)\big] \\ &= \underbrace{\mathbb{E}_q\big[\log p(\mathcal{D} \mid \theta)\big]}_{\text{fit the data}} - \underbrace{\mathrm{KL}\big(q(\theta) \,\|\, p(\theta)\big)}_{\text{stay near the prior}}\end{aligned}$$


> The rest is minus a KL, from q to the prior. Fit the data, but stay near the prior. For our network the second term is the diagonal-Gaussian formula of 2.4; the first needs samples.

▶ Notebook 4.2

Check the identity where p($\mathcal{D}$) is known: a coin



One fixed total



−22.020 + 0.623 = −21.397

A coin with a Beta(5, 3) prior and 17 heads in 30 tosses has an exact posterior and an exact p($\mathcal{D}$). Drag q: the ELBO and the KL trade off, and their sum never changes.

▶ Notebook 5.1

The ELBO for our network, in code


```
mu  = Parameter(0.5 * randn(7))          # 7 means
rho = Parameter(full((7,), log(0.1)))    # 7 log-widths
q = Normal(mu, rho.exp())                # one bell per parameter
theta = q.rsample((16,))                 # 16 sampled networks
log_lik = log_likelihood(theta).mean()   # forward pass
kl = kl_divergence(q, prior).sum()       # closed form
loss = -(log_lik - kl)                   # minus the ELBO
```

> Fourteen parameters, sixteen sampled networks per step, and the closed-form KL to the prior.

▶ Notebook 5.1

The ELBO for our network, in code



```
mu  = Parameter(0.5 * randn(7))          # 7 means
rho = Parameter(full((7,), log(0.1)))    # 7 log-widths
q = Normal(mu, rho.exp())                # one bell per parameter
theta = q.rsample((16,))                 # 16 sampled networks
log_lik = log_likelihood(theta).mean()   # forward pass
kl = kl_divergence(q, prior).sum()       # closed form
loss = -(log_lik - kl)                   # minus the ELBO
```


Adam, learning rate 0.02, 4,000 steps; best of five random starts. No p($\mathcal{D}$) anywhere.

> Every line is computable: rsample, a forward pass and a formula. Never p($\mathcal{D}$).

▶ Notebook 5.1

Press play: q learns the network







Start from random means and narrow widths. Watch q's predictive band and the ELBO.

▶ Notebook 5.1

Press play: q learns the network






The ELBO climbs from −1,235 to about −7.2 and the band settles onto the data. It stays below log p($\mathcal{D}$) ≈ −1.43.

▶ Notebook 5.1


What q learned: one hidden unit switched off



The fitted bell for each parameter, against the N(0, 1) prior in grey.

▶ Notebook 5.1

What q learned: one hidden unit switched off




$v_1 \approx 0.01$ with width 0.035: the first hidden unit is switched off, and its w_1 and b_1 fall back to the prior.

> VI switched the first hidden unit off. Mean-field VI often prunes units like this: every parameter kept away from the prior costs KL.

▶ Notebook 5.1

Mean-field VI, and full-covariance VI

diagonal q (mean-field)



KL 0.51

full-covariance q



KL ≈ 0

$$\text{mean-field: } q(\theta) = \prod_{j=1}^{7} \mathcal{N}(\theta_j; \mu_j, \sigma_j^2)$$


One independent bell per parameter: the diagonal family of 3.3. 7 + 7 = 14 numbers.

> Our q is mean-field: q factorises into one bell per parameter, so it cannot express correlations. In 3.3 that made a diagonal q shrink.

▶ Notebook 5.1

Mean-field VI, and full-covariance VI

diagonal q (mean-field)



KL 0.51

full-covariance q



KL ≈ 0

$$\text{mean-field: } q(\theta) = \prod_{j=1}^{7} \mathcal{N}(\theta_j; \mu_j, \sigma_j^2)$$

One independent bell per parameter: the diagonal family of 3.3. 7 + 7 = 14 numbers.

$$\text{full covariance: } q(\theta) = \mathcal{N}\left(\mu,\, LL^\top\right)$$


One tilted bell in $\mathbb{R}^7$: 7 + 28 = 35 numbers; d(d + 1)/2 in general.

> Full-covariance VI uses one tilted Gaussian over all seven parameters: Σ = LLᵀ, 35 numbers instead of 14. Fine for 7 parameters; hopeless for a million.

▶ Notebook 5.1

Mean-field VI, and full-covariance VI

diagonal q (mean-field)



KL 0.51

full-covariance q



KL ≈ 0

$$\text{mean-field: } q(\theta) = \prod_{j=1}^{7} \mathcal{N}(\theta_j; \mu_j, \sigma_j^2)$$

One independent bell per parameter: the diagonal family of 3.3. 7 + 7 = 14 numbers.

$$\text{full covariance: } q(\theta) = \mathcal{N}\left(\mu,\, LL^\top\right)$$


One tilted bell in $\mathbb{R}^7$: 7 + 28 = 35 numbers; d(d + 1)/2 in general.

Full q: correlation −0.76 between w_2 and v_2; ELBO −7.18 → −6.32.

> Starting from the mean-field answer, it learns that a steeper w_2 goes with a smaller v_2, and the ELBO rises by 0.86 nats: the KL to the posterior falls by exactly that much.

▶ Notebook 5.1

Against the reference: the right curve, bands still too narrow




	ELBO	sd at −2.5	sd at 2.5
mean-field	−7.18	0.110	0.110
MCMC reference	$\log p(\mathcal{D}) \approx -1.43$	0.215	0.168

Mean-field VI gets the mean curve right, but its bands are too narrow beyond the data: the diagonal q of 3.3 shrinking, in seven dimensions.

▶ Notebook 5.1

Against the reference: the right curve, bands still too narrow



	ELBO	sd at −2.5	sd at 2.5
mean-field	−7.18	0.110	0.110
full covariance	−6.32	0.117	0.120
MCMC reference	$\log p(\mathcal{D}) \approx -1.43$	0.215	0.168

Full covariance helps a little. The rest of the gap is the family: the posterior has several modes (swap the hidden units, flip signs), and one Gaussian sits on one.

▶ Notebook 5.2

Six obstacles, six tools

obstacle	tool	where
$p(\mathcal{D})$ is an impossible integral	approximate the posterior with q	1.4 – 1.5
"close" needs a number	KL(q ‖ p)	2.1 – 2.4
a grid cannot search 14 numbers	gradient descent	3.1 – 3.2
no formula for the average	Monte Carlo	2.5, 3.4
no gradient through a draw	$\theta = \mu + \sigma\varepsilon$, rsample()	3.5
KL to the posterior needs $p(\mathcal{D})$	maximise the ELBO	4.1

> Every obstacle today earned a tool, and our network used them all. In Lecture 3 a VAE does the same with one q per image.

▶ Notebook 5.2

Generative models create this inference problem again

A hidden description can generate an observation.

> In an image model, a hidden code z can generate an observed image x, just as the mixture's bump z generated θ in 3.4.

▶ Notebook 5.2

Generative models create this inference problem again


A hidden description can generate an observation.

Next: learn representations for images, then ask whether they can generate new ones.

> Lecture 2 builds autoencoders; Lecture 3 returns to VI with an encoder that outputs q for each image.

▶ Notebook

By hand: two runs of this course, two grade distributions



Appendix: KL by hand. A second route to the same formula.


Grades in last year's run (q) and this year's (p).

> A second route to KL, by hand: the grade distributions of two runs of this course. How different are they, as one number?

KL intuition after ritvikmath, "The KL Divergence: Data Science Basics" (YouTube)

▶ Notebook

By hand: two runs of this course, two grade distributions



Appendix: KL by hand. A second route to the same formula.

Grades in last year's run (q) and this year's (p).

A: 10% → 40%. B: 40% → 10%. C: 50% both years.

> Four times as many A's, a quarter as many B's, the same share of C's.

KL intuition after ritvikmath, "The KL Divergence: Data Science Basics" (YouTube)

Compare grade by grade: ratios

grade	p	q	p/q
A	0.4	0.1	4
B	0.1	0.4	¼
C	0.5	0.5	1



> How many times more common is each grade this year? A: 4. B: ¼. C: 1.

KL intuition after ritvikmath, "The KL Divergence: Data Science Basics" (YouTube)

Compare grade by grade: ratios

grade	p	q	p/q
A	0.4	0.1	4
B	0.1	0.4	¼
C	0.5	0.5	1



Average the ratios? (4 + ¼ + 1) / 3 = 1.75. Identical years would give 1, not 0, and ×4 sits 3 above 1 while ÷4 sits only 0.75 below.

> Averaging raw ratios is lopsided: a fourfold rise counts four times as much as a fourfold fall.

KL intuition after ritvikmath, "The KL Divergence: Data Science Basics" (YouTube)

What should a ratio count for? Find the function

We want a function f of the ratio r with:

$$f(1) = 0$$

$$f(4) = +y, \qquad f(\tfrac{1}{4}) = -y$$

$$f(2 \cdot 2) = f(2) + f(2)$$

$f(r)$	$f(\frac{1}{4})$	$f(1)$	$f(4)$	$f(2) + f(2)$

> What should a ratio count for? No change should count 0; a fourfold rise and a fourfold fall should cancel; and changes should add up.

KL intuition after ritvikmath, "The KL Divergence: Data Science Basics" (YouTube)

What should a ratio count for? Find the function

We want a function f of the ratio r with:

$$f(1) = 0$$

$$f(4) = +y, \qquad f(\tfrac{1}{4}) = -y$$

$$f(2 \cdot 2) = f(2) + f(2)$$

$f(r)$	$f(\frac{1}{4})$	$f(1)$	$f(4)$	$f(2) + f(2)$
$r - 1$	−0.75	0	3	2 ≠ 3 ✗
$r - \frac{1}{r}$	−3.75	0	3.75	3 ≠ 3.75 ✗

> Try a few functions. r − 1 is lopsided. r − 1/r balances ×4 against ÷4, but two doublings do not add up to a quadrupling.

KL intuition after ritvikmath, "The KL Divergence: Data Science Basics" (YouTube)

What should a ratio count for? Find the function

We want a function f of the ratio r with:

$$f(1) = 0$$

$$f(4) = +y, \qquad f(\tfrac{1}{4}) = -y$$

$$f(2 \cdot 2) = f(2) + f(2)$$


$f(r)$	$f(\frac{1}{4})$	$f(1)$	$f(4)$	$f(2) + f(2)$
$r - 1$	−0.75	0	3	2 ≠ 3 ✗
$r - \frac{1}{r}$	−3.75	0	3.75	3 ≠ 3.75 ✗
$\log r$	−1.39	0	1.39	1.39 = 1.39 ✓

> The logarithm passes every test: log 4 = 1.39, log ¼ = −1.39, and log 2 + log 2 = log 4.

KL intuition after ritvikmath, "The KL Divergence: Data Science Basics" (YouTube)

▶ Notebook

Weight each grade by this year's share: that is KL




$$\text{plain average: } \tfrac{1}{3}\left(1.39 - 1.39 + 0\right) = 0$$

> Why not simply average the logs? Here the plain average is exactly 0, yet the two years clearly differ.

KL intuition after ritvikmath, "The KL Divergence: Data Science Basics" (YouTube)

▶ Notebook

Weight each grade by this year's share: that is KL



$$\text{plain average: } \tfrac{1}{3}\left(1.39 - 1.39 + 0\right) = 0$$

$$\text{weighted: } 0.4(1.39) + 0.1(-1.39) + 0.5(0) = 0.42$$

> Weight each grade by how many students got it this year: the A's, now common, count most. The weighted sum is 0.42.

KL intuition after ritvikmath, "The KL Divergence: Data Science Basics" (YouTube)

▶ Notebook

Weight each grade by this year's share: that is KL

p(x) × log ratio, and their sum



$$\text{plain average: } \tfrac{1}{3}\left(1.39 - 1.39 + 0\right) = 0$$

$$\text{weighted: } 0.4(1.39) + 0.1(-1.39) + 0.5(0) = 0.42$$

$$\mathrm{KL}(p \,\|\, q) = \sum_x p(x) \log \frac{p(x)}{q(x)}$$

That weighted average of log ratios is the Kullback–Leibler divergence of p from q.

KL intuition after ritvikmath, "The KL Divergence: Data Science Basics" (YouTube)

▶ Notebook

The unit: nats, or bits

$$\mathrm{KL}(p\|q) = \sum_x p(x)\,\ln\frac{p(x)}{q(x)} = 0.42 \text{ nats}$$

$$\sum_x p(x)\,\log_2\frac{p(x)}{q(x)} = 0.60 \text{ bits}$$

The log's base sets the unit. Natural log: nats. Log base 2: bits. Here 0.42 nats is exactly 0.60 bits; 1 nat ≈ 1.44 bits.

▶ Notebook

The unit: nats, or bits

$$\mathrm{KL}(p\|q) = \sum_x p(x)\,\ln\frac{p(x)}{q(x)} = 0.42 \text{ nats}$$


$$\sum_x p(x)\,\log_2\frac{p(x)}{q(x)} = 0.60 \text{ bits}$$

Read 0.42 nats as $e^{0.42} \approx 1.5$: on average, a grade from this year is 1.5 times more probable under this year's distribution than under last year's.

> 0.42 nats means a typical this-year grade is about 1.5 times more probable under this year's distribution than under last year's.

▶ Notebook

Swap the reference year: a different number




$$\mathrm{KL}(\text{this} \,\|\, \text{last}) = 0.64$$

A different pair of years: last year 1% got an A, this year 25% did. Measured from this year: 0.64 nats.

▶ Notebook

Swap the reference year: a different number



$$\mathrm{KL}(\text{this} \,\|\, \text{last}) = 0.64$$


$$\mathrm{KL}(\text{last} \,\|\, \text{this}) = 0.30$$


Which distribution comes first matters: KL is not symmetric.

> Measured from last year instead: 0.30 nats. Changing the reference changes the answer: KL is not symmetric.

Notebook

Why the reference matters







> From this year's view, A is 25 times more common than before, and a quarter of the class has it: A contributes 0.80.

Notebook

Why the reference matters





The first distribution supplies the weights, so it decides which changes count.

> From last year's view A was rare, so its change barely counts (−0.03); the drop in B dominates instead.

▶ Notebook

Sample grades and watch the average find the KL

grades drawn from this year's p (latest 30)


C	A	A	C	B	A	C	B	C	C
C	B	C	C	B	A	A	C	C	C
A	A	B	B	A	C	C	A	A	B

A: log ratio +1.39 — 411 draws, 41%

B: log ratio −1.39 — 108 draws, 11%

C: log ratio +0.00 — 481 draws, 48%


average of 1,000 log ratios: 0.420



Draw grades from this year's distribution and average their log ratios. The counts approach 40%, 10%, 50%, and the average approaches the exact 0.42.

▶ Notebook

Sampling from the prior: a few lucky draws carry everything




$$p(\mathcal{D}) \approx \frac{1}{S}\sum_{s=1}^{S} p(\mathcal{D} \mid \theta_s)$$

$$\theta_s \sim p(\theta)$$

Average the likelihood over draws from the prior: four seeds, up to ten million draws each.

▶ Notebook

Sampling from the prior: a few lucky draws carry everything



$$p(\mathcal{D}) \approx \frac{1}{S}\sum_{s=1}^{S} p(\mathcal{D} \mid \theta_s)$$


$$\theta_s \sim p(\theta)$$

Ten draws: off by about 1,000 nats. Ten million: about 103 draws carry the estimate, −1.43.

> Almost every prior draw explains the data terribly. Seven parameters just about work; more parameters or more data make it hopeless.

▶ Notebook

A KL value is how fast evidence piles up




$$\mathrm{KL}\big(\mathrm{Bern}(0.8) \,\|\, \mathrm{Bern}(0.5)\big) \approx 0.19 \text{ nats}$$

> A coin lands heads 80% of the time. Each toss favours “0.8” over “fair” by 0.19 nats on average. After 30 tosses, by what factor?

▶ Notebook

A KL value is how fast evidence piles up



$$\mathrm{KL}\big(\mathrm{Bern}(0.8) \,\|\, \mathrm{Bern}(0.5)\big) \approx 0.19 \text{ nats}$$

$$\frac{p(\mathcal{D} \mid 0.8)}{p(\mathcal{D} \mid 0.5)} \approx e^{0.19\,n}$$

The reverse direction, KL(0.5‖0.8) = 0.22, is a different number.

> About e^(30 × 0.19) ≈ 325. KL is the rate at which evidence piles up: 6.9× after 10 tosses, 325× after 30.

▶ Notebook

A sensor has an unknown constant bias

$$\theta \sim \mathcal{N}(0, 1), \qquad y_i \mid \theta \sim \mathcal{N}(\theta, 0.5^2)$$

A sensor reads a hidden bias θ, with noise. Name the unknown, the data and a q.

▶ Notebook

A sensor has an unknown constant bias

$$\theta \sim \mathcal{N}(0, 1), \qquad y_i \mid \theta \sim \mathcal{N}(\theta, 0.5^2)$$

$$p(\mathcal{D}, \theta) = p(\theta) \prod_i p(y_i \mid \theta)$$

> The joint is the prior times the likelihood of each reading: available.

▶ Notebook

A sensor has an unknown constant bias

$$\theta \sim \mathcal{N}(0, 1), \qquad y_i \mid \theta \sim \mathcal{N}(\theta, 0.5^2)$$


$$p(\mathcal{D}, \theta) = p(\theta) \prod_i p(y_i \mid \theta)$$

$$\max_{\psi} \mathbb{E}_{q_\psi(\theta)}[\log p(\mathcal{D}, \theta) - \log q_\psi(\theta)]$$

Maximise the ELBO over q's parameters. The same tools apply.

▶ Notebook


More samples, tighter estimates



Forty repeats at each sample size. The spread shrinks like $1/\sqrt{S}$; the exact value never moves.

▶ Notebook


A check with an answer key: the Beta coin



With the Beta(5, 3) prior we know the posterior: Beta(22, 16). Fit a Beta q from Beta(10, 10) using only the joint.

▶ Notebook


A check with an answer key: the Beta coin



By step 100, q has moved most of the way.

▶ Notebook

A check with an answer key: the Beta coin



After 600 steps: Beta(22.4, 16.2), with KL ≈ 0.0002 to the true posterior.

Notebook

One literal update of the mean

$$\sigma = 2 \text{ fixed}: \qquad L(\mu) = \text{const} + \frac{(\mu - 1)^2}{32}$$

Which term of the Gaussian KL depends on μ?

Notebook

One literal update of the mean

$$\sigma = 2 \text{ fixed}: \qquad L(\mu) = \text{const} + \frac{(\mu - 1)^2}{32}$$

$$\frac{\partial L}{\partial \mu} = \frac{\mu - 1}{16} = -0.0625 \ \text{ at } \mu = 0$$

> Differentiate it at μ = 0.

▶ Notebook

One literal update of the mean

$$\sigma = 2 \text{ fixed}: \qquad L(\mu) = \text{const} + \frac{(\mu - 1)^2}{32}$$

$$\frac{\partial L}{\partial \mu} = \frac{\mu - 1}{16} = -0.0625 \ \text{ at } \mu = 0$$

$$\mu^+ = 0 - 4(-0.0625) = 0.25: \quad \text{KL } 0.349 \to 0.336$$

Take one step with η = 4. What is the best KL with σ stuck at 2?

▶ Notebook

KL ≥ 0 by Jensen's inequality

$$\mathrm{KL}(q\|p) = -\mathbb{E}_q\left[\log \frac{p(\theta)}{q(\theta)}\right]$$

Which function makes Jensen's inequality useful?

▶ Notebook

KL ≥ 0 by Jensen's inequality

$$\mathrm{KL}(q\|p) = -\mathbb{E}_q\left[\log \frac{p(\theta)}{q(\theta)}\right]$$

$$\geq -\log \mathbb{E}_q\left[\frac{p(\theta)}{q(\theta)}\right]$$

Why does the inequality point this way?

▶ Notebook

KL ≥ 0 by Jensen's inequality

$$\mathrm{KL}(q\|p) = -\mathbb{E}_q\left[\log \frac{p(\theta)}{q(\theta)}\right]$$

$$\geq -\log \mathbb{E}_q\left[\frac{p(\theta)}{q(\theta)}\right]$$

$$= -\log 1 = 0$$

What makes the final expectation one?

▶ Notebook

Five draws can give a negative KL estimate

θ_i	$\log q(\theta_i) - \log p(\theta_i)$
5.623	−0.232
0.120	0.269
−5.536	−0.751
2.705	0.217
−2.254	0.030

> Some rows are negative. Could their average be negative?

▶ Notebook

Five draws can give a negative KL estimate

θ_i	$\log q(\theta_i) - \log p(\theta_i)$
5.623	−0.232
0.120	0.269
−5.536	−0.751
2.705	0.217
−2.254	0.030

$$\widehat{\mathrm{KL}}_5 \approx -0.093, \qquad \mathrm{KL}(q\|p) \approx 0.069$$

> Does this contradict KL ≥ 0?

▶ Notebook

The two PML candidates, scored by Monte Carlo

candidate	MC estimate ± 1 s.e.	numerical integral
$N(3, 1^2)$	9.54 ± 0.03	9.49
$N(3, 4.5^2)$	45.02 ± 0.21	45.15

> How large is the difference relative to the sampling uncertainty?

▶ Notebook

The two PML candidates, scored by Monte Carlo

candidate	MC estimate ± 1 s.e.	numerical integral
$N(3, 1^2)$	9.54 ± 0.03	9.49
$N(3, 4.5^2)$	45.02 ± 0.21	45.15

The narrow candidate scores far better. Both sit away from the mixture's mass, so both are poor.

> What role does the numerical integral play?

▶ Notebook

A fixed-noise finite difference checks the path

```
q = Normal(mu, sigma)
theta = q.rsample((S,))
loss = (q.log_prob(theta) - p.log_prob(theta)).mean()
```

What is different about rsample?

▶ Notebook

A fixed-noise finite difference checks the path

```
q = Normal(mu, sigma)
theta = q.rsample((S,))
loss = (q.log_prob(theta) - p.log_prob(theta)).mean()
```

derivative	autograd	finite difference
μ	−0.062466	−0.062466
σ	−0.372223	−0.372223

> What must stay fixed on both sides of the finite difference?

Notebook

Pathwise gradients are one route, not the only one

$$\nabla_\psi \mathbb{E}_{q_\psi}[f_\psi(\theta)] = \mathbb{E}_{q_\psi}[f_\psi(\theta)\nabla_\psi \log q_\psi(\theta) + \nabla_\psi f_\psi(\theta)]$$

> Which term was missing from the naive sample() gradient?

▶ Notebook

Pathwise gradients are one route, not the only one


$$\nabla_\psi \mathbb{E}_{q_\psi}[f_\psi(\theta)] = \mathbb{E}_{q_\psi}[f_\psi(\theta)\nabla_\psi \log q_\psi(\theta) + \nabla_\psi f_\psi(\theta)]$$

A score-function estimator accounts explicitly for the changing sampling distribution.

> Does this identity require θ = μ + σε?

▶ Notebook


A noisy ELBO estimate can cross the evidence



Can a finite-sample ELBO estimate cross the reference line?

▶ Notebook

A noisy ELBO estimate can cross the evidence



162 of 600 sampled estimates lie above the known evidence in this run.

> Which object remains a lower bound?

▶ Notebook

A higher ELBO does not prove a zero gap

$$\mathrm{ELBO}(q_{\mathrm{new}}) > \mathrm{ELBO}(q_{\mathrm{old}})$$

What does a true increase in the exact ELBO establish?

▶ Notebook

A higher ELBO does not prove a zero gap


$$\mathrm{ELBO}(q_{\mathrm{new}}) > \mathrm{ELBO}(q_{\mathrm{old}})$$

The family may not contain the posterior, and optimisation may stop short.

> Does a stable loss establish q = p(θ | $\mathcal{D}$)?

▶ Notebook


Check the fixed total with three Beta candidates



Add the ELBO and the gap. What is fixed?

▶ Notebook


Check the fixed total with three Beta candidates



Does the better candidate raise the ELBO by the amount its KL falls?

▶ Notebook

Check the fixed total with three Beta candidates



What happens at the exact posterior?

▶ Notebook

A mean must say which dimension it averages

object	shape	correct reduction
S scalar Beta draws	[S]	mean over S
S d-dimensional joint log densities	[S]	mean over S
S factorised coordinate log densities	[S, d]	sum over d, then mean over S

> Does every log_prob output still have an event dimension?

▶ Notebook

A mean must say which dimension it averages

object	shape	correct reduction
S scalar Beta draws	[S]	mean over S
S d-dimensional joint log densities	[S]	mean over S
S factorised coordinate log densities	[S, d]	sum over d, then mean over S

MultivariateNormal.log_prob already sums the event dimension. Do not sum the sample axis as well.

> What goes wrong if you sum over S before averaging?

▶ Notebook

An ordered sequence and a count have different evidence

$$p(H = 17 \mid r) = \binom{30}{17} r^{17}(1-r)^{13}$$

Which event contains more possible sequences?

▶ Notebook

An ordered sequence and a count have different evidence

$$p(H = 17 \mid r) = \binom{30}{17} r^{17}(1-r)^{13}$$

observed object	log evidence
one ordered sequence	−21.397
only the head count	−2.796

> Does this constant change the posterior or its KL gap?

▶ Notebook

Check the sampled objective with analytic Beta moments

$$\mathbb{E}_q[\log r] = \mathrm{digamma}(a) - \mathrm{digamma}(a+b)$$

> Which two expectations determine the coin ELBO?

Notebook

Check the sampled objective with analytic Beta moments

$$\mathbb{E}_q[\log r] = \text{digamma}(a) - \text{digamma}(a+b)$$

$$\mathbb{E}_q[\log(1-r)] = \text{digamma}(b) - \text{digamma}(a+b)$$

> Why write digamma rather than ψ here?

Notebook

Check the sampled objective with analytic Beta moments

$$\mathbb{E}_q[\log r] = \mathrm{digamma}(a) - \mathrm{digamma}(a+b)$$

$$\mathbb{E}_q[\log(1-r)] = \mathrm{digamma}(b) - \mathrm{digamma}(a+b)$$

Insert these into expected log likelihood + expected log prior − expected log q.

> How was the identity checked non-circularly?