


Atrial Fibrillation Pathway

This pathway has been developed from published guidance, in collaboration with local cardiologists.

This guidance is to assist GPs in decision making and is not intended to replace clinical judgment.



This pathway is currently under review – as some of the content may be out of date, it should be viewed as an archive document for information only. Please contact Camden MMT for up-to-date information/advice.

Reference  
<https://cks.nice.org.uk/atrial-fibrillation>

Comments & enquiries relating to medication:  
CCCG Medicines Management Team [mmt.camdenccg@nhs.net](mailto:mmt.camdenccg@nhs.net)  
Refer to current BNF or SPC for full medicines information

Clinical Contact for pathway queries: [Camden.pathways@nhs.net](mailto:Camden.pathways@nhs.net)

Pathway created by NCL  
Approved by Clinical Cabinet  
December 2017  
Review due November 2020

| <b>CHA<sub>2</sub>DS<sub>2</sub>Vasc</b>                                | <b>Score</b> | <b>HASBLED</b>                                                                         | <b>Score</b> |
|-------------------------------------------------------------------------|--------------|----------------------------------------------------------------------------------------|--------------|
| Congestive heart failure/LV dysfunct.                                   | 1            | Hypertension (uncontrolled, > 160 mmHg systolic)                                       | 1            |
| Hypertension                                                            | 1            | Chronic liver disease or Bili 2xULN with AST/ALT/ALP 3x ULN                            | 1            |
| Age = 75                                                                | 2            | Abnormal renal function (creatinine =200 umol/L, renal transplant or chronic dialysis) | 1            |
| Diabetes mellitus                                                       | 1            | Stroke                                                                                 | 1            |
| Stroke/TIA/systemic arterial embolism                                   | 2            | History of major bleeding <sup>1</sup> or predisposition                               | 1            |
| Vascular disease (prev. MI, peripheral arterial disease, aortic plaque) | 1            | Labile INRs, time in range less than 60%                                               | 1            |
| Age 65 -74                                                              | 1            | Elderly (age = 65 or frail condition)                                                  | 1            |
| Sex (male 0, female 1)                                                  | F 1          | Drugs (concomitant antiplatelet, NSAIDs etc) or alcohol abuse (1 point each)           | 1 or 2       |
| <b>Total score</b> (maximum score 9)                                    |              | <b>Total score</b> (maximum score 9)                                                   |              |

<sup>1</sup>Bleeding requiring hospitalisation and/or causing decrease in Hb >20 g/L and/or requiring =2 units blood transfusion

| <b>CHA<sub>2</sub>DS<sub>2</sub>-VASc score</b> | <b>Adjusted stroke rate (%/year)</b>  |
|-------------------------------------------------|---------------------------------------|
| 0                                               | 0%                                    |
| 1                                               | 1.3%                                  |
| 2                                               | 2.2%                                  |
| 3                                               | 3.2%                                  |
| 4                                               | 4.0%                                  |
| 5                                               | 6.7%                                  |
| 6                                               | 9.8%                                  |
| 9                                               | 15.2%                                 |
| <b>HAS-BLED score</b>                           | <b>Major bleed s per 100 pt years</b> |
| 0                                               | 1.13                                  |
| 1                                               | 1.02                                  |
| 2                                               | 1.88                                  |
| 3                                               | 3.74                                  |
| 4                                               | 8.70                                  |
| 5                                               | 12.50                                 |
| 6 -9                                            | Insufficient data                     |

**This pathway is currently under review – as some of the content may be out of date, it should be viewed as an archive document for information only. Please contact Camden MMT for up-to-date information/advice.**