

Flash and Continuous Blood Glucose Devices Prescribing

Primary Care Approval Tool

April 25

An Assessment Tool for Prescribing and Monitoring primary care prescribing of Blood Glucose Monitoring Devices

Background

1. The advancement of Flash and Continuous Blood Glucose Monitoring has greatly improved the life of patients with Diabetes and immense benefits and cost saving to the health system.
2. It is important that Staffordshire and Stoke on Trent Flash Glucose Monitoring and Dexcom One Commissioning Policy is utilised to support adherence for the initiation and prescribing of FCGM in the ICS.
3. This document is a Decision Support Tool (DST) to support primary care prescribing of FCGM devices. This will ensure:
 - The correct patient is prescribed FCGM device.
 - The SSOT FCGM Policy is adhered to in the decision to initiate or prescribe at the request of secondary care.
4. The following checklist will be used by primary care teams when the request to initiate FCGM is received from secondary care colleagues.
5. If the request to initiate or continue prescribing in primary care does not conform to policy which was agreed with secondary care, the standard decline to prescribe form will be filled and sent back to the requestor in accordance with SSOT ICB Policy
6. The questions in the DST are the criteria in the SSOT FCGM Policy.

NHS Staffordshire and Stoke-on-Trent Integrated Care Board

Name of Practice: Organisation Name

Practice Code: Organisation National Practice Code

Patient Name: Full Name NHS No: NHS Number

Does the FCGM that has been requested have a GREEN/AMBER or AMBER-I designation on the formulary?

Yes ☐ No ☐

If yes to the above question, then continue the assessment, if no, **complete a decline to prescribe form**

Policy criteria

Is the patient a Type 1 Diabetic clinically indicated as requiring intensive therapy monitoring greater than 8 times daily

Yes ☐ No ☐

Patient has any form of Diabetes on Dialysis AND on insulin (who are clinically indicated as requiring intensive monitoring >8 times per day as demonstrated on a meter download/review over the last 3mths)

Yes ☐ No ☐

Diabetes associated with cystic fibrosis on insulin treatment

Yes ☐ No ☐

Pregnant women with type 1 diabetes

Yes ☐ No ☐

People with Type 1 diabetes unable to routinely self-monitor blood glucose levels due to disability that requires carers to support glucose monitoring and insulin management.

Yes ☐ No ☐

People with Type 1 diabetes for whom the specialist MDT team determines has occupational (e.g. working in insufficiently hygienic conditions to safely facilitate finger-prick testing) or psychosocial circumstances that warrant a 6-month trial with adjunct support.

Yes ☐ No ☐

Patient has Type 1 Diabetes and is a previous self funder of FCGM whose responsible clinician is satisfied that their clinical history has shown improvement to their HBA1c

Yes ☐ No ☐

Patient with Type 1 Diabetes with recurrent severe hypoglycaemia or impaired awareness of hyperglycaemia (as per NICE Guidance and SSOT CGM Policy and FGM Policy)

Yes ☐ No ☐

People with Type 1 diabetes or insulin treated Type 2 diabetes who are living with a learning disability and recorded on their GP Learning Disability register.

Yes ☐ No ☐

If Yes to any of the policy criteria, please issue and put on repeat.

If No to all of the policy criteria, please complete a Decline to Prescribe form – search 'MOT SSOTP Decline to Prescribe' and refer back to prescriber

Name of Clinician: Current User

Date: Short date letter merged