

COVID-19 Vaccinations – Allergy Service Review Referral

This form is for use when an identified patient has a history of allergy requiring an allergy review with a view to vaccination with full resuscitation facilities.

Note: Allergy reviews do not routinely involve direct contact with the patient and may be based on referral details alone.

PATIENT DETAILS			
Patient Name	Given Name Surname	GP Practice	Organisation Name Organisation Full Address (single line)
Patient Address	Home Full Address (single line)	Patient Date of Birth	Date of Birth
		Contact Number	Patient Home Telephone Patient Mobile Telephone
NHS Number	NHS Number	Hospital Number	Hospital Number

ALLERGY REFERRAL CRITERIA			Recommended Action
General Allergy History	Known allergy to polyethylene glycol (PEG)	☐	Refer for Immunology Review
COVID-19 Vaccine Specific Severe Allergy	Immediate (onset within 2 hours) systemic symptoms (anaphylaxis) to mRNA vaccine	☐	Refer for Immunology Review
	Delayed (onset beyond 2 hours) urticaria/angioedema requiring hospital attention, following dose of mRNA vaccine	☐	Refer for Immunology Review
For all other clinical presentations of allergies or history of allergies, further information is available in Appendix 1 as to how these individuals can be vaccinated safely within COVID-19 vaccination sites.			

Previous COVID-19 vaccine doses			
Dose	Date given	COVID-19 vaccine brand received	Adverse reaction/allergy Please include detail on timings of onset of symptoms and treatment received
1 st Dose			
2 nd Dose			
3 rd Primary Dose			
1 st Booster Dose			
2 nd Booster Dose			
Any other relevant information (please attach where available any relevant Allergy Clinic Letters, Ambulance Reports and, A&E notes):			

REFERRING CLINICIAN			
Signature		Date	Short date letter merged
Name (printed)	Free Text Prompt	Email	Organisation E-mail Address
Professional title			
Working at COVID-19 Vaccination Site?	Yes ☐	No ☐	Site Name:
Please send this referral via e-RS as an Advice & Guidance request to Adult Allergy at UHNM .			

Relevant Medical History:
Problems

Medication

Allergies

Appendix 1: Exclusions from referral

EXCLUSIONS FROM REFERRAL – these individuals should be managed within COVID-19 vaccination sites. Please see The Green Book for further advice on managing allergies https://www.gov.uk/government/publications/covid-19-the-green-book-chapter-14a		
	Allergy issue	Action within vaccination sites
History of Allergy	Family history of allergies	Proceed with mRNA vaccination Consider 15 minute observation period Some patients (e.g. those with mastocytosis) may benefit from pre-treatment with antihistamine to reduce allergic symptoms
	Previous allergic reaction (including anaphylaxis) to a food, insect sting and most medicines (where trigger has been identified e.g. Penicillin)	
	Previous non-systemic reaction to a vaccine	
	Hypersensitivity to non-steroidal anti-inflammatory drugs e.g. aspirin, ibuprofen	
	Mastocytosis	
	Known allergy to polysorbate 80 (not in mRNA COVID-19 vaccines)	
Adverse Reactions	Immediate (onset within 2 hours) anaphylaxis to multiple, different drug classes, with trigger unidentified, OR a medicine likely to contain PEG (e.g. depot steroid injection, macrogol laxatives)	Review previous drug history – if a medication containing PEG has been tolerated before then mRNA vaccine can be given as normal
	Anaphylaxis to a vaccine or injected antibody preparation	
	Idiopathic anaphylaxis	
	Delayed (onset beyond 2 hours) urticaria/angioedema requiring medical attention, following dose of Vaxzevria®/AstraZeneca	Proceed with mRNA vaccination Consider 15 minute observation period
	Fainting/vasovagal - strong central pulse with rapid recovery. Malaise may persist, consciousness should recover within a few minutes.	
	Other Non-Allergic Reaction – non-urticarial skin reaction or non-specific symptoms	
	Panic Attack - hyperventilation ± paraesthesiae (numbness and tingling) in arms and legs. Erythematous rash associated with anxiety. Hypotension, pallor or wheezing <u>not</u> present.	Proceed with mRNA vaccination as normal Support and reassure patient
	Swelling or Rash (at any time post-vaccination) - local to injection site only	Proceed with mRNA vaccination Consider pre-treatment with non-sedating anti-histamine, 30 minutes prior to vaccination Observe for at least 30 minutes
	Urticaria/Angiodema - Self-limiting reaction/resolved with oral anti-histamine	Proceed with mRNA vaccination Consider pre-treatment with non-sedating anti-histamine, 30 minutes prior to vaccination Observe for at least 15 minutes
	Systemic symptoms, within 2 hours of vaccination, but no objective symptoms of anaphylaxis: - No respiratory or cardiovascular compromise - Symptoms rapidly resolved with maximum 1 dose of IM adrenaline	Proceed with mRNA vaccination Consider pre-treatment with non-sedating anti-histamine, 30 minutes prior to vaccination Observe for at least 30 minutes