

Referral Form/ Advice & Guidance
CoViD Vaccine Allergy Service (March 2021)
18 years and over

THIS FORM MUST BE COMPLETED IN FULL AND ATTACHED WITH THE APPROPRIATE CLINICAL INFORMATION TO THE E-REFERRAL SERVICE

REFERRALS WITHOUT FORMS WILL BE REJECTED

Patient details		Referring GP details	
Surname	Surname	Referring GP	
Forename(s)	Given Name	Practice name Organisation Name	
Address Home Full Address (single line)		Practice address Organisation Full Address (single line)	
Date of birth	Date of Birth	Telephone number Organisation Telephone Number	
NHS Number	NHS Number	GP practice code Organisation National Practice Code	

Clinicians are reminded that the following can be vaccinated within primary care:

- Previous allergic reaction (including anaphylaxis) to a food, insect sting and most medicines (where trigger has been identified and is not found within the intended Covid vaccine)
- Family history of allergies
- Previous non-systemic reaction to a vaccine
- Hypersensitivity to nonsteroidal anti-inflammatory drugs e.g. aspirin, ibuprofen
- Mastocytosis
- Patients who have allergic reactions to the first dose of COVID-19 vaccine, who have experienced:
 - Swelling or rash local to injection site only within or longer than two hours after vaccination – (Observe for 30 minutes for *second* dose)
 - Urticaria or angioedema more than two hours after vaccination and which was self-limiting or resolved with oral antihistamine (consider pre-treatment with non-sedating antihistamine 30 minutes prior to vaccination)

***Anaphylaxis**

Anaphylaxis is likely when all of the following three criteria are met:

- sudden onset and rapid progression of symptoms
- life-threatening airway and/or breathing and/or circulation problems
- skin and/or mucosal changes (flushing, urticaria, angioedema).

The following supports the diagnosis:

- exposure to a known allergen where the patient is already known to be allergic.

Remember:

- **skin or mucosal changes alone are not a sign of an anaphylactic reaction**
- skin and mucosal changes can be subtle or absent in up to 20% of reactions (some patients can have only a decrease in blood pressure, i.e. a circulation problem)
- there can also be gastrointestinal symptoms (e.g. vomiting, abdominal pain, incontinence).

Most anaphylactic reactions occur in individuals who have no known risk factors.

Systemic symptoms would usually include generalised urticaria and/or angioedema, but not meeting the full diagnostic criteria for anaphylaxis.

Referral criteria – Section 1 or 2 to be completed

Section 1

First dose Covid-19 Vaccination - Managing patients who have a history of allergic reactions (Flow Chart 1)

Patient Characteristics

- Previous allergic reaction (including anaphylaxis*) to a food, insect sting and most medicines (where the trigger has been identified and is not a component of the intended Covid-19 vaccine)
- Family history of allergies
- Previous non-systemic reaction to vaccine (i.e. swelling/rash localised to injection site)
- Hypersensitivity to NSAIDs or aspirin
- Mastocytosis



Proceed with vaccination as normal according to local guidelines

Patient Characteristics

- History of anaphylaxis* to multiple drug classes (not vaccines/biologics)
- Confirmed diagnosis of polyethylene glycol (PEG) allergy or systemic allergic reaction to macrogol
- Systemic reaction or anaphylaxis to medicine likely to have contained PEG
- History of a systemic allergic reaction or anaphylaxis* to a mRNA vaccine
- History of systemic allergic reaction or anaphylaxis* to a medicine containing any component of the vaccine
- History of unexplained/idiopathic anaphylaxis



REFER to Covid vaccine Allergy service via Advice and Guidance
Use referral form and tick referral reason **ONE**

☐

Patient Characteristics

- History of anaphylaxis* or systemic allergic reaction to non-Covid vaccine or biologics (e.g. infliximab, adalimumab, anakinra etc.)




REFER to Covid vaccine Allergy service via Advice and Guidance
Use referral form and tick referral reason **TWO**

☐

Referral criteria – Section 1 or 2 to be completed

Section 2

Second dose Covid-19 Vaccination - Managing patients who have allergic reactions to the first dose of Covid-19 vaccine (Flow Chart 2)



Additional Patient Information	All must apply
This patient is willing to undergo vaccination should it be offered.	☐
To manage expectation, I have discussed with the patient the fact they will be referred for a possible vaccination but there is no guarantee that the intervention will be the required outcome following the consultation with the secondary care specialist.	☐
Depending on the advice the vaccination may take place in either of the following settings, UHDBFT, CRHFT or a local vaccination service	☐

Part B – Reason for referral (a detailed history is preferred to allow the Specialist to make an informed and will reduce requests for additional information)	
Salutations:	Dear colleague,
Preamble/ context:	Consultations Thank you, History should contain: - Date of reaction - Vaccination type & brand e.g. flu / Influvac – Tetra - Details of diagnosis e.g. polysorbate 80 - Details of attending clinician – nurse, GP, Hospital clinician

Problems

Relevant SH & FH:

Smoking status	Single Code Entry: Tobacco smoking consumption
Alcohol	Single Code Entry: Alcohol consumption
Occupation	
Ethnicity	Ethnic Origin
Veteran	Single Code Entry: Military veteran
Detail which might assist timely discharge:	

Medication

Allergies

Useful values:

BP Single Code Entry: O/E - blood pressure reading	Pulse rate Single Code Entry: Pulse rate	Height Single Code Entry: Standing height	Weight Single Code Entry: Body weight	BMI Single Code Entry: Body mass index	HbA1C Single Code Entry: Haemoglobin A1c level - International Federation of Clinical
--	--	---	---	--	---

Name: Given Name Surname NHS No: NHS Number

					Chemistry and Laboratory Medicine standardised
--	--	--	--	--	---

Please embed any attached items here .e.g. Vaccination History
Please note any individual patient requirements here (e.g. Wheelchair user).