

My Maintenance and Reliever Treatment (MART) personal asthma action plan

Name:	Given Name Surname	Height:	Single Code Entry: Standing height	Weight:	Single Code Entry: Body weight
Date of birth:	Date of Birth	Best Peak Flow:		Predicted Peak Flow:	
NHS number:	NHS Number	Allergies:		Triggers:	
Hospital number:	Hospital Number				
GP Name	Usual GP Full Name		Tel:	Organisation Telephone Number	
Practice Nurse name:			Tel:		
Hospital Practitioner			Tel:		

My asthma treatment

Medication / Inhaler Colour	Strength	Dose and Frequency	Device / Type of Medication
Issued by:		Date:	

The Four Stages of Asthma Control

Peak flow recorded from 5 years and above.

Stage 1	Key messages	Recommended Actions
Your asthma is under good control when: - There is no cough, wheeze, chest tightness or shortness of breath. - You are able to do usual activities. - You do not need extra doses of your MART inhaler. - Your peak flow is 75% -100% of best (5 years and above). 100% 75%	Remember: Continue to take your maintenance medication(s).	Take your maintenance treatment as prescribed to keep you well. Maintenance medication (s): Attend a planned asthma review once or twice per year.


Stage 2	Key messages	Recommended Actions
Your asthma control is not good when you have any of the following: - Cough, wheeze, chest tightness or shortness of breath. - Waking at night more than once per week due to asthma symptoms. - You are able to do some, but not all usual activities. - You need extra doses of your MART inhaler. - Your peak flow is 60%-75% of best (5 years and above). 75% 60%	Important: Seek a medical review (GP/111) if extra MART medication is: - Needed three times a week or more. - Needed more than once in 24 hours, even if you feel better.	Maintenance And Reliever Therapy (MART) For symptom relief take 1 puff/inhalation of your MART inhaler. If symptoms are not improving after 3-5 minutes repeat. This should make you feel better and your symptoms should have improved. If your symptoms are not improving move to stage 3.

Stage 3	Key messages	Recommended Actions
---------	--------------	---------------------

You are having an asthma attack when: - Symptoms are more frequent. - MART inhaler is less effective and is needed more than once in 24 hours. - MART inhaler relief is not lasting 4 hours. - You are breathless on normal activity. - Your peak flow is 60% or less (5 years and above). 60%	Important: Seek a medical review (GP/111) within the next 24 hours if: - You have needed to use your MART maximum puffs/inhalations in one go, or you have had your maximum daily amount even if you feel better. - Extra MART medication is not lasting at least 4 hours.	Maintenance And Reliever Therapy (MART) Take 1 more puff/inhalation of your MART inhaler. If symptoms are not improving after 3-5 minutes you can repeat taking 1 more puff/inhalation every 3-5 minutes. You can take up puffs/inhalations to in one go If your symptoms have now improved still arrange a medical review. You may need to start a course of prednisolone You can take a maximum of puffs/inhalations daily (including your regular morning and evening puffs/inhalations). If you have reached your maximum puffs/inhalations in one go and your symptoms are not improving move to stage 4.
--	---	--

Stage 4	Key messages	Recommended Actions
You are having a severe asthma attack when: - Symptoms continue to get worse. - Symptoms are not responding to MART inhaler - You are too breathless or exhausted to speak. - Peak flow is 40% or less (5 years and above). 40%	Important: Call 999 immediately as there is no or poor response to your MART inhaler More than your daily maximum puffs/inhalations of your MART inhaler may be used in this emergency situation.	Try to keep calm, call 999 and tell them you are having a severe asthma attack. Whilst you continue to have symptoms, and you are waiting for the ambulance, you can keep taking 1 additional puff/inhalation every 3-5 minutes. If the ambulance has not arrived after 10 minutes call 999 again. If you don't have your MART inhaler with you, you can use a blue reliever inhaler via spacer. Take one puff every 30–60 seconds up to a maximum of 10 puffs. Call 999 and if the ambulance has not arrived after 10 minutes call 999 again.

Asthma and Your Airways



Handy Tips

- Each inhaler comes with a set amount of doses. Keep a check on the expiry date and dose counter or count out the doses if no dose counter to avoid running out of treatment. Once empty pMDI's will only deliver the propellant.
- Always carry your MART inhaler dry powder device/pMDI and spacer with you including at school and work.
- Do not run out of your inhalers

Inhaler maintenance

- Rinse your mouth and gargle after using steroid Inhalers.
- Spacers need to be washed once weekly in warm soapy water and allow to air dry.
- If you use a spacer, replace this every 6 to 12 months.
- Access www.asthmaandlung.org.uk to check your inhaler technique
- If extra MART medication is needed regularly three times a week or more, arrange a medical review.

Asthma Review

- Please bring all of your medications and personal asthma action plan with you if you are admitted to hospital or if you attend any outpatient appointments.
- Please arrange a review with your GP, or asthma nurse within two working days after an admission to hospital or after receiving emergency care.
- If you have commenced a rescue course of prednisolone, you should contact your GP for a review.
- You will need to attend your asthma reviews and take your medication as directed.