

## Derbyshire Shared Care Pathology Guidelines

### Hypomagnesaemia in Adults

#### Purpose of Guideline

The management of patients with newly diagnosed hypomagnesaemia.

Magnesium is crucial in the appropriate reabsorption of both potassium and calcium in the kidney. Therefore, in some cases of hypocalcaemia and hypokalaemia it is essential to replace magnesium to enable the correction of both calcium and potassium.

#### When is hypomagnesaemia considered a medical emergency?

- Magnesium  $<0.30$  mmol/L      Likely to be medical emergency



Magnesium results  $\leq 0.40$  mmol/L may be telephoned to OOH GP service when the GP practice is closed, dependent on other results.

These limits are for guidance only; the severity of the patient's symptoms, and other factors, e.g. rapidity of the fall in magnesium, serum calcium and potassium concentrations, renal function, will also determine whether the patient requires emergency admission.

#### When is hypomagnesaemia suspected?

- Hypomagnesaemia is more common than people think with probably 90% not identified clinically. Symptoms are similar to hypocalcaemia, and may include neuromuscular issues like muscle twitching, tremors, and weakness, as well as cardiovascular symptoms such as palpitations and arrhythmias. The number and severity of symptoms will be dependent on the degree of hypomagnesaemia. Mild hypomagnesaemia may only present with vague muscular aches and pains.

### Incidence of Hypomagnesaemia

- 2% of the general population
- 7-11% of hospital patients
- Poorly Controlled Diabetes up to 25%
- Critically Ill Patients up to 65%
- Alcoholics up to 80%
- Haematological Malignancies up to 90%

### What are the causes of hypomagnesaemia?

Hypomagnesaemia is usually caused by GI and renal loss and redistribution.

#### Renal causes include:

- Drugs
  - Proton Pump Inhibitors (common cause)
  - Diuretics, especially Loop diuretics
  - Cytotoxic drugs
  - Aminoglycosides
  - Immunosuppressants
  - Theophylline
- Osmotic diuresis (diabetes)
- Inherited Disorders (rare, but include Gitelman's syndrome)

#### GI causes include:

- Reduced Intake
  - Dietary deficiency (rare)
- Reduced Absorption
  - Diseases causing Malabsorption (e.g. Coeliac Disease)
  - Chronic Diarrhoea
  - Laxative Abuse
  - Fistula
  - Short Bowel Syndrome
  - Inherited Transport Defect (rare)

### What happens next?

1. Review the medication - stop PPI and review diuretics (the recommendation is for permanent cessation of PPI which could be substituted by Famotidine. If PPI is needing to be reconsidered for challenging GORD symptoms, seek A&G from gastroenterology or endocrinology)
- 2.
3. The majority of patients who develop hypomagnesaemia are already under the care of hospital physicians, and referral is usually straight forward (e.g. Oncology)
4. For unknown causes of hypomagnesaemia ( $<0.50$  mmol/L), check U&E and calcium with repeat Mg level.

- If repeat Mg  $>0.4$  mmol/L and patient is asymptomatic, immediate referral is not required. Consider oral replacement (as below) and referral to appropriate specialty dependent on the patients' history. Endocrine advice is available by choosing 'Advice and Guidance' from NHS e-Referrals.
- If repeat Mg  $\leq 0.4$  mmol/L or with concurrent hypocalcaemia or hypokalaemia, consider referring patient to SDEC
- Symptomatic patients tend to have more profound hypomagnesaemia (often  $<0.30$  mmol/L) with concomitant hypokalaemia and hypocalcaemia. Consider referral to SDEC in these cases.

## Treatment

Treatment of severe hypomagnesaemia is usually intravenous and should always be carried out in hospital. Intramuscular magnesium injections are very painful and are not recommended.

### Oral administration

This is used for chronic magnesium loss or moderately severe hypomagnesaemia i.e. where there is a serum magnesium level of approximately 0.4 - 0.7 mmol/L and patients are asymptomatic. Give up to 50 mmol/L day.

Suggested initial treatment is;

- Co-magaldrox 10-20 mL qds (although this product is off-label, this remains the preferred first-line treatment)

During oral replacement monitor response by rechecking magnesium 1-2 weekly initially depending on clinical context. Reversal of hypomagnesaemia with oral replacement may take 6-8 weeks. Long term maintenance replacement may be needed if a reversible cause is not found and removed. In these circumstances, aim to maintain normal serum magnesium using the lowest effective dose of oral replacement.

Diarrhoea tends to limit the amount of magnesium that can be given orally; if diarrhoea develops reduce the dose. The aluminium contained in Mucogel may reduce the chance of diarrhoea.

*(Note: The other product occasionally appropriate for use in this context is Magnesium glycerophosphate (1 tablet = 4 mmol Mg, Dose= 1-2 tablets three to four times daily (12-32 mmol/day). Prescription should be reserved for cases of intolerance to Mucogel.)*

***Patients with Renal Impairment should be treated with caution. Suggest obtain renal advice before commencing treatment if CKD 4 – 5.***

## Contacts

CHISCP11  
Revision:7

Date of issue: April 2026  
Expiry date: 30th April 2028  
Approved by: J Monaghan

### **Derby**

Duty Biochemist  
On Call Consultant Biochemist  
Endocrinology Advice  
Consultant Phone Triage

01332 789383 (8am to 7pm, Mon – Fri)  
Via RDH switchboard, 01332 340131 (24/7)  
07879 115507 (9am – 5pm, Mon – Fri)  
Via RDH switchboard, 01332 340131  
(10am to 6pm Monday-Friday)  
01332 788707

MAU and ACC

### **Chesterfield**

Duty Biochemist  
On Call Consultant Biochemist  
A&E, EMU and CDU

01246 512212 (9am to 5pm, Mon – Fri)  
Via CRH switchboard, 01246 277271  
Via CRH switchboard

### **Burton**

Duty Biochemist  
Endocrinology advice

01283 566333 ext 4233  
Via endocrine secretaries 01283 593259

| Document title                    | Hypomagnesaemia in adults |                                                                      |                               |
|-----------------------------------|---------------------------|----------------------------------------------------------------------|-------------------------------|
| Revision                          | 7                         | Document Reference                                                   | CHISCP11                      |
| Reason for Review                 | Routine Review            |                                                                      |                               |
| Review history                    | Version/Date              | Authors                                                              | Reason for review             |
|                                   | V1 Jan 2012               | Dr Rustam Rea, Dr Roger Stanworth, Dr Paul Masters & Dr Nigel Lawson | New Document                  |
|                                   | V2 Mar 2014               | Dr R Stanworth, Dr P Blackwell, Mrs H Seddon                         | Routine Review                |
|                                   | V3 June 2016              | Dr R Stanworth, Dr P Blackwell, Mrs H Seddon                         | Routine Review                |
|                                   | V4 Oct 2018               | Dr R Stanworth, Dr P Blackwell, Mrs H Seddon                         | Routine Review                |
|                                   | V5 Oct 2021               | Dr D Hughes, Dr A Ugur, Dr P Shillo, Dr P Blackwell, Mrs H Seddon    | Routine Review                |
|                                   | V6 Apr 2024               | Dr R Stanworth, Dr P Blackwell, Mrs H Seddon                         | Routine Review                |
| Approval                          | Date                      | Name(s)                                                              | Designation                   |
| Secondary Care UHDB               | 22.04.26                  | Dr A Ugur, Dr D Hughes UHDB Biochemistry/Endocrinology MDT           | Consultant Endocrinologists   |
| Secondary Care CRH                | 09.04.26                  | Dr P Shillo                                                          | Consultant Endocrinologist    |
| Primary Care                      | 27.04.26                  | Dr P Blackwell                                                       | GP                            |
| Laboratory (Derbyshire Pathology) | 28.04.26                  | Ms S Knowles                                                         | Consultant Clinical Scientist |

CHISCP11  
Revision:7

Date of issue:April 2026  
Expiry date:30th April 2028  
Approved by:J Monaghan