

Written Assignment 3

Niraj Venkat

Exercise 1

An *analytic* function has infinitely many derivatives in its domain. For our proof, we only need a function to be twice-differentiable.

We state Liouville's theorem:

Suppose f is analytic in the open set that contains a ball $B_r(z_0)$ of radius r centered at z_0 , and that $|f(z)| \leq c$ holds on $\partial B_r(z_0)$ for some constant c . Then for all $k \geq 0$ we have:

$$|f^{(k)}(z_0)| \leq \frac{k! c}{r^k}$$

Simply put: if f is bounded, f must be constant.

This holds for *harmonic* functions as well, in particular our function $\phi : M \rightarrow \mathbb{R}$ where $k = 2$ and $\Delta\phi = 0$.

Exercise 2

This admits different proofs:

- Suppose we had a function f such that $\Delta f = \Delta\phi = c$. Then by linearity of the Laplacian we have:

$$\Delta(\phi - f) = \Delta(f - \phi) = 0$$

Liouville's Theorem says that both functions $\phi - f$ and $f - \phi$ must be constant.
 $c = 0$ is the only value that satisfies this condition.

- Stokes' theorem (divergence theorem) says the following:

For any vector field, we can relate the divergence of the vector field within the volume V to the outward flux of the vector field through the surface $S = \partial V$.

In our case, we apply this theorem to the vector field which is the gradient $\nabla\phi$.

The Laplacian ($\Delta = \nabla \cdot \nabla$) is the divergence of gradient, so for some compact domain M without boundary ($\partial M = \emptyset$) we have:

$$\begin{aligned} \int_M \nabla \cdot \nabla\phi \, dV &= \int_{\partial M} \nabla\phi \cdot \hat{n} \, dS = 0 \\ \implies 0 &= \int_M \Delta\phi \, dV = \int_M c \, dV = c \, \text{vol}(M) \end{aligned}$$

Because M has some non-negative volume, c cannot be an arbitrary constant. c has to be zero.

Exercise 3

According to Stokes' theorem:

$$\begin{aligned}
 \int_{\partial} g \star df &= \int d(g \star df) \\
 &= \int dg \wedge \star df + \int g \wedge d \star df \\
 &= \int dg \wedge \star df + \int g \wedge \star \Delta f \\
 &= \langle \nabla g, \nabla f \rangle + \langle g, \Delta f \rangle
 \end{aligned}$$

On the other hand:

$$\int_{\partial} g \star df = \langle g, N \cdot \nabla f \rangle_{\partial}$$

This proves Green's first identity:

$$\langle \Delta f, g \rangle = -\langle \nabla f, \nabla g \rangle + \langle N \cdot \nabla f, g \rangle_{\partial}$$

Exercise 4

If there is no boundary:

$$\langle g, N \cdot \nabla f \rangle_{\partial} = 0$$

If $f = g$, then Green's first identity becomes:

$$\begin{aligned}
 \langle \Delta f, f \rangle &= -\langle \nabla f, \nabla f \rangle + 0 \\
 &= -\|\nabla f\|^2 \leq 0
 \end{aligned}$$

This means the operator Δ is negative semidefinite, and $-\Delta$ is positive semidefinite (PSD).

The set of PSD matrices is convex, as such many tools in numerical linear algebra (eg: semidefinite programming) can be used to find the minimum of the quadratic functions they describe (eg: quadratic forms $\vec{x}^T A \vec{x}$).

Exercise 5

Using figure on the next page as a reference:

$$\begin{aligned}
 w &= b_1 + b_2 \\
 h &= |\overline{ip}| \\
 \cot \alpha &= \frac{b_1}{h} \\
 \cot \beta &= \frac{b_2}{h} \\
 \frac{w}{h} &= \cot \alpha + \cot \beta
 \end{aligned}$$


Exercise 6

Triangle ijk has base $b = |e| = b_1 + b_2$ and height h .

Let e be the edge vector for the base edge jk , with i as the opposite vertex.

The interpolating hat functions ϕ are similar to their smooth counterpart, the Dirac delta δ . It is equal to one at their associated vertex and zero at all other vertices. We associate ϕ with vertex i .

We can expand the interpolating hat function ϕ in a Taylor series expansion about the point x_0 . Since the hat function is linear, all higher order terms go to zero.



$$\phi(x) = \phi(x_0) + \nabla\phi \cdot (x - x_0)$$

Plugging our vertices into this expression:

$$\phi(i) = 1 \tag{1}$$

$$\phi(j) = \phi(i) + \nabla\phi \cdot (j - i) = 0 \tag{2}$$

$$\phi(k) = \phi(i) + \nabla\phi \cdot (k - i) = 0 \tag{3}$$

Subtracting (3) - (2):

$$\begin{aligned} \phi(k) - \phi(j) &= \nabla\phi \cdot (k - i) - \nabla\phi \cdot (j - i) \\ &= \nabla\phi \cdot (k - j) = 0 \end{aligned}$$

We get that direction of $\nabla\phi$ is perpendicular to e . Next we calculate the magnitude of $\nabla\phi$ by substituting (1) into (2):

$$\begin{aligned} 0 &= \phi(i) + \nabla\phi \cdot (j - i) = 0 \\ &= 1 + \nabla\phi \cdot (j - i) \\ 1 &= \nabla\phi \cdot (i - j) \\ &= |\nabla\phi| |i - j| \cos(\pi - \alpha) \\ &= |\nabla\phi| h \\ |\nabla\phi| &= \frac{1}{h} = \frac{b}{bh} = \frac{|e|}{2A} \end{aligned}$$

We call e^\perp the vector perpendicular to e with magnitude $|e|$. This gives our desired gradient:

$$\nabla\phi = \frac{e^\perp}{2A}$$

Exercise 7

Using our previous result $\nabla\phi = \frac{e^\perp}{2A}$:

$$\begin{aligned} \langle \nabla\phi, \nabla\phi \rangle &= \frac{e^\perp \cdot e^\perp}{4A^2} \\ &= \frac{b^2}{4b^2h^2} \\ &= \frac{1}{2} \frac{b}{2h} \\ &= \frac{1}{2} (\cot \alpha + \cot \beta) \end{aligned}$$

We consider vertices i and j with opposite edge vectors making angle θ (labelled β in our figure). The opposite edge vectors as labelled in our figure are $e_i = \overrightarrow{jk}$ and $e_j = \overrightarrow{ki}$

$$\begin{aligned}\langle \nabla \phi_i, \nabla \phi_j \rangle &= \frac{e_i^\perp \cdot e_j^\perp}{4A} \\ e_i^\perp \cdot e_j^\perp &= |e_i||e_j| \cos(\pi - \theta) = -|e_i||e_j| \cos \theta \\ A &= \frac{1}{2}(e_j \times -e_i) = \frac{1}{2}(e_i \times e_j) = \frac{1}{2}|e_i||e_j| \sin(\pi - \theta) = \frac{1}{2}|e_i||e_j| \sin \theta\end{aligned}$$

Altogether:

$$\langle \nabla \phi_i, \nabla \phi_j \rangle = -\frac{1}{2} \cot \theta$$

In our diagram, primal mesh elements are solid, dual are dashed.

Since each dual edge connects the centers of two circles with the corresponding primal edge as a common chord, each dual edge is a perpendicular bisector of the primal edge.

This can easily be proven:

- Let the two circumcircles have circumcentres O and O' where AB is the common chord, with intersection point P .
- $OA = OB$ and $O'A = O'B$ (radii of same circle)
So, $\triangle OAO'$ and $\triangle OBO'$ are congruent.
- With this we can prove $AP = BP$ and $\angle APO = \angle BPO = \frac{\pi}{2}$.
- $\therefore OO'$ is the perpendicular bisector of AB .

Using inscribed angle theorem, we get:

$$\alpha = \frac{1}{2} \angle AOB \text{ and } \beta = \frac{1}{2} \angle AO'B.$$

Using angle bisector theorem, we get:

$$\angle AOO' = \frac{1}{2} \angle AOB = \alpha \text{ and } \angle BO'O = \frac{1}{2} \angle BO'A = \beta.$$

From the previous result relating the ratio w/h to sum of cotangents we have:

$$\begin{aligned}\frac{OO'}{AP} &= \frac{OO'}{AB/2} \\ \frac{OO'}{AB} &= \frac{1}{2}(\cot \alpha + \cot \beta) \\ \implies \frac{|e_{ij}^*|}{|e_{ij}|} &= \frac{1}{2}(\cot \alpha + \cot \beta)\end{aligned}$$

