

Measles, Mumps, Rubella (MMR) Immunity Profile

Test Details

METHODOLOGY

See individual tests.

RESULT TURNAROUND TIME

1 - 2 days

Turnaround time is defined as the usual number of days from the date of pickup of a specimen for testing to when the result is released to the ordering provider. In some cases, additional time should be allowed for additional confirmatory or additional reflex tests. Testing schedules may vary.

Test Includes

Measles IgG antibodies; Mumps IgG antibodies; Rubella IgG antibodies

Use

This test is used to aid in the determination of serological status to measles, mumps, and rubella viruses. A positive result generally indicates exposure to virus or previous vaccination. A positive result is considered adequate laboratory evidence of immunity.

Custom Additional Information

Measles

Acceptable presumptive evidence of Measles Immunity (at least one of the following):

- Documentation of age-appropriate vaccination against measles virus.
- Laboratory evidence of immunity (IgG in serum; equivocal results should be considered negative).
- Laboratory confirmation of measles.
- Birth before 1957.

Persons who have measles-specific IgG antibody that is detectable by any commonly used serological assay are considered to have adequate laboratory evidence of measles immunity. Persons with an equivocal serologic test result do not have adequate presumptive evidence of immunity and should be considered susceptible unless they have other evidence of measles immunity.

Documented age-appropriate vaccination supersedes the results of subsequent serologic testing. If a person who has two documented doses of measles-containing vaccine is tested serologically and is determined to have negative or equivocal measles results, it is **not** recommended that the person receive an additional dose of vaccine. Such persons should be considered to have presumptive evidence of immunity.

Mumps

Acceptable presumptive evidence of Mumps Immunity (at least one of the following):

- Documentation of age-appropriate vaccination against mumps virus.
- Laboratory evidence of immunity (IgG in serum; equivocal results should be considered negative).
- Laboratory confirmation of disease.
- Born before 1957.

Rubella

Acceptable presumptive evidence of Rubella Immunity (at least one of the following):

- Documentation of 1-dose of live rubella virus-containing vaccine.
- Laboratory evidence of immunity (IgG in serum; equivocal results should be considered negative).
- Laboratory confirmation of disease.
- Born before 1957 (except women of childbearing age who could become pregnant).

Specimen Requirements

Specimen

Serum

Volume

2 mL

Minimum Volume

1 mL

Container

Red-top tube or gel-barrier tube

Reference Range

See individual tests.

Storage Instructions

Refrigerate.

Causes for Rejection

Grossly hemolyzed, lipemic and icteric samples; samples with visible particulate matter or gross bacterial contamination

References

Centers for Disease Control and Prevention. Manual for the Surveillance of Vaccine-Preventable Diseases. CDC web site.

<https://www.cdc.gov/vaccines/pubs/surv-manual/index.html>. Accessed June 2019.

Centers for Disease Control and Prevention. Measles (Rubeola): For Healthcare Professionals. CDC web site.

<https://www.cdc.gov/measles/hcp/>. Accessed June 2019.

McLean HQ, Fiebelkorn AP, Temte JL, Wallace GS; Centers for Disease Control and Prevention. Prevention of measles, rubella, congenital rubella syndrome, and mumps, 2013: summary recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep*. 2013 Jun 14;62(RR-04):1-34. PubMed 23760231

(<http://www.ncbi.nlm.nih.gov/pubmed/23760231>)