

Computational Methods of Optimization

Final Exam(Nov 22, 2024)

Time: 120 minutes

Instructions

- Answer all questions
- Please write your answer in the spaces provided. Answers written outside will not be graded.
- Rough work can be done in designated spaces.
- This is a closed book exam.

Name: _____


Index:

SRNO:

Degree:

Question:	1	2	3	4	5	Total
Points:	10	10	10	5	10	45
Score:						

Rough Work

A large, empty rectangular box with a thin black border, occupying the central portion of the page. It is intended for students to show their rough work or calculations.

In the following, let $\mathbf{I} = [\mathbf{e}_1, \dots, \mathbf{e}_d]$ be a $d \times d$ matrix with \mathbf{e}_j be the j th column. Also $\mathbf{x} = [x_1, x_2, \dots, x_d]^\top \in \mathbb{R}^d$ and $\|\mathbf{x}\| = \sqrt{\mathbf{x}^\top \mathbf{x}}$. Set of real symmetric $d \times d$ matrices will be denoted by \mathcal{S}_d . $[n]$ will denote the set $\{1, 2, \dots, n\}$ Let

$$h(\mathbf{x}) = \frac{1}{2} \mathbf{x}^\top Q \mathbf{x} + b^\top \mathbf{x} + c \quad (\text{QUAD})$$

$\mathbf{x} \in \mathbb{R}^d, Q \in \mathcal{S}_d, b \in \mathbb{R}^d, c \in \mathbb{R}$. For a function $f: \mathbb{R}^d \rightarrow \mathbb{R}, f \in \mathcal{C}_L^1$ the Algorithm DESCENT will refer to the following iteration

$$\mathbf{x}^{(k+1)} = \mathbf{x}^{(k)} + \alpha_k \mathbf{u}^{(k)} \quad (\text{DESCENT})$$

where $\mathbf{u}^{(k)\top} \nabla f(\mathbf{x}^{(k)}) < 0$.

1. True or False.

(a) (1 point) Let $h(\mathbf{x})$ be defined as in (QUAD). For any $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$

$$h(\mathbf{y}) - h(\mathbf{x}) = \nabla h(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) + \frac{1}{2} (\mathbf{y} - \mathbf{x})^\top Q (\mathbf{y} - \mathbf{x})$$

is true only if Q is positive semi-definite or positive definite. **F**

(b) (1 point) Let $\mathbf{u} \in \mathbb{R}^d, \mathbf{u} \neq 0, a > 0$. Consider minimizing $\mathbf{u}^\top \mathbf{x}$ subject to $\|\mathbf{x}\| \leq a$. Optimum value is reached at \mathbf{x}^* where $\|\mathbf{x}^*\| = \frac{a}{2}$ for some \mathbf{u} . **F**

(c) (1 point) Consider running the descent algorithm in (DESCENT) by setting $\mathbf{u}^{(k)} = -\nabla f(\mathbf{x}^{(k)}), \alpha_k = \frac{1}{L}$ with $f(\mathbf{x}) = h(\mathbf{x})$ where $h(\mathbf{x})$ is defined by QUAD. Starting from an arbitrary point $\hat{\mathbf{x}}$ after 5 iterations, we can be assured that

$$h(\mathbf{x}^{(5)}) - h(\mathbf{x}^*) \leq \left(\frac{3}{4}\right)^5 (h(\hat{\mathbf{x}}) - h(\mathbf{x}^*))$$

Assume the eigenvalues of Q are all greater than 1 but less than 7. And assume \mathbf{x}^* is the global minimum. **F**

(d) (1 point) Conjugate Gradient algorithm finds the global unconstrained minimum of h in atmost d iterations **T**.

(e) (1 point) Convergence of Newton's method for minimizing h when the Hessian is positive definite depends on the eigen-values of Q . **F**.

(f) (1 point) Let $\mathbf{z} \notin C$ where $C \subset \mathbb{R}^d$ is a closed convex set. There is only one $\mathbf{x}^* \in C$ such that $\|\mathbf{x}^* - \mathbf{z}\| < \|\mathbf{x} - \mathbf{z}\|$ for all $\mathbf{x} \in C$. **T**

(g) (1 point) For a constrained optimization problem with objective function $f(\mathbf{x})$ it was found that the at \mathbf{x}^* , the optimal point, $\nabla f(\mathbf{x}^*) \neq 0$. Then at least one of the constraints must be active **T**

(h) (1 point) The set of Feasible Descent Directions at $\hat{\mathbf{x}}$ is not empty if $\hat{\mathbf{x}}$ is a KKT point.

(i) (1 point) Gradient projection algorithm in finite number of iterations will find the global minimum for a convex programming problem **F**

(j) (1 point) For a given constrained problem with objective function f it was found that there exists a dual feasible point with dual objective function value -1 and a primal feasible point, $\bar{\mathbf{x}}$ with primal point $f(\bar{\mathbf{x}}) = .5$. Then $f(\bar{\mathbf{x}}) - f(\mathbf{x}^*) \leq 1.5$ **T**

Rough Work

Rough Work

2. Let $f \in \mathcal{C}_L^1$ over \mathbb{R}^d and convex with $L = 5$. For some $\mu > 0$, the function f satisfies the following for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$.

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) + \frac{1}{2} \|\mathbf{x} - \mathbf{y}\|^2$$

Consider minimizing f over \mathbb{R}^d using (DESCENT) using $\mathbf{u}^{(k)} = -\nabla f(\mathbf{x}^{(k)})$, $\alpha_k = \alpha$. Let \mathbf{x}^* be the global minimum

- (a) (2 points) Suppose $\mathbf{x}^{(0)}$ be a point such that $\|\nabla f(\mathbf{x}^{(0)})\| = \sqrt{2}$. Find an upper bound on $\|\mathbf{x}^{(0)} - \mathbf{x}^*\|^2$. Credit will be given to the lowest upper bound.

Solution: Minimizing both sides we obtain

$$f(\mathbf{x}^{(0)}) - f(\mathbf{x}^*) \leq \frac{1}{2} \|\nabla f(\mathbf{x}^{(0)})\|^2 = 1$$

Also since $\nabla f(\mathbf{x}^*) = 0$,

$$f(\mathbf{x}^{(0)}) - f(\mathbf{x}^*) \geq \frac{1}{2} \|\mathbf{x}^{(0)} - \mathbf{x}^*\|^2.$$


- (b) (2 points) At each iteration find a lower bound on the decrease, $f(\mathbf{x}^{(k)}) - f(\mathbf{x}^{(k+1)})$ in terms of $\|\nabla f(\mathbf{x}^{(k)})\|^2$. Find the range of α where decrease is guaranteed?

Solution: The relationship is

$$f(\mathbf{x}^{(k)}) - f(\mathbf{x}^{(k+1)}) \geq \alpha \left(1 - \frac{L}{2} \alpha\right) \|\nabla f(\mathbf{x}^{(k)})\|^2$$

Thus range of $\alpha \in (0, \frac{2}{5})$.

Rough Work

A large, empty rectangular box with a thin black border, occupying the majority of the page below the 'Rough Work' header. It is intended for students to show their calculations or work for the problem on the next page.

- (c) (3 points) Let $\Delta_k = f(\mathbf{x}^{(k)}) - f(\mathbf{x}^*)$. After T iterations derive a bound, more specifically find A and B

$$\log \frac{\Delta_T}{\Delta_0} \leq A + BT?$$

Solution:

$$\Delta_k - \Delta_{k+1} \geq \alpha(1 - \frac{5}{2}\alpha) \|\nabla f(\mathbf{x}^{(k)})\|^2$$

$$\Delta_k \leq \frac{1}{2} \|\nabla f(\mathbf{x}^{(k)})\|^2$$

Define $\beta = 2\alpha(1 - \frac{5}{2}\alpha)$

$$\Delta_{k+1} \leq (1 - \beta)\Delta_k \leq e^{-\beta} \Delta_k$$

$$\Delta_T \leq e^{-T\beta} \Delta_0$$

Thus $A = 0$ and $B = -\beta$

- (d) (3 points) Similarly find A, B such that $\log \|\mathbf{x}^{(T)} - \mathbf{x}^*\|^2 \leq A + BT$. ?

Solution: For all k

$$\frac{1}{2} \|\mathbf{x}^{(k)} - \mathbf{x}^*\|^2 \leq \Delta_k$$

$$\|\mathbf{x}^{(k)} - \mathbf{x}^*\|^2 \leq 2e^{-T\beta} \Delta_0 \leq 2e^{-T\beta} \frac{1}{2} \|\nabla f(\mathbf{x}^{(0)})\|^2$$

Substituting the value of the norm of the gradient we obtain $A = \log 2, B = -\beta$ as defined earlier.

Rough Work

3. Consider $f : \mathbb{R}^3 \rightarrow \mathbb{R} \in \mathcal{C}_L^1$ a convex function and the problem,

$$\begin{aligned} \min_{\mathbf{x} \in \mathbb{R}^3} \quad & f(\mathbf{x}) \\ \text{s.t.} \quad & \frac{1}{3}(x_1x_2 + x_3x_2 + x_1x_2) \leq 1.5, \quad x_1 + ax_2 \geq b, \quad x_2 + x_3 \leq 10, x_3 \geq 0.5 \end{aligned}$$

Suppose you are given a point $\hat{\mathbf{x}} = [1, 1, 1]^\top$ and oracle access to f , namely you have the value $f(\hat{\mathbf{x}}) = 3$, $\nabla f(\hat{\mathbf{x}}) = [1, 10, 0]^\top$.

(a) (5 points) Can you choose $a, b \in \mathbb{R}$ so that $\hat{\mathbf{x}}$ is a KKT point.

Solution: $\hat{\mathbf{x}}$ is feasible. Since $\nabla f(\hat{\mathbf{x}})$ is not zero it means that there must be atleast one active constraint. Only possible candidate is the constraint $x_1 + ax_2 \geq b$. To satisfy KKT conditions

$$\nabla f(\hat{\mathbf{x}}) = \lambda[1, a, 0]^\top$$

Thus $\lambda = 1, \lambda a = 10$. Hence $a = 10$. Since the constraint is active $b = 1 + 10 \cdot 1 = 11$.

(b) (5 points) The global optimal value of f is strictly less than 3. **F**. Justify

Solution: Consider the problem

$$\min_{\mathbf{x} \in \mathbb{R}^3} \quad f(\mathbf{x}), \quad \text{s.t. } x_1 + 10x_2 \geq 11$$

Since $\hat{\mathbf{x}}$ is a KKT point for this problem and hence it is globally optimal. Since $\hat{\mathbf{x}}$ is also feasible for the original problem which has more constraints, it must be the case $f(\hat{\mathbf{x}})$ must be less than or equal to the global optimum of the original problem.

Rough Work

Rough Work

4. (5 points) Let $f, g_1, g_2, g_3 : \mathbb{R}^3 \rightarrow \mathbb{R}$ be \mathcal{C}^1 functions. Consider the following problem.

$$\min_{\mathbf{x}} f(\mathbf{x}) \text{ subject to } g_i(\mathbf{x}) \geq 0, i \in \{1, 2, 3\}$$

However the inequalities maybe in the wrong direction. Given the following partial information about a point $\mathbf{x}^{(0)}$ we need to determine if the inequalities are in the proper directions.

- $\mathbf{x}^{(0)}$ is a KKT point
- $g_1(\mathbf{x}^{(0)}) = 0, g_2(\mathbf{x}^{(0)}) = 0, g_3(\mathbf{x}^{(0)}) = -4$
- $\nabla f(\mathbf{x}^{(0)}) = [4, -12, 4]^\top, \nabla g_1(\mathbf{x}^{(0)}) = [1, -2, 0]^\top, \nabla g_2(\mathbf{x}^{(0)}) = [0, 1, -2]^\top, \nabla g_3(\mathbf{x}^{(0)}) = [-1, 1, 10]^\top$

From the above information can you decide if the inequalities are in the proper direction? If they are not please correct them? Give reasons for your answer.

Solution: The inequalities g_2, g_3 are not in the proper direction. Since $\mathbf{x}^{(0)}$ is a KKT point, it is also feasible. This implies that $g_3(\mathbf{x}) \leq 0$. Hence

$$\nabla f(\mathbf{x}^{(0)}) = \lambda_1 \nabla g_1(\mathbf{x}^{(0)}) + \lambda_2 \nabla g_2(\mathbf{x}^{(0)}) + \lambda_3 \nabla g_3(\mathbf{x}^{(0)})$$

$\lambda_3 = 0$ since g_3 is not active. Assuming that the inequalities are in the correct direction both the λ_i are positive. However the given values indicate $\lambda_1 = 4, \lambda_2 = -2$ is the only possibility. This indicates that g_2 is also in the wrong direction.

Rough Work

Rough Work

5. Let $\mathbf{u} \in \mathbb{R}^d$ such that all coordinates are distinct non-zero values with $a = \min_i u_i, b = \max_i u_i$. Consider the linear programming problem

$$\max_{\mathbf{x} \in \mathbb{R}^d} \mathbf{u}^\top \mathbf{x}, \text{ subject to } \sum_{i=1}^d x_i \leq 1, x_i \geq 0, i \in \{1, \dots, d\}$$

- (a) (5 points) Suppose it is given that the constraint $\sum_{i=1}^d x_i \leq 1$ is not active at optimality. What conclusions can you draw about a, b and the optimal value.

Solution: This is a convex problem and hence a KKT point is sufficient for optimality. Since it is given that the constraint is not active then from the KKT conditions

$$-u_i = \lambda_i \quad \forall i \in [d]$$

where $\lambda_i \geq 0$. This implies that $u_i \leq 0$. Since none of the coordinates are 0 this implies that $a < b < 0$ and furthermore due to KKT conditions $\lambda_i x_i = 0 \implies x_i = 0$. Hence $\mathbf{u}^\top \mathbf{x} = 0$ at optimality.

- (b) (5 points) Repeat the same question supposing that the constraint $\sum_{i=1}^d x_i \leq 1$ is active at optimality.

Solution: The KKT conditions yield the following

$$-u_i = -\lambda + \lambda_i \quad \forall i \in [d]$$

where $\lambda, \lambda_i \geq 0$. All $\lambda_i > 0$ for all i is not possible as it will force all $x_i = 0$ which render the constraint to be inactive. Thus there exists at least one l such that $\lambda_l = 0$. For such l , $u_l = \lambda$ and hence $b > 0$. Also note that $\lambda \geq u_i$ for all i . Thus at KKT point $\lambda = b, \lambda_l = 0, \lambda_i = 0, i \neq l$. At optimality then $x_l = 1$ and $x_i = 0, i \neq l$. Nothing can be said about a .

Rough Work

Rough Work

Rough Work

Rough Work

Rough Work

Rough Work