

1CD-0-3
carcinoma, infiltrating lobular, NOS 8520/3
Site: breast, NOS C50.9 lw
8/17/11

Final Diagnosis

Breast, left, total mastectomy: Infiltrating lobular carcinoma, focal solid variant, Nottingham grade II (of III) [tubules 3/3, nuclei 1/3, mitoses 2/3, Nottingham score 6/9], forming a 11.5 x 7.0 x 6.0 cm mass extensively involving the breast [AJCCpT3]. Lobular carcinoma in situ with focal intraluminal necrosis is present. The tumor does not involve the nipple, overlying skin, or underlying chest wall. All surgical resection margins, after re-excision of the deep margin, are negative for tumor (minimum tumor free margin, 0.8 cm, deep margin).

Lymph node, left axillary sentinel, biopsy: One (of 1) axillary sentinel lymph node is positive for metastatic carcinoma with extensive involvement of the subcapsular sinus [AJCCpN1(sn)]. Immunohistochemical cytokeratin stain was performed on the paraffin embedded sentinel lymph node tissue and confirms the H&E impression. Extranodal extension is not present.

Lymph nodes, left low axillary, dissection: Multiple (17) axillary lymph nodes are negative for metastatic carcinoma.

Seen with Dr.

UUID: 5411AC07-09E4-40C3-844F-2DFAA216D947
TCGA-AR-A2LJ-01A-PR

Redacted



Criteria	Yes	No
Diagnosis Discrepancy		☒
Primary Tumor Site Discrepancy		☒
HIPAA Discrepancy		☒
Prior Malignancy History		☒
Dual/Synchronous Primary Notes		☒
Case is (circle): <u>NOT QUALIFIED</u> / DISQUALIFIED		
Reviewer Initials <u>lw</u> Date Reviewed: <u>8/17/11</u>		