

ICD-0-3

Carcinoma, infiltrating lobular, NOS 8520/3

Path Site: Breast, UOQ C50.4

1/19/11 *lw*

CQCF Site: Breast, NOS C50.9

Diagnosis:

1. Polymorphocellular undifferentiated breast, G II (maximum tumor diameter 2.8 cm) showing infiltrative growth, with marked chronic secondary inflammation, focal invasion of lymph vessels and spread to just below the skin.

The conventional histological findings are consistent with the polymorphocellular variant of invasive lobular breast carcinoma. The nodular tumor infiltration in the axillary tissue described under 6. is to be interpreted as a lymph node metastasis extending beyond the capsule into the surrounding soft tissue. This gives rise to the following summary tumor classification: G II, pT2pN1a(1/14)L1V0R0, negative for estrogen and progesterone receptors, positive for C-erb B-2 oncoprotein.

UUID:E306BE26-0358-4BE4-AEF6-AD42CC0DA020
TCGA-A8-A0A7-01A-PR

Redacted



Criteria	Yes	No
Diagnosis Discrepancy		☒
Primary Tumor Site Discrepancy		☒
HIPAA Discrepancy		☒
Prior Malignancy History		☒
Dual/Synchronous Primary Noted		☒
Cases (Circle):	QUALIFIED	DISQUALIFIED
Revised: <i>lw</i>	Date Reviewed: <i>5/2/11</i>	