


Criteria	Yes	No
Diagnosis Discrepancy		☒
Primary Tumor Site Discrepancy		☒
HPA Discrepancy		☒
Prior Malignancy History		☒
Dual/Synchronous Primary Noted		☒
Case is (circle):	QUALIFIED	DISQUALIFIED
Reviewer Initials: RB	Date Reviewed: 4/25/11	

UUID: 492DDF00-858B-4420-A43F-4FED2DF61C15

TCGA-AR-A255-01A-PR

Redacted



Final Diagnosis

Breast, right, wide local excision: Infiltrating ductal carcinoma, Nottingham grade II (of III) [tubules 3/3, nuclei 3/3, mitoses 1/3; Nottingham score 7/9], forming a mass (1.3 x 1.3 x 1.2 cm) [AJCC pT1c]. Ductal carcinoma in situ, intermediate nuclear grade, comprises <5% of the tumor volume. Angiolymphatic invasion is absent. Biopsy site changes are present. All surgical resection margins are negative for tumor (minimum tumor free margin, 0.7 cm, inferior margin).

Lymph nodes, right axillary sentinel, excision: Multiple (2) right axillary sentinel lymph nodes are negative for metastatic carcinoma [AJCC pN0 (i-) (sn)]. Blue dye is identified in right axillary sentinel lymph node No. 1. Blue dye is not identified in right axillary sentinel lymph node No. 2.

Immunohistochemical cytokeratin stain was performed on the paraffin embedded sentinel lymph node tissue and confirms the H&E impression.

Faxitron done.

ICD-0-3

carcinoma, infiltrating duct, nos 8500/3

Site: breast, nos C50.9

4/25/11