

DIAGNOSIS:

- AXILLARY SENTINEL LYMPH NODE #1, BIOPSY -
 ONE LYMPH NODE WITH METASTATIC CARCINOMA MEASURING 1.5 MM IN GREATEST DIMENSION (1/1).
 B. NO EXTRACAPSULAR EXTENSION IS IDENTIFIED.

PART 2: RIGHT BREAST AT 9 O'CLOCK, SEGMENTAL MASTECTOMY.

- A. MULTIFOCAL INVASIVE DUCTAL CARCINOMA, NOTTINGHAM GRADE 2 (NUCLEAR GRADE 3, TUBULE SCORE 3, MITOTIC ACTIVITY SCORE 1, TOTAL SCORE 7/9).
 B. MAIN TUMOR MASS MEASURES 2.2 CM AND IS ASSOCIATED WITH BIOPSY SITE CHANGES.
 C. A 2.5 MM SATELLITE NODULE OF INVASIVE TUMOR IS PRESENT NEAR THE LATERAL ASPECT OF THE SPECIMEN, AWAY FROM THE MAIN TUMOR MASS.
 D. THIRD TUMOR FOCUS IS SEEN IN PART 3 (see comment).
 E. LYMPHOVASCULAR SPACE INVASION IS IDENTIFIED.
 F. PERINEURAL INVASION IS IDENTIFIED.
 G. TUMOR NODULES CONTAIN CENTRAL FIBROSIS WITH PSEUDO DECIDUALIZED STROMAL HYPERPLASIA (PASH) LIKE STROMAL REACTION.
 H. DUCTAL CARCINOMA IN SITU (DCIS), NUCLEAR GRADE 3 WITH COMEDONECROSIS AND CALCIFICATIONS.
 I. MARGINS ARE NEGATIVE FOR NEOPLASM (INCLUDING BOTH TUMORS IN PART 1); INVASIVE CARCINOMA WAS CLOSEST TO THE ANTERIOR MARGIN (3 MM) AND INTRAVASCULAR TUMOR WAS CLOSEST TO THE SUPERIOR MARGIN (2 MM).
 J. CHANGES CONSISTENT WITH PREVIOUS BIOPSY SITE.
 K. TUMOR PREVIOUSLY REPORTED TO BE POSITIVE FOR ER AND PR, AND NEGATIVE FOR HER-2/NEU.
 L. PATHOLOGIC STAGE: pT2 pN2a pMX.

PART 3: RIGHT AXILLARY CONTENTS, DISSECTION.

- A. SATELLITE NODULE OF INVASIVE DUCTAL CARCINOMA AND DUCTAL CARCINOMA IN SITU, SIMILAR IN APPEARANCE TO TUMOR SEEN IN PART 2 (9.5 MM).
 B. SEVEN LYMPH NODES, POSITIVE FOR METASTATIC ADENOCARCINOMA MEASURING UP TO APPROXIMATELY 6 MM EACH (7/13).
 C. EXTRACAPSULAR EXTENSION IS PRESENT AND MEASURES LESS THAN 1 MM.
 D. EXTENSIVE INTRAVASCULAR TUMOR.

CASE SYNOPSIS:

SYNOPTIC - PRIMARY INVASIVE CARCINOMA OF BREAST

LATERALITY:

Right

PROCEDURE:

Segmental

LOCATION:

Upper outer quadrant

SIZE OF TUMOR:

Lower outer quadrant

MULTICENTRICITY/MULTIFOCALITY OF INVASIVE FOCI:

Maximum dimension invasive component: 2.2 cm

TUMOR AGGREGATE SIZE:

Yes

TUMOR TYPE (invasive component):

Sum of the sizes of multiple invasive tumors: 3.4 cm

NOTTINGHAM SCORE:

Ductal adenocarcinoma, NOS

Nuclear grade: 3

Tubule formation: 3

ANGIOLYMPHATIC INVASION:

Total Nottingham score: 7

DERMAL LYMPHATIC INVASION:

Nottingham grade (1, 2, 3): 2

CALCIFICATION:

Yes

TUMOR TYPE, IN SITU:

Not applicable

Yes, malignant zones

Cribriform

Comedo

SURGICAL MARGINS INVOLVED BY INVASIVE COMPONENT:

No

Distance of invasive tumor to closest margin: 2 mm

SURG MARGINS INVOLVED BY IN SITU COMPONENT:

No

Distance of in situ disease to closest margin: 6 mm

LYMPH NODES POSITIVE:

8

LYMPH NODES EXAMINED:

14

METHOD(S) OF LYMPH NODE EXAMINATION:

H/E stain

SENTINEL NODE METASTASIS:

Yes

SIZE OF NODAL METASTASES:

Diameter of largest lymph node metastasis: 6 mm

LYMPH NODE METASTASIS(-ES) WITH EXTRACAPSULAR EXTENSION:

Yes

T STAGE, PATHOLOGIC:

pT2

N STAGE, PATHOLOGIC:

pN2a

M STAGE, PATHOLOGIC:

pMX

ESTROGEN RECEPTORS:

positive

PROGESTERONE RECEPTORS:

positive

HER2/NEU:

Zero or 1+

100-0-3

carcinoma, infiltrating ductal, NOS 8500/3
 Site: breast, NOS C50.9 3/13/11

UUID: E4CBDC29-63E2-4CF0-B02F-16CA9C69F26E
 TCGA-BH-A0DS-01A-PR

Redacted



Criteria	Yes	No
Diagnosis Discrepancy		2
Primary Tumor Site Discrepancy		2
IPAA Discrepancy		2
Prior Malignancy History		
Local/Synchronous Primary Tumor		
Case is (Little):		
Reviewer Initials:		

DATE REVIEWED: 3/13/11
 REVIEWER: [Signature]