

ICD-0-3

Carcinoma, infiltrating lobular, NOS 8520/3

Pth Site: Breast, UOQ C50.4

1/19/11 *hw*

CQCF Site: Breast, NOS C50.9

Diagnosis:

Skin and fatty tissue with small foci of resorptive inflammatory reactions and also incipient scarring.

But no evidence of carcinoma. A regional R0 status is therefore likely to have been attained.

Diagnosis:

Invasive lobular breast carcinoma (maximum tumor diameter 2.7 cm, malignancy grade 2) with an invasive satellite node and multicentric, in parts confluent LIN formations (grade I). Remaining glandular body fibrolipomatous and with cystic duct ectasia, also small focal areas of adenosis. Submitted tissue samples of cutaneous/subcutaneous tissue histologically unremarkable.

Tumor-free resection margins. Minimum width of dorsal safety margin 0.6 cm.

Tumor classification: pT2m (maximum diameter 2.7 cm), pN0 sn (0/2) (compare xxxxx), MX, R0, G2 (L0, V0).

UUID: E4D30842-383A-40BF-BB91-511A90227655
TCGA-A8-A0AB-01A-PR

Redacted



Criteria	Yes	No
Diagnosis Discrepancy		☒
Primary Tumor Site Discrepancy		☒
HIPAA Discrepancy		☒
Prior Malignancy History		☒
Dual/Synchronous Primary Nodules		☒
Case is (circle):	QUALIFIED	DISQUALIFIED
Reviewed by:	<i>hw</i>	<i>5/31/11</i>