

Surgical Pathology Report

Name: [REDACTED]
DOB: [REDACTED]
Gender: F
MRN: [REDACTED]
Location:
Physician:

Case #: [REDACTED]
Collected:
Received:
Reported:
Copy To:

Pathologic Interpretation:

- A. BREAST, EXCISIONAL/INCISIONAL BIOPSY:**
- INVASIVE MAMMARY CARCINOMA, poorly differentiated, Nottingham grade 3 (3+3+2=8), 2.0 cm in greatest dimension.
- Margins are free of tumor.
- Tumor cells are reported to be negative for ER, PR and Her2 by immunohistochemistry performed on patient's previous biopsy
- See tumor summary.
- B. SENTINEL NODE #1 FS:**
- No carcinoma seen in one lymph node (0/1).
- Keratin immunostain is pending.
- C. SENTINEL NODE #2 FS:**
- Metastatic carcinoma in one lymph node (1/1).
- Largest metastatic deposit, 2.5 cm.
- Perinodal extension is not identified.
- D. SENTINEL NODE #3 FS:**
- No carcinoma seen in one lymph node (0/1).
- Keratin immunostain is pending.
- E. AXILLARY CONTENT RIGHT SIDE:**
- No carcinoma seen in seven lymph nodes (0/7).

ICD-O-3

Carcinoma, infiltrating duct, NOS 8500/3
Site: breast, NOS C50.9 3/12/11 lw

UUID: 2E77693B-FAAD-4CCC-BD4F-3D2470C25143
TCGA-EW-A1PH-01A-PR

Redacted



Tumor Summary

Specimen:

- Partial breast

Procedure:

- Excision without wire-guided localization

Lymph Node Sampling:

- Sentinel lymph node(s)
- Axillary dissection (partial or complete dissection)

Specimen Integrity:

- Multiple designated specimen (eg, main excision and identified margins)

Specimen Size:

- Greatest dimension: 7 cm
- * Additional dimension: 6 x 4 cm.

Laterality:

- Right

Tumor Size: Size of Largest Invasive Carcinoma

- Greatest dimension of largest focus of invasion over 0.1 cm: 2 cm
- * Additional dimensions: 2 x 1.6 cm

Tumor Focality:

- Single focus of invasive carcinoma

Macroscopic and Microscopic Extent of tumor:

- Skin: Skin is not present
- Skeletal muscle: No skeletal muscle present
- Ductal Carcinoma In (DCIS): No DCIS is present
- Lobular Carcinoma In Situ (LCIS): Not identified.

Histologic Type of Invasive Carcinoma:

- Invasive carcinoma

Histologic Grade: Nottingham Histologic Score

- Glandular (Acinar)/Tubular Differentiation: Score 3
- Nuclear Pleomorphism: Score 3

SURGICAL PATHOL Report

Mitotic Count: Score 2

Overall Grade: Grade 3: score of 8

Margins:

- Margins uninvolved by invasive carcinoma
- Distance from closest margin: 1 mm (anterior, posterior, lateral)

*Specify margins:

- Distance from superior margin: 3 mm
- Distance from inferior margin: 2.5 mm
- Distance from anterior margin: 1 mm
- Distance from posterior margin: 1 mm
- Distance from medial margin: 2 mm
- Distance from lateral margin: 1 mm

Treatment Effect: Response to Presurgical (Neoadjuvant) therapy:

- In the Breast: No definite response to presurgical therapy in the invasive carcinoma
- In the Lymph Nodes: No definite response to presurgical therapy in metastatic carcinoma

Lymph-Vascular Invasion: Not identified.

Dermal Lymph-Vascular Invasion: No skin present

Lymph Nodes:

- Number of sentinel lymph node examined: 3
- Total number of lymph nodes examined (sentinel and Nonsentinel): 10
- Number of lymph nodes with macrometastases (>0.2 cm): 1

*Extranodal extension: Not identified

*Method of processing: Hematoxylin and eosin (H&E), one level

Staging:

Tumor: pT1c

Regional Lymph Nodes: pN1

Distant Metastasis: Not applicable

Ancillary Studies:

Estrogen Receptor:

- Performed on another specimen:

Results: No immunoreactive tumor cells present

Her2

Results: No immunoreactive tumor cells present.

- Performed on another specimen:

Results: Negative (Score 0)

Preliminary AJCC Classification (7th Edition) pT1c pN1a Mx/a

(Final AJCC classification pending evaluation of keratin stain on sentinel nodes)

NOTES: Immunohistochemical analysis of the specimen was performed using the following specific reagents (ASRs) validated by our laboratory. These ASRs are clinically useful indicators that do not require FDA approval. These clones are used: ID5 (anti-PR), 636 (anti-PR), A485 (anti-HER2), H-EGFR (anti-EGFR). Immunohistochemical stains are used with formalin or molecular fixed, paraffin embedded tissue. Detection is by Envision Method. The results are read by a pathologist as positive or negative.

As the attending pathologist, I attest that I: (i) Examined the relevant preparation(s) for the specimen(s); and (ii) Rendered the diagnosis(es).

Electronically Signed Out By

Procedures/Addenda

Addendum

Date Ordered:
Date Complete:
Date Reported:

Status: Signed Out

Addendum Diagnosis

- B. SENTINEL NODE #1, FS:
- Immunohistochemistry for keratin is negative for carcinoma.

- D. SENTINEL NODE #3), FS:
 - Immunohistochemistry for keratin is negative for carcinoma.

Intraoperative Consultation

- B. Sentinel node #1 (), FS: No carcinoma seen
 C. Sentinel node #2 FS: Metastatic carcinoma to one lymph node
 D. Sentinel node #3 , FS: No carcinoma seen

MC

MD

Clinical History:

None Provided

Operation Performed

Right breast lumpectomy with sentinel node biopsy and possible node dissection

Pre Operative Diagnosis:

Breast cancer

Specimen(s) Received:

- A: Breast, excisional/incisional biopsy
 B: Sentinel node #1 , FS
 C: Sentinel node #2 (, FS
 D: Sentinel node #3 (, FS
 E: Axillary content right side

Gross Description:

- A. Received in formalin is a 43 gram, 7.0 x 6.0 x 4.0 cm lumpectomy specimen, oriented with one stitch superior, two stitches lateral and three stitches deep. Resection margins are inked as follows: Superior blue, inferior green, medial red, lateral orange, anterior yellow, posterior black. On serial sections through the specimen, there is a tan-white indurated mass with ill-defined borders, measuring 2.0 x 2.0 x 1.6 cm. This lesion is situated less than 1 mm from the anterior resection margin, 1 mm from the lateral resection margin, 7 mm from the medial margin, 4 mm from the deep margin, and 20 cm from superior margin and 2.5 cm from the inferior margin. No other lesions are identified. Sections submitted as follows:
- 1 Lesion in relation to anterior and posterior margin
 - 2 Lateral margin
 - 3 Inferior and medial margin
 - 4 Superior margin
 - 5-8 Lesion submitted in toto
 - 9&10 Additional sections of the specimen with fibrotic area
- B. Received fresh is a segment of tan-grey soft tissue, measuring 1.0 x 1.0 x 0.4 cm. Bisected and submitted in toto in one cassette for frozen.
- C. Received fresh is a segment of tan-brown soft tissue, measuring 0.7 x 0.3 x 0.2 cm. Submitted in toto in one cassette for frozen.
- D. Received fresh is a segment of tan-grey soft tissue, measuring 1.5 x 0.7 x 0.4 cm. Submitted in toto in one cassette for frozen.

SURGICAL PATHOL Report

- E. Received in formalin is a tan-yellow fibroadipose tissue, measuring 11.0 x 6.0 x 2.0 cm. Examination of the specimen reveals multiple possible lymph nodes. Sections submitted as follows:
- 1-3 One lymph node bisected per cassette
 - 4&5 Two lymph nodes in toto per cassette

MD (

Criteria	Yes	NO/NA
Diagnosis Discrepancy		NA/NA
Primary Tumor Site Discrepancy		NA/NA
stPAA Discrepancy		NA/NA
Prior Malignancy History		NA/NA
Local/Synchronous Primary Noted		NA/NA
Case is (circle):	QUALIFIED	DISQUALIFIED
Reviewer Initials	3/12/11	