Statistical Thermodynamics - HW 7 - 2D Ising Model

Prof. Yael Roichman, Nir Sherf

December 2024

Introduction

The Ising model is a mathematical model of ferromagnetism in statistical mechanics. This exercise explores the Ising model in two scenarios: a non-interaction model and a nearest-neighbor interaction model.

In this exercise you will code a simulation using the Monte-Carlo Metropolis-Hastings method for sampling from the canonical ensemble. You are free to write your simulation in any programming language of your choosing (C/Python/Matlab etc..) we recommend that you build upon the demo file given to you in the lecture.

Part 1: Non-Interaction Model

Questions

- 1.1 What is the average energy for a spin in a 2D grid with $\mu = 1, h = 1K_bT$, T = const? What is the average magnetization $\langle s \rangle$? Reach the expression analytically.
- 1.2 Does the simulation reach equilibrium? Show a figure that supports this claim.
- 1.3 How does the simulation differ when we increase the number of spins or iterations?
- 1.4 Plot the distribution of the system for different ratios $\frac{h}{K_bT}$. Explain what you observe in text.
- 1.5 What are the fluctuations in energy and magnetization for each ratio? Show a figure that compares them and explain their meaning from a **physical and statistical** point of view.
- 1.6 Change the initial conditions for the simulation (random grid, spin-up grid, spin-down grid). Does the system reach the same **macroscopic** state?
- 1.7 Does the simulation match the analytical result from 1.1?

Part 2: Nearest-Neighbor Interactions

Questions

- 2.1 Starting from different initial grids (like in 1.6), calculate the average magnetization and average energy for $J = 5K_bT$, assuming h = 0. Does the system reach the same equilibrium state?
- 2.2 Repeat the calculation for high temperatures, $J = \frac{1}{2}K_bT$. What changes in the equilibrium state?
- 2.3 What happens when an external field is added? Must answer in text, Optional: Perform a simulation.
- 2.4 The Ising Model is a classical model in statistical mechanics.
 - (a) What physical phenomena does it describe?
 - (b) Did you observe these phenomena in your simulations?
 - (c) Can you suggest a way to calculate the temperature (ratio between field and temperature) at which this phase change happens?