

Azoreductase producer	Purification steps	References
Shewanella xiamenensis BC01	• Recombinant AzrS collected via centrifugation • Sonication • Centrifugation • His-Trap affinity chromatography • Elution with sodium phosphate buffer made with imidazole & NaCl • Harvested cells were treated with Lysozyme DNase • Lysate clarified by centrifugation	[50]
Bacillus licheniformis	• Supernatant was added with ammonium sulfate (4°C) and EDTA (0.5mM) • Ion-exchange chromatography • Affinity chromatography • Membrane bound azoreductase	[39]
Shewanella sp. Strain IFN4	• Crude extract subjected to (NH₄)₂SO₄ precipitation 25% to 70% saturation • Desalt via desalting column PD-10 • Ion Exchange (Q-Sepharose Fast Flow) • E.coli (BL 21 and Rosetta-gami) transformed with mutated plasmid for improving thermal stability of Azoreductase • Transformed bacteria was harvested via centrifugation	[14]
Halomonas elongate	• Sonication • Centrifugation and supernatant loaded on nickel column • Elution of protein with 100mM imidazole • Buffer exchanged to remove imidazole via Amicon ultra-15 centrifugal filter unit • Culture centrifuged; sonication with lysis buffer • Centrifugation; supernatant loaded on pre-equilibrated Ni-Nitrilotriacetic acid column	[26]
Chromobacterium violaceum	• Elution with elution buffer (Imidazole in Tris-HCl with NaCl) • Dialyzed purified protein for removal of imidazole • SDS-PAGE (analysis of purified protein & size estimation)	[43]
Bacillus wakoensis A01	• Single step affinity chromatographic purification • Mechanistic and crystallographic studies	[34]