

Table 10
Skin sensitization humans

Material	Method	Concentration	Subjects	Results	References
Allyl cinnamate	MAX	4% (2760 µg/cm ²) in petrolatum	22 male volunteers	No reactions	RIFM (1975c)
α-Amylcinnamyl acetate ^a	MAX	8% (5520 µg/cm ²) in petrolatum	25 male and female volunteers	No reactions	RIFM (1975d)
α-Amylcinnamyl alcohol	HRIPT	3% (3543 µg/cm ²) in 3:1 diethyl phthalate:ethanol	31 male and 74 female volunteers	No reactions	RIFM (2004a)
α-Amylcinnamyl alcohol	MAX	8% (5520 µg/cm ²) in petrolatum	25 male volunteers	No reactions	RIFM (1973d)
α-Amylcinnamyl alcohol	Modified Draize	8% in petrolatum 8% in ethyl alcohol	78 volunteers	No (0/78) reactions with 8% in petrolatum 1/78 reactions with 8% in ethyl alcohol	Marzulli and Maibach (1980)
Benzyl cinnamate	HRIPT	4% (4720 µg/cm ²) in 1:3 ethanol: diethyl phthalate	25 male and 76 female volunteers	No reactions	RIFM (2005a)
Benzyl cinnamate	MAX	8% (5520 µg/cm ²) in petrolatum	25 male volunteers	No reactions	RIFM (1972c)
Benzyl cinnamate	MAX	8% (5520 µg/cm ²) in petrolatum	25 male and female volunteers	No reactions	RIFM (1975d)
Butyl cinnamate	MAX	4% (2760 µg/cm ²) in petrolatum	25 male and female volunteers	No reactions	RIFM (1977b)
Cinnamyl acetate	MAX	5% (3450 µg/cm ²) in petrolatum	25 male volunteers	No reactions	RIFM (1972c)
Cinnamyl benzoate	MAX	5% (3450 µg/cm ²) in petrolatum	25 male and female volunteers	No reactions	RIFM (1975d)
Cinnamyl butyrate	MAX	4% (2760 µg/cm ²) in petrolatum	29 male volunteers	No reactions	RIFM (1976b)
Cinnamyl cinnamate	MAX	4% (2760 µg/cm ²) in petrolatum	25 male and female volunteers	No reactions	RIFM (1974d)
Cinnamyl formate	MAX	4% (2760 µg/cm ²) in petrolatum	25 male volunteers	No reactions	RIFM (1973d)
Cinnamyl isobutyrate	MAX	4% (2760 µg/cm ²) in petrolatum	31 male volunteers	No reactions	RIFM (1977c)
Cinnamyl isovalerate	MAX	2% (1380 µg/cm ²) in petrolatum	25 male volunteers	No reactions	RIFM (1973d)
Cinnamyl propionate	MAX	4% (2760 µg/cm ²) in petrolatum	25 male volunteers	No reactions	RIFM (1973d)
Cinnamyl tiglate	MAX	4% (2760 µg/cm ²) in petrolatum	24 male volunteers	1 questionable reaction which was negative at re-test 5 months later	RIFM (1975c)
Ethyl cinnamate	MAX	4% (2760 µg/cm ²) in petrolatum	25 male volunteers	No reactions	RIFM (1973d)
Isoamyl cinnamate	MAX	8% (5520 µg/cm ²) in petrolatum	25 male and female volunteers	No reactions	RIFM (1974d)
Isobutyl cinnamate	MAX	8% (5520 µg/cm ²) in petrolatum	24 male volunteers	No reactions	RIFM (1975c)
Isopropyl cinnamate	MAX	6% (4140 µg/cm ²) in petrolatum	28 male and female volunteers	No reactions	RIFM (1982b)
Linalyl cinnamate	MAX	8% (5520 µg/cm ²) in petrolatum	25 male volunteers	No reactions	RIFM (1973d)
Methyl cinnamate	MAX	2% (1380 µg/cm ²) in petrolatum	25 male volunteers	No reactions	RIFM (1970a)
Methyl cinnamate	MAX	10% (6900 µg/cm ²) in petrolatum	25 male and female volunteers	No reactions	RIFM (1975d)
α-Methylcinnamic alcohol	MAX	2% (1380 µg/cm ²) in petrolatum	25 male and female volunteers	No reactions	RIFM (1974d)
Phenethyl cinnamate	MAX	2% (1380 µg/cm ²) in petrolatum	25 male and female volunteers	No reactions	RIFM (1975d)
Phenethyl cinnamate	MAX	2% (1380 µg/cm ²) in petrolatum	25 male and female volunteers	No reactions	RIFM (1975d)

^a α-Amylcinnamyl acetate is not one of the materials being reviewed, but it is included in this table because it is structurally related.