

Material	Method	Concentration	Species	Results	References
a-Amylcinnamyl alcohol	Open Epicutaneous Test	8% (vehicle not specified by material)	male & female guinea pigs	No reactions (no further data reported)	Klecak (1979, 1985)
α-Amylcinnamyl alcohol	Local Lymph Node Assay	1%, 2.5%, 5%, 10% & 25% in 1:3 ethanol:diethyl phthalate	4 female CBA/Ca/Ola/Hsd mice per group	Negative EC3 > 25% (6250 µg/cm²)	RIFM (2004b)
a-Amylcinnamyl alcohol	Local Lymph Node Assay	7.5%, 15% & 30% in 1:3 diethyl phthalate:ethanol	5 female CBA/J f mice per group	Negative EC3 30% (7500 µg/cm²)	RIFM (2004c)
Benzyl cinnamate	Maximization test	A subirritant concentration in petrolatum	Male & female outbred Himalayan guinea pigs	Sensitization was observed (no further data reported)	Klecak et al. (1977)
Benzyl cinnamate	Freund's Complete Adjuvant Test	A subirritant concentration in petrolatum	Male & female outbred Himalayan guinea pigs	Sensitization was observed (no further data reported)	Klecak et al. (1977)
Benzyl cinnamate	Open Epicutaneous Test	0.3% & 3.0% (vehicle not specified)	Male & female outbred Himalayan guinea pigs	Sensitization was observed (no further data reported)	Klecak et al. (1977)
Benzyl cinnamate	Open Epicutaneous Test	3.0% (vehicle not specified)	6-8 guinea pigs	No reactions (no further data reported)	Klecak (1979)
Benzyl cinnamate	Modified Draize test	0.1% in saline	Male & female outbred Himalayan guinea pigs	Sensitization was observed (no further data reported)	Klecak et al. (1977)
Benzyl cinnamate	Modified Freund's Complete Adjuvant Test	3% and 10% in acetone	10 female Pirbright guinea pigs	1/10 reactions plus 3 questionable reactions at 10% 1/10 reactions plus 2 questionable reactions at 3%	Hausen and Wollenweber (1988) Hausen et al. (1995)
Benzyl cinnamate	Local Lymph Node Assay	2.5%, 5%, 10%, 25% & 50% in 1:3 ethanol: diethyl phthalate	4 female CBA/Ca mice	EC3 18.44% (4610 µg/cm²)	RIFM (2005b)
Cinnamyl cinnamate	Modified Freund's complete adjuvant test	3% & 10% in acetone	Guinea pigs	Weak sensitization was observed at both concentrations (no further details given)	Hausen et al. (1992)
Ethyl cinnamate	Open epicutaneous test	4% (vehicle not specified)	6-8 guinea pigs	No reactions	(1995) Klecak (1979) Klecak (1985)
Methyl cinnamate	Maximization test	a subirritant concentration in petrolatum	Male & female outbred Himalayan guinea pigs	Sensitization effects were observed (no further details given)	Klecak et al. (1977)
Methyl cinnamate	Freund's complete adjuvant test	a subirritant concentration in petrolatum	Male & female outbred Himalayan guinea pigs	Sensitization effects were observed (no further details given)	Klecak et al. (1977)
Methyl cinnamate	Modified Freund's complete adjuvant test	10% in acetone	guinea pigs	No reactions (no further details given)	Hausen et al. (1992)
Methyl cinnamate	Open epicutaneous test	30% (vehicle not specified)	Male & female outbred Himalayan guinea pigs	Sensitization effects were observed (no further details given)	Klecak et al. (1977)
Methyl cinnamate	Open epicutaneous test	2.0% (vehicle not specified)	6-8 guinea pigs	No reactions (no further details given)	Klecak (1979)
Methyl cinnamate	Open epicutaneous test	10% (vehicle not specified)	6-8 guinea pigs	No reactions (no further details given)	Klecak (1985)
Methyl cinnamate	Intradermal sensitization test	0.1% in 5% ethyl alcohol in distilled water	Male albino guinea pigs	No reactions	RIFM (1971b)
Methyl cinnamate	Modified Draize test	0.1% in saline	Male & female outbred Himalayan guinea pigs	Sensitization effects were observed (no further details given)	Klecak et al. (1977)
α-Methyl cinnamic alcohol	Maximization test	75% test material v/v in arachis oil BP	Dunkin Hartley albino guinea pigs	No reactions	RIFM (1997b)