

INCREASED RISK - People who have a history of polyps on prior colonoscopy

Risk category	When to test	Recommended test(s)	Comment
People with small rectal hyperplastic polyps	Same age as those at average risk	Colonoscopy, or other screening options at same intervals as for those at average risk	Those with hyperplastic polyposis syndrome are at increased risk for adenomatous polyps and cancer and should have more intensive follow-up.
People with 1 or 2 small (less than 1 cm) tubular adenomas with low-grade dysplasia	5 to 10 years after the polyps are removed	Colonoscopy	Time between tests should be based on other factors such as prior colonoscopy findings, family history, and patient and doctor preferences.
People with 3 to 10 adenomas, or a large (at least 1 cm) adenoma, or any adenomas with high-grade dysplasia or villous features	3 years after the polyps are removed	Colonoscopy	Adenomas must have been completely removed. If colonoscopy is normal or shows only 1 or 2 small tubular adenomas with low- grade dysplasia, future colonoscopies can be done every 5 years.
People with more than 10 adenomas on a single exam	Within 3 years after the polyps are removed	Colonoscopy	Doctor should consider possible genetic syndrome (such as FAP or Lynch syndrome).
People with sessile adenomas that are removed in pieces	2 to 6 months after adenoma removal	Colonoscopy	If entire adenoma has been removed, further testing should be based on doctor's judgment.