

[Description](#) | [Editorial](#) | [Submissions](#)


42. Trapping Rain Water

Solved 

Hard  Topics  Companies

Given n non-negative integers representing an elevation map where the width of each bar is 1, compute how much water it can trap after raining.

Example 1:



Input: height = [0,1,0,2,1,0,1,3,2,1,2,1]

Output: 6

Explanation: The above elevation map (black section) is represented by array [0,1,0,2,1,0,1,3,2,1,2,1]. In this case, 6 units of rain water (blue section) are being trapped.

Example 2:

Input: height = [4,2,0,3,2,5]

Output: 9


42. Trapping Rain Water

Solved

Hard Topics Companies

Given n non-negative integers representing an elevation map where the width of each bar is 1, compute how much water it can trap after raining.

Example 1:





Input: height = [0,1,0,2,1,0,1,3,2,1,2,1]
Output: 6
Explanation: The above elevation map (black section) is represented by array [0,1,0,2,1,0,1,3,2,1,2,1]. In this case, 6 units of rain water (blue section) are being trapped.

Example 2:

Input: height = [4,2,0,3,2,5]
Output: 9

$[4, 2, 0, 3, 2, 5]$
Total Sum = $(2 \times 5) + 1 + 2 + 1$



$ngl = 5, 5$
 $ngl = 4, 5$
 $gap = 3, 3$
 $min = 3, 4 = 3$
 $1 - 3 = 1$
 $= 1 \times 1 = 1$
 $= 1 \times 1 = 1$

$n \times n$ $7/12$

The Celebrity Problem

Difficulty: Medium Accuracy: 38.33% Submissions: 323K+ Points: 4 Average Time: 30m

A celebrity is a person who is known to all but does not know anyone at a party. A party is being organized by some people. A square matrix `mat[][]` ($n \times n$) is used to represent people at the party such that if an element of **row i and column j is set to 1** it means **i th person knows j th person**. You need to return the **index of the celebrity** in the party, if the celebrity does not exist, return -1.

Note: Follow 0-based indexing.

Examples:

Input: `mat[][] = [[1, 1, 0], [0, 1, 0], [0, 1, 1]]`
Output: 1
Explanation: 0th and 2nd person both know 1st person. Therefore, 1 is the celebrity person.


Input: `mat[][] = [[1, 1], [1, 1]]`
Output: -1
Explanation: Since both the people at the party know each other. Hence none of them is a celebrity person.

Input: `mat[][] = [[1]]`
Output: 0

Constraints:

$1 \leq \text{mat.size}() \leq 1000$
 $0 \leq \text{mat}[i][j] \leq 1$
 $\text{mat}[i][i] == 1$

	0	1	2	3	4	5
0	1	1	1	1	1	1
1	0	1	0	0	0	0
2	1	1	1	0	0	0
3	0	1	1	0	0	0
4	0	1	0	0	1	1
5	1	1	1	1	1	1



if $(\text{arr}[p_1][p_2] == 1) \{$
 $\text{star} = p_2$

$1 \rightarrow 0$
 $p_1 \quad p_2$

Let

$\text{star} = p_1$

2

Description Editorial Submissions

56. Merge Intervals

Solved

Medium Topics Companies

Given an array of `intervals` where `intervals[i] = [starti, endi]`, merge all overlapping intervals, and return an array of the non-overlapping intervals that cover all the intervals in the input.

Example 1:

```
Input: intervals = [[1,3],[2,6],[8,10],[15,18]]
```

```
Output: [[1,6],[8,10],[15,18]]
```

Explanation: Since intervals [1,3] and [2,6] overlap, merge them into [1,6].

Example 2:

```
Input: intervals = [[1,4],[4,5]]
```

Output: `[[1,5]]`

Explanation: Intervals $[1,4]$ and $[4,5]$ are considered overlapping.

Constraints:

- $1 \leq \text{intervals.length} \leq 10^4$
- $\text{intervals}[i].\text{length} = 2$
- $0 \leq \text{start}_i \leq \text{end}_i \leq 10^4$

$$, \left[\begin{matrix} \checkmark \\ 5 \end{matrix} \begin{matrix} 3 \\ 0 \end{matrix} \right], \begin{matrix} 1 \\ \end{matrix} \left[\begin{matrix} 2 \\ 1 \end{matrix} \begin{matrix} 4 \\ 0 \end{matrix} \right], \begin{matrix} 2 \\ \end{matrix} \left[\begin{matrix} 6 \\ 2 \end{matrix} \begin{matrix} 10 \\ 0 \end{matrix} \right], \begin{matrix} 3 \\ \end{matrix} \left[\begin{matrix} 11 \\ 3 \end{matrix} \begin{matrix} 15 \\ 0 \end{matrix} \right], \quad , \begin{matrix} 5 \\ \end{matrix} \left[\begin{matrix} 21 \\ 5 \end{matrix} \begin{matrix} 25 \\ 0 \end{matrix} \right]$$
$$I_1 \Rightarrow [11, 69]$$
$$I_1 \rightarrow [(2), 5]$$
$$212 = 19$$
$$\begin{aligned} & C^{21, 25}) \\ & \Gamma_{11}^{14}) \\ & [1, 10] \end{aligned}$$
$$\checkmark \quad \checkmark$$
$$s_2 < s_1 \rightarrow \checkmark$$

735. Asteroid Collision

Solved 

Medium  Topics  Companies  Hint

We are given an array `asteroids` of ~~integers~~ representing asteroids in a row. The indices of the asteroid in the array represent their relative position in space.

For each asteroid, the absolute value represents its size, and the sign represents its direction (positive meaning right, ~~negative meaning left~~). Each asteroid moves at the same speed.

Find out the state of the asteroids after all collisions. If two asteroids meet, the smaller one will explode. If both are the same size, both will explode. Two asteroids moving in the same direction will never meet.

Example 1:

Input: ~~asteroids = [5,10,-5]~~ asteroids = [5,10,-5]

Output: [5,10]

Explanation: The 10 and -5 collide resulting in 10. The 5 and 10 never collide.

Example 2:

Input: asteroids = [8,-8]

Output: []

Explanation: The 8 and -8 collide exploding each other.

Example 3:

Input: asteroids = [10,2,-5]

Output: [10]

Explanation: The 2 and -5 collide resulting in -5. The 10 and -5 collide resulting in 10.

$[5, 2, -10, 10, -2] \rightarrow [-5, 10]$

$\begin{bmatrix} -10 \\ -10 \end{bmatrix}$

-10