

Recent progress in the monotiling conjecture

Nishant Chandgotia

Tata Institute of Fundamental Research, Bangalore

December

Some notation: Convolutions and tilings

Given two functions $f, g : \mathbb{Z}^d \rightarrow \mathbb{C}$ we write

$$f \star g(x) := \sum_{y \in \mathbb{Z}^d} f(x - y)g(y).$$

Some notation: Convolutions and tilings

Given two functions $f, g : \mathbb{Z}^d \rightarrow \mathbb{C}$ we write

$$f \star g(x) := \sum_{y \in \mathbb{Z}^d} f(x - y)g(y).$$

Given $y \in \mathbb{Z}^d$ note that $(f \star \delta_y)(x) = f(x - y)$.

Some notation: Convolutions and tilings

Given two functions $f, g : \mathbb{Z}^d \rightarrow \mathbb{C}$ we write

$$f \star g(x) := \sum_{y \in \mathbb{Z}^d} f(x - y)g(y).$$

Given $y \in \mathbb{Z}^d$ note that $(f \star \delta_y)(x) = f(x - y)$. Thus convolving by a delta function is just the shift-operation.

Some notation: Convolutions and tilings

Given two functions $f, g : \mathbb{Z}^d \rightarrow \mathbb{C}$ we write

$$f \star g(x) := \sum_{y \in \mathbb{Z}^d} f(x - y)g(y).$$

Given $y \in \mathbb{Z}^d$ note that $(f \star \delta_y)(x) = f(x - y)$. Thus convolving by a delta function is just the shift-operation.

Let $F \subset \mathbb{Z}^d$ be a finite set. We say that F tiles \mathbb{Z}^d at level k if there exists a co-tiler $A \subset \mathbb{Z}^d$ such that

$$1_A \star 1_F = \sum_{y \in A} \delta_y \star 1_F = k.$$

Some notation: Convolutions and tilings

Given two functions $f, g : \mathbb{Z}^d \rightarrow \mathbb{C}$ we write

$$f \star g(x) := \sum_{y \in \mathbb{Z}^d} f(x - y)g(y).$$


Given $y \in \mathbb{Z}^d$ note that $(f \star \delta_y)(x) = f(x - y)$. Thus convolving by a delta function is just the shift-operation.

Let $F \subset \mathbb{Z}^d$ be a finite set. We say that F tiles \mathbb{Z}^d at level k if there exists a co-tiler $A \subset \mathbb{Z}^d$ such that


$$1_A \star 1_F = \sum_{y \in A} \delta_y \star 1_F = k.$$

Tiling at level 1 is just called tiling.

Tiling by a box can is always periodic but can aperiodic in one direction: Aperiodic in vertical direction



Tiling by a box can is always periodic but can aperiodic in one direction: Aperiodic in horizontal direction



Some notation: Convolutions and tilings

$\delta_y \star 1_F$ is just the translate of the set F .

Let $F \subset \mathbb{Z}^d$ be a finite set. We say that F tiles \mathbb{Z}^d at level k if there exists a co-tiler $A \subset \mathbb{Z}^d$ such that

$$1_A \star 1_F = \sum_{y \in A} \delta_y \star 1_F = k.$$

In other words, there are translates of the set F (given by elements of A) which cover the \mathbb{Z}^d lattice k times.

Some notation: Discrete derivatives, periodicities and polynomials

Let $y \in \mathbb{Z}^d \setminus \{0\}$ and consider the discrete derivative in the direction y by

$$\Delta_y(f)(x) = (\delta_y - \delta_0) \star f(x) = f(x - y) - f(x).$$

Some notation: Discrete derivatives, periodicities and polynomials

Let $y \in \mathbb{Z}^d \setminus \{0\}$ and consider the discrete derivative in the direction y by

$$\Delta_y(f)(x) = (\delta_y - \delta_0) \star f(x) = f(x - y) - f(x).$$

f is periodic in the direction y if and only if $\Delta_y(f) = 0$.

Some notation: Discrete derivatives, periodicities and polynomials

Let $y \in \mathbb{Z}^d \setminus \{0\}$ and consider the discrete derivative in the direction y by

$$\Delta_y(f)(x) = (\delta_y - \delta_0) \star f(x) = f(x - y) - f(x).$$

f is periodic in the direction y if and only if $\Delta_y(f) = 0$.

f is **periodic** if and only if its discrete derivative is zero in some direction.

Some notation: Discrete derivatives, periodicities and polynomials

Let $y \in \mathbb{Z}^d \setminus \{0\}$ and consider the discrete derivative in the direction y by

$$\Delta_y(f)(x) = (\delta_y - \delta_0) \star f(x) = f(x - y) - f(x).$$

f is periodic in the direction y if and only if $\Delta_y(f) = 0$.

f is **periodic** if and only if its discrete derivative is zero in some direction.

As with usual derivatives $\Delta_y^k(f) = 0$ if and only if $(f(x + ny); n \in \mathbb{N})$ is a polynomial in n for all $x \in \mathbb{Z}^d$.

Monotiling Conjecture

F tiles \mathbb{Z}^d if there exists A such that

$$1_A \star 1_F = \sum_{y \in A} \delta_y \star 1_F = 1.$$

F tiles \mathbb{Z}^d periodically if there is $A \subset \mathbb{Z}^d$ such that $1_A \star 1_F = 1$ and 1_A is periodic in some direction.

Monotiling Conjecture

F tiles \mathbb{Z}^d if there exists A such that

$$1_A \star 1_F = \sum_{y \in A} \delta_y \star 1_F = 1.$$

F tiles \mathbb{Z}^d periodically if there is $A \subset \mathbb{Z}^d$ such that $1_A \star 1_F = 1$ and 1_A is periodic in some direction.

Question

Suppose F tiles \mathbb{Z}^d . Does F tile \mathbb{Z}^d periodically?

Who cares?

Question

Suppose F tiles \mathbb{Z}^d . Does F tile \mathbb{Z}^d periodically?

- ① Raised by Lagarias and Wang for tilings of \mathbb{R}^d and comes about naturally in connection with a question in harmonic analysis (the Fuglede's conjecture).

Who cares?

Question

Suppose F tiles \mathbb{Z}^d . Does F tile \mathbb{Z}^d periodically?

- ① Raised by Lagarias and Wang for tilings of \mathbb{R}^d and comes about naturally in connection with a question in harmonic analysis (the Fuglede's conjecture).
- ② If there is a tiling whose set of periodicities forms a lattice of full rank then there is an algorithm to decide given a finite set F whether or not it tiles \mathbb{Z}^d (following Hao Wang's ideas from 1961).

Who cares?

Question

Suppose F tiles \mathbb{Z}^d . Does F tile \mathbb{Z}^d periodically?

- ① Raised by Lagarias and Wang for tilings of \mathbb{R}^d and comes about naturally in connection with a question in harmonic analysis (the Fuglede's conjecture).
- ② If there is a tiling whose set of periodicities forms a lattice of full rank then there is an algorithm to decide given a finite set F whether or not it tiles \mathbb{Z}^d (following Hao Wang's ideas from 1961).
- ③ The answer is no if we take multiple tiles instead of one. (This can be done by adapting Robinson's construction of an aperiodic shift of finite type from 1971).

Who cares?

Question

Suppose F tiles \mathbb{Z}^d . Does F tile \mathbb{Z}^d periodically?

- ① Raised by Lagarias and Wang for tilings of \mathbb{R}^d and comes about naturally in connection with a question in harmonic analysis (the Fuglede's conjecture).
- ② If there is a tiling whose set of periodicities forms a lattice of full rank then there is an algorithm to decide given a finite set F whether or not it tiles \mathbb{Z}^d (following Hao Wang's ideas from 1961).
- ③ The answer is no if we take multiple tiles instead of one. (This can be done by adapting Robinson's construction of an aperiodic shift of finite type from 1971).
- ④ Related to Golomb-Welch conjecture in linear codes.

And what did all the care result in?

Question

Suppose F tiles \mathbb{Z}^d . Does F tile \mathbb{Z}^d periodically?

And what did all the care result in?

Question

Suppose F tiles \mathbb{Z}^d . Does F tile \mathbb{Z}^d periodically?

- ① True if $|F|$ is prime (Szegedy).

And what did all the care result in?

Question

Suppose F tiles \mathbb{Z}^d . Does F tile \mathbb{Z}^d periodically?

- ① True if $|F|$ is prime (Szegedy).
- ② True for \mathbb{R} (instead of \mathbb{Z}^d) (Leptin-Müller 1991, also look at Kolountzakis- Lagarias 1996 and Lev-Kolountzakis 2020).

And what did all the care result in?

Question

Suppose F tiles \mathbb{Z}^d . Does F tile \mathbb{Z}^d periodically?

- ① True if $|F|$ is prime (Szegedy).
- ② True for \mathbb{R} (instead of \mathbb{Z}^d) (Leptin-Müller 1991, also look at Kolountzakis- Lagarias 1996 and Lev-Kolountzakis 2020).
- ③ True if F is a topological disc (for $d = 2$, Kenyon - 1992) and convex polytopes (Venkov - 1962). (For \mathbb{R}^d ; $d \geq 2$)

And what did all the care result in?

Question

Suppose F tiles \mathbb{Z}^d . Does F tile \mathbb{Z}^d periodically?

- ① True if $|F|$ is prime (Szegedy).
- ② True for \mathbb{R} (instead of \mathbb{Z}^d) (Leptin-Müller 1991, also look at Kolountzakis- Lagarias 1996 and Lev-Kolountzakis 2020).
- ③ True if F is a topological disc (for $d = 2$, Kenyon - 1992) and convex polytopes (Venkov - 1962). (For \mathbb{R}^d ; $d \geq 2$)
- ④ True for $d = 2$ (Bhattacharya - 2019).

And what did all the care result in?

Question

Suppose F tiles \mathbb{Z}^d . Does F tile \mathbb{Z}^d periodically?

- ① True if $|F|$ is prime (Szegedy).
- ② True for \mathbb{R} (instead of \mathbb{Z}^d) (Leptin-Müller 1991, also look at Kolountzakis- Lagarias 1996 and Lev-Kolountzakis 2020).
- ③ True if F is a topological disc (for $d = 2$, Kenyon - 1992) and convex polytopes (Venkov - 1962). (For \mathbb{R}^d ; $d \geq 2$)
- ④ True for $d = 2$ (Bhattacharya - 2019).
- ⑤ Parallel developments took place with regard to a similar conjecture (called the Nivat's conjecture) with Kari-Szabados and Kari-Moutot.

And what did all the care result in?

Question

Suppose F tiles \mathbb{Z}^d . Does F tile \mathbb{Z}^d periodically?

- ① True if $|F|$ is prime (Szegedy).
- ② True for \mathbb{R} (instead of \mathbb{Z}^d) (Leptin-Müller 1991, also look at Kolountzakis- Lagarias 1996 and Lev-Kolountzakis 2020).
- ③ True if F is a topological disc (for $d = 2$, Kenyon - 1992) and convex polytopes (Venkov - 1962). (For \mathbb{R}^d ; $d \geq 2$)
- ④ True for $d = 2$ (Bhattacharya - 2019).
- ⑤ Parallel developments took place with regard to a similar conjecture (called the Nivat's conjecture) with Kari-Szabados and Kari-Moutot.
- ⑥ Wide open for higher dimensions.

What did Bhattacharya prove?

Before we understand this, we need another definition.

What did Bhattacharya prove?


Before we understand this, we need another definition.

A set $A \subset \mathbb{Z}^d$ is called **weakly-periodic** if there is a lattice $\Lambda \subset \mathbb{Z}^d$ such that for all its cosets $x + \Lambda$,


$$(x + \Lambda) \cap A$$

is periodic.

When all tilings are periodic?




When some of them are weakly periodic but not periodic



Shape F

The even sublattice can be tiled so that it is aperiodic in the horizontal direction


When some of them are weakly periodic but not periodic



Shape F

The even sublattice can be tiled so that it is aperiodic in the horizontal direction while the odd sublattice can be tiled so that it is aperiodic in the vertical direction.

When some of them are weakly periodic but not periodic



Shape F

The even sublattice can be tiled so that it is aperiodic in the horizontal direction while the odd sublattice can be tiled so that it is aperiodic in the vertical direction.

The resulting configuration will be weakly periodic but not periodic.

What did Bhattacharya prove?

Note that the set of A such that $1_A \star 1_F = 1$ is invariant under translations by \mathbb{Z}^d . In fact it can be considered as a compact subset of the $\{0, 1\}^{\mathbb{Z}^2}$ and has the structure of a shift of finite type.

What did Bhattacharya prove?

Note that the set of A such that $1_A \star 1_F = 1$ is invariant under translations by \mathbb{Z}^d . In fact it can be considered as a compact subset of the $\{0, 1\}^{\mathbb{Z}^2}$ and has the structure of a shift of finite type.

For $d = 2$, Bhattacharya said that with respect to any invariant measure, every co-tiler A is weakly periodic.

What did Bhattacharya prove?

Note that the set of A such that $1_A \star 1_F = 1$ is invariant under translations by \mathbb{Z}^d . In fact it can be considered as a compact subset of the $\{0, 1\}^{\mathbb{Z}^2}$ and has the structure of a shift of finite type.

For $d = 2$, Bhattacharya said that with respect to any invariant measure, every co-tiler A is weakly periodic.

From this result he used some standard ideas to show that a periodic tiling exists.

And what have Greenfeld and Tao done?

Theorem

Let $F \subset \mathbb{Z}^2$ be a finite set. Let $A \subset \mathbb{Z}^2$ be such that $1_A \star 1_F = 1$. Then A is weakly periodic.

And what have Greenfeld and Tao done?

Theorem

Let $F \subset \mathbb{Z}^2$ be a finite set. Let $A \subset \mathbb{Z}^2$ be such that $1_A \star 1_F = 1$. Then A is weakly periodic.

Further, the results are effective.

And what have Greenfeld and Tao done?

Theorem

Let $F \subset \mathbb{Z}^2$ be a finite set. Let $A \subset \mathbb{Z}^2$ be such that $1_A \star 1_F = 1$. Then A is weakly periodic.

Further, the results are effective.

Theorem

There is $F \subset \mathbb{Z}^2$ be a finite set for which there is a level-4 co-tiler, meaning,

$$1_F \star 1_A = 4$$

but A is not weakly periodic.

And what have Greenfeld and Tao done?

Theorem

Let $F \subset \mathbb{Z}^2$ be a finite set. Let $A \subset \mathbb{Z}^2$ be such that $1_A \star 1_F = 1$. Then A is weakly periodic.

Further, the results are effective.

Theorem

There is $F \subset \mathbb{Z}^2$ be a finite set for which there is a level-4 co-tiler, meaning,


$$1_F \star 1_A = 4$$

but A is not weakly periodic.

Fundamentally the ideas being used are not very different from that by Bhattacharya but the usage is much more elegant and clearer.

Level-4 tiling which is not weakly periodic

$$1_F := 1_{(0,0),(1,0)} \star 1_{(0,0),(0,2)} \star 1_{(0,0),(2,-2)}.$$



Level-4 tiling which is not weakly periodic

- 1 Consider a group $K := \mathbb{R}/\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ and $\alpha \in \mathbb{R}/\mathbb{Z}$ irrational.

Level-4 tiling which is not weakly periodic

- 1 Consider a group $K := \mathbb{R}/\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ and $\alpha \in \mathbb{R}/\mathbb{Z}$ irrational.
- 2 Consider the rotation on K given by

$$(m_1, m_2)(\beta, x, y) = (\beta + (m_1 + m_2)\alpha, x + m_1, y + m_2).$$

Level-4 tiling which is not weakly periodic

- 1 Consider a group $K := \mathbb{R}/\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ and $\alpha \in \mathbb{R}/\mathbb{Z}$ irrational.
- 2 Consider the rotation on K given by

$$(m_1, m_2)(\beta, x, y) = (\beta + (m_1 + m_2)\alpha, x + m_1, y + m_2).$$

- 3 Find a set $B \subset K$ such that

$$1_F \star 1_B := 4.$$

Level-4 tiling which is not weakly periodic

- 1 Consider a group $K := \mathbb{R}/\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ and $\alpha \in \mathbb{R}/\mathbb{Z}$ irrational.
- 2 Consider the rotation on K given by

$$(m_1, m_2)(\beta, x, y) = (\beta + (m_1 + m_2)\alpha, x + m_1, y + m_2).$$

- 3 Find a set $B \subset K$ such that

$$1_F \star 1_B := 4.$$

- 4 The visit times to B , that is,

$$A := \{(m_1, m_2) : (m_1, m_2)(0, 0, 0) \in B\}$$

has the property

$$1_F \star 1_A = 4.$$

Level-4 tiling which is not weakly periodic

- 1 Consider a group $K := \mathbb{R}/\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ and $\alpha \in \mathbb{R}/\mathbb{Z}$ irrational.
- 2 Consider the rotation on K given by

$$(m_1, m_2)(\beta, x, y) = (\beta + (m_1 + m_2)\alpha, x + m_1, y + m_2).$$

- 3 Find a set $B \subset K$ such that

$$1_F \star 1_B := 4.$$

- 4 The visit times to B , that is,


$$A := \{(m_1, m_2) : (m_1, m_2)(0, 0, 0) \in B\}$$

has the property


$$1_F \star 1_A = 4.$$

- 5 Because α is irrational, for most choices of B , the level 4 co-tiler A is not weakly periodic.

The set B is the union of



and



Greenfeld and Tao ask whether there exists a periodic tiling exists
for tilings of all level?

Greenfeld and Tao ask whether there exists a periodic tiling exists for tilings of all level?

I conjecture that all tilings of a certain level of a set F arise as visit times of a rotation of a compact group.

Greenfeld and Tao ask whether there exists a periodic tiling exists for tilings of all level?

I conjecture that all tilings of a certain level of a set F arise as visit times of a rotation of a compact group.

While it is false as stated, a precise conjecture close to it can be formulated along these lines (this is done at the end).

The main theorem

A set $A \subset \mathbb{Z}^d$ is called **weakly-periodic** if there is a lattice $\Lambda \subset \mathbb{Z}^d$ such that for all its cosets $x + \Lambda$,

$$(x + \Lambda) \cap A$$

is periodic.

Theorem

Let $F \subset \mathbb{Z}^2$ be a finite set. Let $A \subset \mathbb{Z}^2$ be such that $1_A \star 1_F = 1$. Then A is weakly periodic.

The main theorem

A set $A \subset \mathbb{Z}^d$ is called **weakly-periodic** if there is a lattice $\Lambda \subset \mathbb{Z}^d$ such that for all its cosets $x + \Lambda$,

$$(x + \Lambda) \cap A$$

is periodic.

Theorem

Let $F \subset \mathbb{Z}^2$ be a finite set. Let $A \subset \mathbb{Z}^2$ be such that $1_A \star 1_F = 1$. Then A is weakly periodic.

We will concentrate on $d = 2$ and level 1 tilings though a lot of ideas go through to higher dimensions and higher level tilings.

Dilation lemma (starting point for Szegedy, Bhattacharya, Kari-Szabados)

Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

If q is the product of all primes less than or equal to $|F|$ then for $r \equiv 1 \pmod q$ we have

$$1_{rF} \star 1_A = 1.$$

Dilation lemma (starting point for Szegedy, Bhattacharya, Kari-Szabados)

Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

If q is the product of all primes less than or equal to $|F|$ then for $r \equiv 1 \pmod q$ we have

$$1_{rF} \star 1_A = 1.$$

If F tiles \mathbb{Z}^d by a set A then rF also tiles \mathbb{Z}^d by the same set A for chosen numbers r .

Dilation lemma

Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

If q is the product of all primes less than or equal to $|F|$ then for $r \equiv 1 \pmod q$ we have

$$1_{rF} \star 1_A = 1.$$

Dilation lemma

Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

If q is the product of all primes less than or equal to $|F|$ then for $r \equiv 1 \pmod q$ we have

$$1_{rF} \star 1_A = 1.$$

Let p be a prime greater than $|F|$. We have $(1_F)^p \star 1_A = |F|^{p-1}$.

Dilation lemma

Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

If q is the product of all primes less than or equal to $|F|$ then for $r \equiv 1 \pmod q$ we have

$$1_{rF} \star 1_A = 1.$$

Let p be a prime greater than $|F|$. We have $(1_F)^p \star 1_A = |F|^{p-1}$.

Going modulo p kills all the multinomial terms. Further Fermat's little theorem gives us

$$1_{pF} \star 1_A \pmod p = 1 \pmod p.$$

Dilation lemma

Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

If q is the product of all primes less than or equal to $|F|$ then for $r \equiv 1 \pmod q$ we have

$$1_{rF} \star 1_A = 1.$$

Let p be a prime greater than $|F|$. We have $(1_F)^p \star 1_A = |F|^{p-1}$.

Going modulo p kills all the multinomial terms. Further Fermat's little theorem gives us

$$1_{pF} \star 1_A \pmod p = 1 \pmod p.$$

Finally $1_{pF} \star 1_A$ is bounded by $|pF| = |F|$ and hence $1_{pF} \star 1_A = 1$. The same fact for r now follows since the prime factors of r are greater than $|F|$.

We will always assume that $0 \in F$.

Periodic decomposition

We assume $0 \in F$. Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

If q is the product of all primes less than or equal to $|F|$ then

$$1_{(tq+1)F} * 1_A = 1.$$

Periodic decomposition

We assume $0 \in F$. Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

If q is the product of all primes less than or equal to $|F|$ then

$$1_{(tq+1)F} \star 1_A = 1.$$

This implies that

$$1_A = 1 - \sum_{f \in F \setminus \{0\}} \delta_{(tq+1)f} \star 1_A = 1 - \sum_{f \in F \setminus \{0\}} \frac{1}{N} \sum_{t=1}^N \delta_{(tq+1)f} \star 1_A.$$

Periodic decomposition

We assume $0 \in F$. Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

If q is the product of all primes less than or equal to $|F|$ then

$$1_{(tq+1)F} \star 1_A = 1.$$

This implies that

$$1_A = 1 - \sum_{f \in F \setminus \{0\}} \delta_{(tq+1)f} \star 1_A = 1 - \sum_{f \in F \setminus \{0\}} \frac{1}{N} \sum_{t=1}^N \delta_{(tq+1)f} \star 1_A.$$

By taking $N \rightarrow \infty$ and choosing a convergent subsequence we have

$$1_A = 1 - \sum_{f \in F \setminus \{0\}} \phi_f$$

Periodic decomposition

We assume $0 \in F$. Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

If q is the product of all primes less than or equal to $|F|$ then

$$1_{(tq+1)F} * 1_A = 1.$$

This implies that

$$1_A = 1 - \sum_{f \in F \setminus \{0\}} \delta_{(tq+1)f} \star 1_A = 1 - \sum_{f \in F \setminus \{0\}} \frac{1}{N} \sum_{t=1}^N \delta_{(tq+1)f} \star 1_A.$$

By taking $N \rightarrow \infty$ and choosing a convergent subsequence we have

$$1_A = 1 - \sum_{f \in F \setminus \{0\}} \phi_f$$

where $\phi_f : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in the qf direction.

A similar step is taken by Kari-Szabados by taking an ultrafilter limit and by Bhattacharya by considering an ergodic invariant measure to start with. The dilation lemma also finds its presence in the work of Szegedy.

Periodic decomposition

Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

We had

$$1_A = 1 - \sum_{f \in F \setminus \{0\}} \phi_f.$$

where $\phi_f : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in the f direction.

Periodic decomposition

Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

We had

$$1_A = 1 - \sum_{f \in F \setminus \{0\}} \phi_f.$$

where $\phi_f : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in the f direction.

In fact $0 \leq \sum_{f \in F \setminus \{0\}} \phi_f \leq 1$.

Periodic decomposition

Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

We had

$$1_A = 1 - \sum_{f \in F \setminus \{0\}} \phi_f.$$

where $\phi_f : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in the f direction.

In fact $0 \leq \sum_{f \in F \setminus \{0\}} \phi_f \leq 1$. By grouping terms which are linearly dependent we have that

$$1_A = 1 - \sum_{j=1}^m \phi_j$$

where $\phi_j : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in a direction h_j .

Periodic decomposition

Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

We had

$$1_A = 1 - \sum_{f \in F \setminus \{0\}} \phi_f.$$

where $\phi_f : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in the f direction.

In fact $0 \leq \sum_{f \in F \setminus \{0\}} \phi_f \leq 1$. By grouping terms which are linearly dependent we have that

$$1_A = 1 - \sum_{j=1}^m \phi_j$$

where $\phi_j : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in a direction h_j . Here h_j 's are linearly independent and multiples of vectors in F .

Periodic decomposition

Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

We had

$$1_A = 1 - \sum_{f \in F \setminus \{0\}} \phi_f.$$

where $\phi_f : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in the f direction.

In fact $0 \leq \sum_{f \in F \setminus \{0\}} \phi_f \leq 1$. By grouping terms which are linearly dependent we have that

$$1_A = 1 - \sum_{j=1}^m \phi_j$$

where $\phi_j : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in a direction h_j . Here h_j 's are linearly independent and multiples of vectors in F . Note that we have written 1_A as a sum of periodic terms.

Periodic decomposition

Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

We had

$$1_A = 1 - \sum_{f \in F \setminus \{0\}} \phi_f.$$

where $\phi_f : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in the f direction.

In fact $0 \leq \sum_{f \in F \setminus \{0\}} \phi_f \leq 1$. By grouping terms which are linearly dependent we have that

$$1_A = 1 - \sum_{j=1}^m \phi_j$$

where $\phi_j : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in a direction h_j . Here h_j 's are linearly independent and multiples of vectors in F . Note that we have written 1_A as a sum of periodic terms. We want to write it as a sum of 0, 1 valued periodic terms.

Periodic decomposition

Let $F \subset \mathbb{Z}^2$ be a finite set and $A \subset \mathbb{Z}^2$ be such that

$$1_F \star 1_A = 1.$$

Then

$$1_A = 1 - \sum_{j=1}^m \phi_j$$

where each $\phi_j : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in a direction h_j ; the h_j are linearly independent.

Polynomials everywhere

$$1_A = 1 - \sum_{j=1}^m \phi_j$$

where each $\phi_j : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in a direction h_j ; the h_j are linearly independent.

Polynomials everywhere

$$1_A = 1 - \sum_{j=1}^m \phi_j$$

where each $\phi_j : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in a direction h_j ; the h_j are linearly independent.

Choose e such that it is linearly independent of all the h_i and

$$e \in \langle h_i, h_j \rangle$$

for all i, j .

Polynomials everywhere

$$1_A = 1 - \sum_{j=1}^m \phi_j$$

where each $\phi_j : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in a direction h_j ; the h_j are linearly independent.

Choose e such that it is linearly independent of all the h_i and

$$e \in \langle h_i, h_j \rangle$$

for all i, j .

Consider the function $P_{x,j} : \mathbb{Z} \rightarrow [0, 1]$ given by

$$P_{x,j}(n) := \phi_j(x + ne).$$

Some intuition about the $P_{x,j}$

$$1_A = 1 - \sum_{j=1}^m \phi_j.$$

$\phi_j : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in linearly independent directions h_j . e is linearly independent of all h_i and in $\langle h_i, h_j \rangle$ for all i, j . Let

$$P_{x,j}(n) := \phi_j(x + ne).$$

These are bounded. If these function $n \rightarrow P_{x,j}(n)$ were polynomials then these sequences would be either periodic or equidistributed.

Some intuition about the $P_{x,j}$

$$1_A = 1 - \sum_{j=1}^m \phi_j.$$

$\phi_j : \mathbb{Z}^2 \rightarrow [0, 1]$ is periodic in linearly independent directions h_j . e is linearly independent of all h_i and in $\langle h_i, h_j \rangle$ for all i, j . Let

$$P_{x,j}(n) := \phi_j(x + ne).$$

These are bounded. If these function $n \rightarrow P_{x,j}(n)$ were polynomials then these sequences would be either periodic or equidistributed.

Let us see why they are polynomials.

$P_{x,j}$ is a nice polynomial

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is periodic in the direction h_j .

$$P_{x,j}(n) := \phi_j(x + ne)$$

where e is linearly independent of all h_i and in $\langle h_i, h_j \rangle$ for all i, j .

$P_{x,j}$ is a nice polynomial

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is periodic in the direction h_j .

$$P_{x,j}(n) := \phi_j(x + ne)$$

where e is linearly independent of all h_i and in $\langle h_i, h_j \rangle$ for all i, j .

Lemma

Let $x \in \mathbb{Z}^2$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$.
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m - 2$.
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

$P_{x,j}$ is a nice polynomial

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is periodic in the direction h_j .

$$P_{x,j}(n) := \phi_j(x + ne)$$

where e is linearly independent of all h_i and in $\langle h_i, h_j \rangle$ for all i, j .

Lemma

Let $x \in \mathbb{Z}^2$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$.
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m - 2$.
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

(1) follows immediately from \star .

$P_{x,j}$ is a nice polynomial

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is periodic in the direction h_j .

$$P_{x,j}(n) := \phi_j(x + ne)$$

where e is linearly independent of all h_i and in $\langle h_i, h_j \rangle$ for all i, j .

Lemma

Let $x \in \mathbb{Z}^2$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$.
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m - 2$.
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

(1) follows immediately from \star . For (2) we need to show $\Delta_e^{m-1} \phi_j = 0 \bmod 1$.

$$\Delta_e^{m-1} \phi_j = 0 \bmod 1$$

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle h_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne)$$

where e is linearly independent of all h_i and in $\langle h_i, h_j \rangle$ for all i, j .

We have $e = a_i h_i + b_j h_j$.

$$\Delta_e^{m-1} \phi_j = 0 \bmod 1$$

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle h_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne)$$

where e is linearly independent of all h_i and in $\langle h_i, h_j \rangle$ for all i, j .

We have $e = a_i h_i + b_j h_j$. Since ϕ_i is $\langle h_i \rangle$ periodic we have

$$0 = \Delta_{a_i h_i} \phi_i = \Delta_{e - b_j h_j} \phi_i.$$

$$\Delta_e^{m-1} \phi_j = 0 \bmod 1$$

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle h_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne)$$

where e is linearly independent of all h_i and in $\langle h_i, h_j \rangle$ for all i, j .

We have $e = a_i h_i + b_j h_j$. Since ϕ_i is $\langle h_i \rangle$ periodic we have

$$0 = \Delta_{a_i h_i} \phi_i = \Delta_{e - b_j h_j} \phi_i.$$

But since we have ϕ_j is $\langle h_j \rangle$ -periodic we have

$$\Delta_e \phi_j = \Delta_{e - b_j h_j} \phi_j.$$

$$\Delta_e^{m-1} \phi_j = 0 \bmod 1$$

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle h_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne)$$

where e is linearly independent of all h_i and in $\langle h_i, h_j \rangle$ for all i, j .

We have $e = a_i h_i + b_j h_j$. Since ϕ_i is $\langle h_i \rangle$ periodic we have

$$0 = \Delta_{a_i h_i} \phi_i = \Delta_{e - b_j h_j} \phi_i.$$

But since we have ϕ_j is $\langle h_j \rangle$ -periodic we have

$$\Delta_e \phi_j = \Delta_{e - b_j h_j} \phi_j.$$

Thus from \star we have that

$$\Delta_e^{m-1} \phi_j \bmod 1 = \left(\prod_{i \neq j} \Delta_{e - b_i h_i} \right) (\phi_j) \bmod 1 = \left(\prod_{i \neq j} \Delta_{e - b_i h_i} \right) \left(\sum_{t=1}^m \phi_t \right) \bmod 1 = 0.$$

$P_{x,j}$ is a nice polynomial

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle h_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne)$$

where e is linearly independent of all h_i and in $\langle h_i, h_j \rangle$ for all i, j .

Lemma

Let $x \in \mathbb{Z}^2$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$. ✓
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m - 2$. ✓
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

$P_{x,j}$ is a nice polynomial

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle h_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne)$$

where e is linearly independent of all h_i and in $\langle h_i, h_j \rangle$ for all i, j .

Lemma

Let $x \in \mathbb{Z}^2$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$. ✓
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m - 2$. ✓
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

We want to bound $P_{x,i}(n_1) + P_{x,j}(n_2) = \phi_i(x + n_1 e) + \phi_j(x + n_2 e)$ for part (3).

Bounding $P_{x,i}(n_1) + P_{x,j}(n_2)$

$$P_{x,i}(n_1) + P_{x,j}(n_2) = \phi_i(x + n_1 e) + \phi_j(x + n_2 e).$$

Bounding $P_{x,i}(n_1) + P_{x,j}(n_2)$

$$P_{x,i}(n_1) + P_{x,j}(n_2) = \phi_i(x + n_1 e) + \phi_j(x + n_2 e).$$

ϕ_i is periodic in h_i direction and ϕ_j is periodic in h_j direction.

Bounding $P_{x,i}(n_1) + P_{x,j}(n_2)$

$$P_{x,i}(n_1) + P_{x,j}(n_2) = \phi_i(x + n_1 e) + \phi_j(x + n_2 e).$$

ϕ_i is periodic in h_i direction and ϕ_j is periodic in h_j direction.

Since $1_A = 1 - \sum_{j=1}^m \phi_j$ it follows that

$$\phi_i(\tilde{x}) + \phi_j(\tilde{x}) \leq 1$$

for all \tilde{x} .

Bounding $P_{x,i}(n_1) + P_{x,j}(n_2)$

$$P_{x,i}(n_1) + P_{x,j}(n_2) = \phi_i(x + n_1 e) + \phi_j(x + n_2 e).$$

ϕ_i is periodic in h_i direction and ϕ_j is periodic in h_j direction.

Since $1_A = 1 - \sum_{j=1}^m \phi_j$ it follows that


$$\phi_i(\tilde{x}) + \phi_j(\tilde{x}) \leq 1$$

for all \tilde{x} .


So it is enough to show that there is some \tilde{x} such that

$x + n_1 e - \tilde{x}$ is a multiple of h_i and $x + n_2 e - \tilde{x}$ is a multiple of h_j .

So it is enough to show that there is some \tilde{x} such that $x + n_1 e - \tilde{x}$ is a multiple of h_i and $x + n_2 e - \tilde{x}$ is a multiple of h_j .



So it is enough to show that there is some \tilde{x} such that $x + n_1 e - \tilde{x}$ is a multiple of h_i and $x + n_2 e - \tilde{x}$ is a multiple of h_j .



This follows because e is linearly independent of h_i, h_j and is in $\langle h_i, h_j \rangle$.

A set $A \subset \mathbb{Z}^d$ is called **weakly-periodic** if there is a lattice $\Lambda \subset \mathbb{Z}^d$ such that for all its cosets $x + \Lambda$,

$$(x + \Lambda) \cap A$$

is periodic.

Theorem

Let $F \subset \mathbb{Z}^2$ be a finite set. Let $A \subset \mathbb{Z}^2$ be such that $1_A \star 1_F = 1$. Then A is weakly periodic.

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle h_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne) \text{ where } e \text{ satisfies}$$

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle h_j \rangle$ periodic.

$P_{x,j}(n) := \phi_j(x + ne)$ where e satisfies

- ① e is linearly independent of all h_i .
- ② Choose N such that $e \in N\mathbb{Z}^2 \subset \langle h_i, h_j \rangle$ for all i, j .
- ③ M is chosen such that $M\mathbb{Z}^2 \subset \langle e, h_j \rangle$ for all j .

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle h_j \rangle$ periodic.

$P_{x,j}(n) := \phi_j(x + ne)$ where e satisfies

- ① e is linearly independent of all h_i .
- ② Choose N such that $e \in N\mathbb{Z}^2 \subset \langle h_i, h_j \rangle$ for all i, j .
- ③ M is chosen such that $M\mathbb{Z}^2 \subset \langle e, h_j \rangle$ for all j .

Let $\Lambda = \langle e, NMh_1 \rangle$. We will prove $A \cap (x + \Lambda)$ is periodic.

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle h_j \rangle$ periodic.

$P_{x,j}(n) := \phi_j(x + ne)$ where e satisfies

- ① e is linearly independent of all h_i .
- ② Choose N such that $e \in N\mathbb{Z}^2 \subset \langle h_i, h_j \rangle$ for all i, j .
- ③ M is chosen such that $M\mathbb{Z}^2 \subset \langle e, h_j \rangle$ for all j .

Let $\Lambda = \langle e, NMh_1 \rangle$. We will prove $A \cap (x + \Lambda)$ is periodic. Rewriting \star we get that

$$1_A(x + se + tNMh_1) = 1 - \sum_{j=1}^m \phi_j(x + se + tNMh_1).$$

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle h_j \rangle$ periodic.

$P_{x,j}(n) := \phi_j(x + ne)$ where e satisfies

- ① e is linearly independent of all h_i .
- ② Choose N such that $e \in N\mathbb{Z}^2 \subset \langle h_i, h_j \rangle$ for all i, j .
- ③ M is chosen such that $M\mathbb{Z}^2 \subset \langle e, h_j \rangle$ for all j .

Let $\Lambda = \langle e, NMh_1 \rangle$. We will prove $A \cap (x + \Lambda)$ is periodic. Rewriting \star we get that

$$1_A(x + se + tNMh_1) = 1 - \sum_{j=1}^m \phi_j(x + se + tNMh_1).$$

But $NMh_1 = a_i h_i + b_i e$.

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle h_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne) \text{ where } e \text{ satisfies}$$

- ① e is linearly independent of all h_i .
- ② Choose N such that $e \in N\mathbb{Z}^2 \subset \langle h_i, h_j \rangle$ for all i, j .
- ③ M is chosen such that $M\mathbb{Z}^2 \subset \langle e, h_j \rangle$ for all j .

Let $\Lambda = \langle e, NMh_1 \rangle$. We will prove $A \cap (x + \Lambda)$ is periodic. Rewriting \star we get that

$$1_A(x + se + tNMh_1) = 1 - \sum_{j=1}^m \phi_j(x + se + tNMh_1).$$

But $NMh_1 = a_i h_i + b_i e$. Hence

$$\begin{aligned} \phi_i(x + se + tNMh_1) &= \phi_i(x + (s + b_i t)e + ta_i h_i) \\ &= \phi_i(x + (s + b_i t)e) \\ &= P_{x,i}(s + b_i t). \end{aligned}$$

$$\text{Thus } 1_A(x + se + tNMh_1) = 1 - \sum_{j=1}^m P_{x,j}(s + b_j t).$$

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle qh_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne).$$

Properties of $P_{x,j}$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$.
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m - 2$.
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

We had $NMh_1 = a_j h_j + b_j e$ and

$$1_A(x + se + tNMh_1) = 1 - \sum_{j=1}^m P_{x,j}(s + b_j t).$$

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle qh_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne).$$

Properties of $P_{x,j}$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$.
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m - 2$.
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

We had $NMh_1 = a_j h_j + b_j e$ and

$$1_A(x + se + tNMh_1) = 1 - \sum_{j=1}^m P_{x,j}(s + b_j t).$$

Case 1: $\sup_{n \in \mathbb{N}} P_{x,j}(n) = 1$ for some $1 \leq j \leq m$.

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle qh_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne).$$

Properties of $P_{x,j}$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$.
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m - 2$.
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

We had $NMh_1 = a_j h_j + b_j e$ and

$$1_A(x + se + tNMh_1) = 1 - \sum_{j=1}^m P_{x,j}(s + b_j t).$$

Case 1: $\sup_{n \in \mathbb{N}} P_{x,j}(n) = 1$ for some $1 \leq j \leq m$. By the third property we have that $P_{x,i} = 0$ for all $i \neq j$.

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle qh_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne)$$

Properties of $P_{x,j}$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$.
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m - 2$.
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

We had $NMh_1 = a_j h_j + b_j e$ and

$$1_A(x + se + tNMh_1) = 1 - P_{x,j}(s + b_j t).$$

Case 1 ($\sup_{n \in \mathbb{N}} P_{x,j}(n) = 1$) : $1_A(x + (s + b_j t)e + ta_j h_j) = 1 - P_{x,j}(s + b_j t).$

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle qh_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne)$$

Properties of $P_{x,j}$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$.
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m - 2$.
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

We had $NMh_1 = a_j h_j + b_j e$ and

$$1_A(x + se + tNMh_1) = 1 - P_{x,j}(s + b_j t).$$

Case 1 ($\sup_{n \in \mathbb{N}} P_{x,j}(n) = 1$) : $1_A(x + (s + b_j t)e + ta_j h_j) = 1 - P_{x,j}(s + b_j t)$. Set $s + b_j t = s'$.

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle qh_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne)$$

Properties of $P_{x,j}$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$.
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m-2$.
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

We had $NMh_1 = a_j h_j + b_j e$ and

$$1_A(x + se + tNMh_1) = 1 - P_{x,j}(s + b_j t).$$

Case 1 ($\sup_{n \in \mathbb{N}} P_{x,j}(n) = 1$) : $1_A(x + (s + b_j t)e + ta_j h_j) = 1 - P_{x,j}(s + b_j t)$. Set $s + b_j t = s'$. We have that

$$1_A(x + s'e + ta_j h_j) = 1 - P_{x,j}(s').$$

This is true for all s' and t .

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle qh_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne)$$

Properties of $P_{x,j}$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$.
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m - 2$.
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

We had $NMh_1 = a_j h_j + b_j e$ and

$$1_A(x + se + tNMh_1) = 1 - P_{x,j}(s + b_j t).$$

Case 1 ($\sup_{n \in \mathbb{N}} P_{x,j}(n) = 1$) : $1_A(x + (s + b_j t)e + ta_j h_j) = 1 - P_{x,j}(s + b_j t)$. Set $s + b_j t = s'$. We have that

$$1_A(x + s'e + ta_j h_j) = 1 - P_{x,j}(s').$$

This is true for all s' and t . Thus $A \cap (x + \Lambda)$ is $\langle a_j h_j \rangle$ -periodic.

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle qh_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne).$$

Properties of $P_{x,j}$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$.
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m - 2$.
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

We had $NMh_1 = a_j h_j + b_j e$ and

$$1_A(x + se + tNMh_1) = 1 - \sum_{j=1}^m P_{x,j}(s + b_j t).$$

Case 2 ($\sup_{n \in \mathbb{N}} P_{x,j}(n) < 1$ for all j) :

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle qh_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne).$$

Properties of $P_{x,j}$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$.
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m - 2$.
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

We had $NMh_1 = a_j h_j + b_j e$ and

$$1_A(x + se + tNMh_1) = 1 - \sum_{j=1}^m P_{x,j}(s + b_j t).$$

Case 2 ($\sup_{n \in \mathbb{N}} P_{x,j}(n) < 1$ for all j) : By the third property of $P_{x,j}$ we have for all but one $1 \leq j \leq m$ (say j_0), $P_{x,j}(n) \in [0, 1/2]$.

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle qh_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne).$$

Properties of $P_{x,j}$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$.
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m - 2$.
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

We had $NMh_1 = a_j h_j + b_j e$ and

$$1_A(x + se + tNMh_1) = 1 - \sum_{j=1}^m P_{x,j}(s + b_j t).$$

Case 2 ($\sup_{n \in \mathbb{N}} P_{x,j}(n) < 1$ for all j) : By the third property of $P_{x,j}$ we have for all but one $1 \leq j \leq m$ (say j_0), $P_{x,j}(n) \in [0, 1/2]$. By Weyl equidistribution, $P_{x,j}$ is periodic for $j \neq j_0$.

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle qh_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne).$$

Properties of $P_{x,j}$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$.
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m-2$.
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

We had $NMh_1 = a_j h_j + b_j e$ and

$$1_A(x + se + tNMh_1) = 1 - \sum_{j=1}^m P_{x,j}(s + b_j t).$$

Case 2 ($\sup_{n \in \mathbb{N}} P_{x,j}(n) < 1$ for all j) : By the third property of $P_{x,j}$ we have for all but one $1 \leq j \leq m$ (say j_0), $P_{x,j}(n) \in [0, 1/2]$. By Weyl equidistribution, $P_{x,j}$ is periodic for $j \neq j_0$. By the first property we have that P_{x,j_0} must also be periodic.

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle qh_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne).$$

Properties of $P_{x,j}$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$.
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m - 2$.
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

We had $NMh_1 = a_j h_j + b_j e$ and

$$1_A(x + se + tNMh_1) = 1 - \sum_{j=1}^m P_{x,j}(s + b_j t).$$

Case 2 ($\sup_{n \in \mathbb{N}} P_{x,j}(n) < 1$ for all j) : By the third property of $P_{x,j}$ we have for all but one $1 \leq j \leq m$ (say j_0), $P_{x,j}(n) \in [0, 1/2]$. By Weyl equidistribution, $P_{x,j}$ is periodic for $j \neq j_0$. By the first property we have that P_{x,j_0} must also be periodic. Suppose $P_{x,j}$ have common period Q_m .

Proof of weak periodicity

$$1_A = 1 - \sum_{j=1}^m \phi_j \quad \star$$

where $\phi_j : \mathbb{Z}^d \rightarrow [0, 1]$ is $\langle qh_j \rangle$ periodic.

$$P_{x,j}(n) := \phi_j(x + ne).$$

Properties of $P_{x,j}$.

- ① One has $\sum_{j=1}^m P_{x,j}(n) = 0 \bmod 1$ for all $n \in \mathbb{Z}$.
- ② For each $1 \leq j \leq m$, the map $n \mapsto P_{x,j}(n) \bmod 1$ is a polynomial of degree at most $m - 2$.
- ③ For any $1 \leq i < j \leq m$, one has $\sup_{n \in \mathbb{Z}} P_{x,i}(n) + \sup_{n \in \mathbb{Z}} P_{x,j}(n) \leq 1$.

We had $NMh_1 = a_j h_j + b_j e$ and

$$1_A(x + se + tNMh_1) = 1 - \sum_{j=1}^m P_{x,j}(s + b_j t).$$

Case 2 ($\sup_{n \in \mathbb{N}} P_{x,j}(n) < 1$ for all j) : By the third property of $P_{x,j}$ we have for all but one $1 \leq j \leq m$ (say j_0), $P_{x,j}(n) \in [0, 1/2]$. By Weyl equidistribution, $P_{x,j}$ is periodic for $j \neq j_0$. By the first property we have that P_{x,j_0} must also be periodic. Suppose $P_{x,j}$ have common period Q_m . Then $(x + \Lambda) \cap A$ is $\langle Q_m MNh_1 \rangle$ -periodic and $\langle Q_m e \rangle$ -periodic.

What is left to do?

- ① Understanding the structure of higher level tilings.
 - ① Suppose a finite shape can tile at level k . Can it tile periodically at level k .
 - ② Given $h \in \mathbb{Z}^2$ we will say that $A \subset \mathbb{Z}^2$ is a Bohr open set in the direction h if for all $x \in \mathbb{Z}^2$, the set

$$\{n \in \mathbb{Z} : x + nh \in A\}$$

is the visit times of a compact group rotation to an open set.

Does there exist a full-rank sublattice Λ such that for all $A \in X_c$ and $y \in \mathbb{Z}^2$, $A \cap (y + \Lambda)$ is Bohr open in some direction?

- ② Higher dimensions.