

Four-Cycle Free Graphs and Entropy Minimality

Nishant Chandgotia

September, 2015

Outline

- Shifts of Finite Type
- Entropy
- Entropy Minimality
- The 3-coloured Chessboard.
- Universal Covers

Full Shift

Full Shift

- Let \mathcal{A} be a finite set (with the discrete topology). We will call it the **alphabet**.

Full Shift

- Let \mathcal{A} be a finite set (with the discrete topology). We will call it the **alphabet**.
- The space $\mathcal{A}^{\mathbb{Z}^d}$ is the set of all **configurations** on \mathbb{Z}^d . It is called the **full shift**.

Full Shift

- Let \mathcal{A} be a finite set (with the discrete topology). We will call it the **alphabet**.
- The space $\mathcal{A}^{\mathbb{Z}^d}$ is the set of all **configurations** on \mathbb{Z}^d . It is called the **full shift**.
- It is a compact space under the product topology with a natural \mathbb{Z}^d -action given by translations (also called **shifts**) of configurations.

Shift Spaces

Shift Spaces

- Given a finite subset $A \subset \mathbb{Z}^d$, a **pattern** on A is a function $a : A \rightarrow \mathcal{A}$.

Shift Spaces

- Given a finite subset $A \subset \mathbb{Z}^d$, a **pattern** on A is a function $a : A \rightarrow \mathcal{A}$.
- Let \mathcal{F} be a set of patterns (referred to as a **forbidden list**).

Shift Spaces

- Given a finite subset $A \subset \mathbb{Z}^d$, a **pattern** on A is a function $a : A \rightarrow \mathcal{A}$.
- Let \mathcal{F} be a set of patterns (referred to as a **forbidden list**).
- Denote

$$X_{\mathcal{F}} := \{x \in \mathcal{A}^{\mathbb{Z}^d} \mid \text{translates of patterns from } \mathcal{F} \text{ do not occur in } x\}.$$

Shift Spaces

- Given a finite subset $A \subset \mathbb{Z}^d$, a **pattern** on A is a function $a : A \rightarrow \mathcal{A}$.
- Let \mathcal{F} be a set of patterns (referred to as a **forbidden list**).
- Denote

$$X_{\mathcal{F}} := \{x \in \mathcal{A}^{\mathbb{Z}^d} \mid \text{translates of patterns from } \mathcal{F} \text{ do not occur in } x\}.$$

- A **shift space** is a subset $X \subset \mathcal{A}^{\mathbb{Z}^d}$ such that there exists a forbidden list \mathcal{F} satisfying $X = X_{\mathcal{F}}$.

Examples: Full Shift

Examples: Full Shift

- $\mathcal{A} = \{0, 1\}$, $\mathcal{F} = \emptyset$. $X_{\mathcal{F}} = \{0, 1\}^{\mathbb{Z}^d}$.

Examples: Hard Square Model

1	0	0	0	0
0	0	0	0	0
1	0	1	0	0
0	0	0	1	0
0	1	0	0	0

A Pattern

Examples: Hard Square Model

1	0	0	0	0
0	0	0	0	0
1	0	1	0	0
0	0	0	1	0
0	1	0	0	0

A Pattern

- $d = 2$

Examples: Hard Square Model

1	0	0	0	0
0	0	0	0	0
1	0	1	0	0
0	0	0	1	0
0	1	0	0	0

A Pattern

- $d = 2$
- $\mathcal{A} = \{0, 1\}$

Examples: Hard Square Model

1	0	0	0	0
0	0	0	0	0
1	0	1	0	0
0	0	0	1	0
0	1	0	0	0

A Pattern

- $d = 2$
- $\mathcal{A} = \{0, 1\}$
- $\mathcal{F} = \{11, \frac{1}{1}\}$

Examples: Hard Square Model

1	0	0	0	0
0	0	0	0	0
1	0	1	0	0
0	0	0	1	0
0	1	0	0	0

A Pattern

- $d = 2$
- $\mathcal{A} = \{0, 1\}$
- $\mathcal{F} = \{11, \frac{1}{1}\}$
- By \mathbb{Z}^d , we will refer to the Cayley graph of \mathbb{Z}^d with standard generators. Thus \mathbb{Z}^2 is the grid.

Examples: Hard Square Model

1	0	0	0	0
0	0	0	0	0
1	0	1	0	0
0	0	0	1	0
0	1	0	0	0

A Pattern

- $d = 2$
- $\mathcal{A} = \{0, 1\}$
- $\mathcal{F} = \{11, \begin{smallmatrix} 1 \\ 1 \end{smallmatrix}\}$
- By \mathbb{Z}^d , we will refer to the Cayley graph of \mathbb{Z}^d with standard generators. Thus \mathbb{Z}^2 is the grid.
- $X_{\mathcal{F}} =$
 {configurations in 0 and 1 where two 1's cannot be adjacent}.

Examples: Even Shift

00001000010011000000100000000100000

Examples: Even Shift

00001000010011000000100000000100000

- $d = 1$

Examples: Even Shift

000010000100110000000100000000100000

- $d = 1$
- $\mathcal{A} = \{0, 1\}$

Examples: Even Shift

000010000100110000000100000000100000

- $d = 1$
- $\mathcal{A} = \{0, 1\}$
- $\mathcal{F} = \{10^{2i-1}1 \mid i \in \mathbb{N}\}$

Examples: Even Shift

0000100001001100000001000000000100000

- $d = 1$
- $\mathcal{A} = \{0, 1\}$
- $\mathcal{F} = \{10^{2i-1}1 \mid i \in \mathbb{N}\}$
- $X_{\mathcal{F}} =$
 $\{0, 1 \text{ sequences such that the gap between any two } 1\text{'s is even}\}.$

Examples: Even Shift

000010000100110000000100000000100000

- $d = 1$
- $\mathcal{A} = \{0, 1\}$
- $\mathcal{F} = \{10^{2i-1}1 \mid i \in \mathbb{N}\}$
- $X_{\mathcal{F}} =$
 $\{0, 1 \text{ sequences such that the gap between any two } 1\text{'s is even}\}.$
- Note that \mathcal{F} is infinite.

Examples: Even Shift

000010000100110000000100000000100000

- $d = 1$
- $\mathcal{A} = \{0, 1\}$
- $\mathcal{F} = \{10^{2i-1}1 \mid i \in \mathbb{N}\}$
- $X_{\mathcal{F}} =$
 $\{0, 1 \text{ sequences such that the gap between any two } 1\text{'s is even}\}.$
- Note that \mathcal{F} is infinite.
- It can be proved that \mathcal{F} cannot be chosen to finite!

Shifts of Finite Type

Shifts of Finite Type

- A shift space X is called a **shift of finite type** if there exists a finite set \mathcal{F} such that $X = X_{\mathcal{F}}$.

Shifts of Finite Type

- A shift space X is called a **shift of finite type** if there exists a finite set \mathcal{F} such that $X = X_{\mathcal{F}}$.
- Both the hard square model and the full shift are shifts of finite type.

Shifts of Finite Type

- A shift space X is called a **shift of finite type** if there exists a finite set \mathcal{F} such that $X = X_{\mathcal{F}}$.
- Both the hard square model and the full shift are shifts of finite type.
- The even shift is not a shift of finite type.

Nearest Neighbour Shifts of Finite Type

Nearest Neighbour Shifts of Finite Type

- A shift space X is called a nearest neighbour shift of finite type if there exists a set of patterns \mathcal{F} such that

Nearest Neighbour Shifts of Finite Type

- A shift space X is called a nearest neighbour shift of finite type if there exists a set of patterns \mathcal{F} such that
 - $X = X_{\mathcal{F}}$.

Nearest Neighbour Shifts of Finite Type

- A shift space X is called a nearest neighbour shift of finite type if there exists a set of patterns \mathcal{F} such that
 - $X = X_{\mathcal{F}}$.
 - \mathcal{F} consists of patterns on edges of \mathbb{Z}^d .

Nearest Neighbour Shifts of Finite Type

- A shift space X is called a nearest neighbour shift of finite type if there exists a set of patterns \mathcal{F} such that
 - $X = X_{\mathcal{F}}$.
 - \mathcal{F} consists of patterns on edges of \mathbb{Z}^d .
- The hard square model is a nearest neighbour shift of finite type.

Examples: Non-Attacking Kings

1	0	1	0	0	0
0	0	0	0	1	0
0	0	0	0	0	0
0	0	1	0	0	1
0	0	0	0	0	0
0	0	0	0	1	0

Examples: Non-Attacking Kings

1	0	1	0	0	0
0	0	0	0	1	0
0	0	0	0	0	0
0	0	1	0	0	1
0	0	0	0	0	0
0	0	0	0	1	0

- $d = 2$, $\mathcal{A} = \{0, 1\}$, $\mathcal{F} = \{11, \begin{smallmatrix} 1 \\ 1 \end{smallmatrix}, \begin{smallmatrix} 1 & 1 \\ 1 & 1 \end{smallmatrix}\}$

Examples: Non-Attacking Kings

1	0	1	0	0	0
0	0	0	0	1	0
0	0	0	0	0	0
0	0	1	0	0	1
0	0	0	0	0	0
0	0	0	0	1	0

- $d = 2$, $\mathcal{A} = \{0, 1\}$, $\mathcal{F} = \{11, \begin{smallmatrix} 1 \\ 1 \end{smallmatrix}, \begin{smallmatrix} 1 & 1 \\ 1 & 1 \end{smallmatrix}\}$
- $X_{\mathcal{F}}$ is a shift of finite type but not at a nearest neighbour shift of finite type.

Examples: Non-Attacking Kings

1	0	1	0	0	0
0	0	0	0	1	0
0	0	0	0	0	0
0	0	1	0	0	1
0	0	0	0	0	0
0	0	0	0	1	0

- $d = 2$, $\mathcal{A} = \{0, 1\}$, $\mathcal{F} = \{11, \begin{smallmatrix} 1 \\ 1 \end{smallmatrix}, \begin{smallmatrix} 1 & 1 \\ 1 & 1 \end{smallmatrix}\}$
- $X_{\mathcal{F}}$ is a shift of finite type but not at a nearest neighbour shift of finite type.
- Any shift of finite type can be recoded into a nearest neighbour shift of finite type (for a different alphabet).

When is a Nearest Neighbour Shift of Finite Type
Non-Empty?

When is a Nearest Neighbour Shift of Finite Type
Non-Empty?

- Let $d = 1$.

When is a Nearest Neighbour Shift of Finite Type Non-Empty?

- Let $d = 1$.
- \mathcal{F} tells us what letter can sit next to another letter.

When is a Nearest Neighbour Shift of Finite Type Non-Empty?

- Let $d = 1$.
- \mathcal{F} tells us what letter can sit next to another letter.
- Given \mathcal{F} we can build a directed graph \mathcal{G} :

When is a Nearest Neighbour Shift of Finite Type Non-Empty?

- Let $d = 1$.
- \mathcal{F} tells us what letter can sit next to another letter.
- Given \mathcal{F} we can build a directed graph \mathcal{G} :
 - The vertices are the elements of \mathcal{A}

When is a Nearest Neighbour Shift of Finite Type Non-Empty?

- Let $d = 1$.
- \mathcal{F} tells us what letter can sit next to another letter.
- Given \mathcal{F} we can build a directed graph \mathcal{G} :
 - The vertices are the elements of \mathcal{A}
 - There is an edge from $v \in \mathcal{A}$ to $w \in \mathcal{A}$ if and only if $vw \notin \mathcal{F}$.

When is a Nearest Neighbour Shift of Finite Type Non-Empty?

- Let $d = 1$.
- \mathcal{F} tells us what letter can sit next to another letter.
- Given \mathcal{F} we can build a directed graph \mathcal{G} :
 - The vertices are the elements of \mathcal{A}
 - There is an edge from $v \in \mathcal{A}$ to $w \in \mathcal{A}$ if and only if $vw \notin \mathcal{F}$.
- $X_{\mathcal{F}}$ is non-empty if and only if there is an infinite path in \mathcal{G} .

When is a Nearest Neighbour Shift of Finite Type Non-Empty?

- Let $d = 1$.
- \mathcal{F} tells us what letter can sit next to another letter.
- Given \mathcal{F} we can build a directed graph \mathcal{G} :
 - The vertices are the elements of \mathcal{A}
 - There is an edge from $v \in \mathcal{A}$ to $w \in \mathcal{A}$ if and only if $vw \notin \mathcal{F}$.
- $X_{\mathcal{F}}$ is non-empty if and only if there is an infinite path in \mathcal{G} .
- It is decidable.

When is a Nearest Neighbour Shift of Finite Type
Non-Empty?

When is a Nearest Neighbour Shift of Finite Type
Non-Empty?

- Let $d \geq 2$.

When is a Nearest Neighbour Shift of Finite Type Non-Empty?

- Let $d \geq 2$.
- Given \mathcal{F} there exists no algorithm to decide when $X_{\mathcal{F}}$ is non-empty.

When is a Nearest Neighbour Shift of Finite Type Non-Empty?

- Let $d \geq 2$.
- Given \mathcal{F} there exists no algorithm to decide when $X_{\mathcal{F}}$ is non-empty.
- So we deal with a more restricted class of shift spaces.

Hom-shifts

Hom-shifts

- Let \mathcal{H} be a finite undirected graph without multiple edges.

Hom-shifts

- Let \mathcal{H} be a finite undirected graph without multiple edges.
- Consider a shift space $X_{\mathcal{H}} := \text{Hom}(\mathbb{Z}^d, \mathcal{H})$; symbols on adjacent vertices of \mathbb{Z}^d form an edge in the graph \mathcal{H} .


Hom-shifts

- Let \mathcal{H} be a finite undirected graph without multiple edges.
- Consider a shift space $X_{\mathcal{H}} := \text{Hom}(\mathbb{Z}^d, \mathcal{H})$; symbols on adjacent vertices of \mathbb{Z}^d form an edge in the graph \mathcal{H} .
- A **hom-shift** is a shift space X for which there exists a graph \mathcal{H} such that $X = X_{\mathcal{H}}$.

Hom-shifts

- Let \mathcal{H} be a finite undirected graph without multiple edges.
- Consider a shift space $X_{\mathcal{H}} := \text{Hom}(\mathbb{Z}^d, \mathcal{H})$; symbols on adjacent vertices of \mathbb{Z}^d form an edge in the graph \mathcal{H} .
- A **hom-shift** is a shift space X for which there exists a graph \mathcal{H} such that $X = X_{\mathcal{H}}$.

Examples: (Hard Square model)



Graph \mathcal{H}


1	0	0	0	0
0	0	0	0	0
1	0	1	0	0
0	0	0	1	0
0	1	0	0	0

A Pattern


Hom-shifts

- Let \mathcal{H} be a finite undirected graph without multiple edges.
- Consider a shift space $X_{\mathcal{H}} := \text{Hom}(\mathbb{Z}^d, \mathcal{H})$; symbols on adjacent vertices of \mathbb{Z}^d form an edge in the graph \mathcal{H} .
- A **hom-shift** is a shift space X for which there exists a graph \mathcal{H} such that $X = X_{\mathcal{H}}$.

Examples: (3-colourings)



Graph \mathcal{H}



A Pattern

Hom-shifts are Nearest Neighbour Shifts of Finite Type

Hom-shifts are Nearest Neighbour Shifts of Finite Type

- Let $d = 2$

Hom-shifts are Nearest Neighbour Shifts of Finite Type

- Let $d = 2$
- Let \mathcal{H} be a finite undirected graph.

Hom-shifts are Nearest Neighbour Shifts of Finite Type

- Let $d = 2$
- Let \mathcal{H} be a finite undirected graph.
- $\mathcal{A} =$ vertex set of \mathcal{H} .

Hom-shifts are Nearest Neighbour Shifts of Finite Type

- Let $d = 2$
- Let \mathcal{H} be a finite undirected graph.
- \mathcal{A} = vertex set of \mathcal{H} .
- $\mathcal{F} = \{vw, \overset{v}{w} \mid v, w \in \mathcal{A} \text{ and } (v, w) \text{ does not form an edge in } \mathcal{H}\}$.

Hom-shifts are Nearest Neighbour Shifts of Finite Type

- Let $d = 2$
- Let \mathcal{H} be a finite undirected graph.
- \mathcal{A} = vertex set of \mathcal{H} .
- $\mathcal{F} = \{vw, \overset{v}{\underset{w}{\curvearrowright}} \mid v, w \in \mathcal{A} \text{ and } (v, w) \text{ does not form an edge in } \mathcal{H}\}$.
- Then $X_{\mathcal{H}} = X_{\mathcal{F}}$


Hom-shifts are Nearest Neighbour Shifts of Finite Type

- Let $d = 2$
- Let \mathcal{H} be a finite undirected graph.
- \mathcal{A} = vertex set of \mathcal{H} .
- $\mathcal{F} = \{vw, \overset{v}{\underset{w}{\curvearrowright}} \mid v, w \in \mathcal{A} \text{ and } (v, w) \text{ does not form an edge in } \mathcal{H}\}$.
- Then $X_{\mathcal{H}} = X_{\mathcal{F}}$
- This is a nearest neighbour shift of finite type for which every direction has the same constraint.


Hom-shifts are Nearest Neighbour Shifts of Finite Type

- Let $d = 2$
- Let \mathcal{H} be a finite undirected graph.
- $\mathcal{A} =$ vertex set of \mathcal{H} .
- $\mathcal{F} = \{vw, \overset{v}{\underset{w}{\curvearrowright}} \mid v, w \in \mathcal{A} \text{ and } (v, w) \text{ does not form an edge in } \mathcal{H}\}.$
- Then $X_{\mathcal{H}} = X_{\mathcal{F}}$
- This is a nearest neighbour shift of finite type for which every direction has the same constraint.
- $X_{\mathcal{H}}$ is non-empty if and only if \mathcal{H} has an edge.

Examples: Domino Tilings



Examples: Domino Tilings



- What is the alphabet?

Examples: Domino Tilings

L	R	U	L	R
L	R	D	U	U
U	L	R	D	D
D	U	U	L	R
U	D	D	U	U
D	L	R	D	D

Examples: Domino Tilings

L	R	U	L	R
L	R	D	U	U
U	L	R	D	D
D	U	U	L	R
U	D	D	U	U
D	L	R	D	D

- $\mathcal{A} = \{U, D, L, R\}$.

Examples: Domino Tilings

L	R	U	L	R
L	R	D	U	U
U	L	R	D	D
D	U	U	L	R
U	D	D	U	U
D	L	R	D	D


- $\mathcal{A} = \{U, D, L, R\}$.
- $\mathcal{F} = \{UR, DR, RR, LL, LU, LD, \overset{U}{U}, \overset{D}{D}, \overset{U}{L}, \overset{U}{R}, \overset{L}{D}, \overset{R}{D}\}$

Examples: Domino Tilings

L	R	U	L	R
L	R	D	U	U
U	L	R	D	D
D	U	U	L	R
U	D	D	U	U
D	L	R	D	D

- $\mathcal{A} = \{U, D, L, R\}$.
- $\mathcal{F} = \{UR, DR, RR, LL, LU, LD, \frac{U}{U}, \frac{D}{D}, \frac{U}{L}, \frac{U}{R}, \frac{L}{D}, \frac{R}{D}\}$
- The constraints in different directions are different. It is not a hom-shift.

Shift Spaces Schematic



Entropy

Entropy

The **language** $\mathcal{L}(X)$ of a shift space is the set of patterns appearing in elements of X .

Entropy

The **language** $\mathcal{L}(X)$ of a shift space is the set of patterns appearing in elements of X .

$\{1, 2, \dots, n\}^d$ is an n -sided square box in \mathbb{Z}^d .

Entropy

The **language** $\mathcal{L}(X)$ of a shift space is the set of patterns appearing in elements of X .

$\{1, 2, \dots, n\}^d$ is an n -sided square box in \mathbb{Z}^d .

Let $\mathcal{L}_n(X) := \mathcal{L}(X) \cap \mathbb{A}^{\{1,2,\dots,n\}^d}$ be patterns which appear on an n -sided square box.

Entropy

The **language** $\mathcal{L}(X)$ of a shift space is the set of patterns appearing in elements of X .

$\{1, 2, \dots, n\}^d$ is an n -sided square box in \mathbb{Z}^d .

Let $\mathcal{L}_n(X) := \mathcal{L}(X) \cap \mathcal{A}^{\{1,2,\dots,n\}^d}$ be patterns which appear on an n -sided square box.

The **topological entropy** of X is defined as

Entropy

The **language** $\mathcal{L}(X)$ of a shift space is the set of patterns appearing in elements of X .

$\{1, 2, \dots, n\}^d$ is an n -sided square box in \mathbb{Z}^d .

Let $\mathcal{L}_n(X) := \mathcal{L}(X) \cap \{1, 2, \dots, n\}^d$ be patterns which appear on an n -sided square box.

The **topological entropy** of X is defined as

$$h_{\text{top}}(X) := \lim_{n \rightarrow \infty} \frac{\log |\mathcal{L}_n(X)|}{n^d}.$$

Entropy

The **language** $\mathcal{L}(X)$ of a shift space is the set of patterns appearing in elements of X .

$\{1, 2, \dots, n\}^d$ is an n -sided square box in \mathbb{Z}^d .

Let $\mathcal{L}_n(X) := \mathcal{L}(X) \cap \{1, 2, \dots, n\}^d$ be patterns which appear on an n -sided square box.

The **topological entropy** of X is defined as

$$h_{\text{top}}(X) := \lim_{n \rightarrow \infty} \frac{\log |\mathcal{L}_n(X)|}{n^d}.$$

(The limit always exists.)

Examples: Full Shift

Examples: Full Shift

- Let $X = \mathbb{A}^{\mathbb{Z}^d}$.

Examples: Full Shift

- Let $X = \mathfrak{a}^{\mathbb{Z}^d}$.
- $|\mathcal{L}_n(X)| = |\mathfrak{a}^{\{1,2,\dots,n\}^d}|$

Examples: Full Shift

- Let $X = \mathfrak{A}^{\mathbb{Z}^d}$.
- $|\mathcal{L}_n(X)| = |\mathfrak{A}^{\{1,2,\dots,n\}^d}|$
- $$h_{top}(X) := \lim_{n \rightarrow \infty} \frac{\log |\mathcal{L}_n(X)|}{n^d} = \lim_{n \rightarrow \infty} \frac{\log |\mathfrak{A}^{\{1,2,\dots,n\}^d}|}{n^d} = \log |\mathfrak{A}|.$$

Examples: Entropy of Shifts of Finite Type in One Dimension

Examples: Entropy of Shifts of Finite Type in One Dimension

- The entropy of a nearest neighbour shift of finite type is the logarithm of the greatest eigenvalue of a non-negative matrix.

Examples: Entropy of Shifts of Finite Type in One Dimension

- The entropy of a nearest neighbour shift of finite type is the logarithm of the greatest eigenvalue of a non-negative matrix.
- For instance, if \mathcal{H} is a finite undirected graph and $\lambda_{\mathcal{H}}$ is the greatest eigenvalue of its adjacency matrix then


$$h_{top}(X_{\mathcal{H}}) = \log \lambda_{\mathcal{H}}.$$

Examples: Entropy of Shifts of Finite Type in One Dimension

- The entropy of a nearest neighbour shift of finite type is the logarithm of the greatest eigenvalue of a non-negative matrix.
- For instance, if \mathcal{H} is a finite undirected graph and $\lambda_{\mathcal{H}}$ is the greatest eigenvalue of its adjacency matrix then

$$h_{top}(X_{\mathcal{H}}) = \log \lambda_{\mathcal{H}}.$$

- The entropy of the hard square model is $\log \left(\frac{\sqrt{5}+1}{2} \right)$. Hence it is also called the golden mean shift.




Examples: Entropy of Shifts of Finite Type in One Dimension

- The entropy of a nearest neighbour shift of finite type is the logarithm of the greatest eigenvalue of a non-negative matrix.
- For instance, if \mathcal{H} is a finite undirected graph and $\lambda_{\mathcal{H}}$ is the greatest eigenvalue of its adjacency matrix then

$$h_{top}(X_{\mathcal{H}}) = \log \lambda_{\mathcal{H}}.$$

- The entropy of the hard square model is $\log \left(\frac{\sqrt{5}+1}{2} \right)$. Hence it is also called the golden mean shift.
- The entropy of the space of 3-colourings is $\log(2)$.



Examples: Entropy of Shifts of Finite Type in Higher Dimensions

Examples: Entropy of Shifts of Finite Type in Higher Dimensions

- For $d \geq 2$ even finding good approximations of the entropy of nearest neighbour shifts of finite type can be difficult.

Examples: Entropy of Shifts of Finite Type in Higher Dimensions

- For $d \geq 2$ even finding good approximations of the entropy of nearest neighbour shifts of finite type can be difficult.
- (*Hochman and Meyerovitch, '07*) The set of entropies of nearest neighbour shifts of finite type for $d > 1$ is the set of non-negative **right recursively enumerable numbers**: those numbers for which there is an algorithm to generate rational numbers approximating it from above.

Examples: Entropy of Shifts of Finite Type in Higher Dimensions

- For $d \geq 2$ even finding good approximations of the entropy of nearest neighbour shifts of finite type can be difficult.
- (*Hochman and Meyerovitch, '07*) The set of entropies of nearest neighbour shifts of finite type for $d > 1$ is the set of non-negative **right recursively enumerable numbers**: those numbers for which there is an algorithm to generate rational numbers approximating it from above.
- Thus for most general shifts of finite type there is no hope of obtaining a 'reasonable' closed-form expression for the entropy.

Examples: Entropy of Shifts of Finite Type in Higher Dimensions

- For $d \geq 2$ even finding good approximations of the entropy of nearest neighbour shifts of finite type can be difficult.
- (*Hochman and Meyerovitch, '07*) The set of entropies of nearest neighbour shifts of finite type for $d > 1$ is the set of non-negative **right recursively enumerable numbers**: those numbers for which there is an algorithm to generate rational numbers approximating it from above.
- Thus for most general shifts of finite type there is no hope of obtaining a 'reasonable' closed-form expression for the entropy.
- Closed forms are known for very few examples.

Entropy for Hom-Shifts

Entropy for Hom-Shifts

- (*Friedlander, 1997*) There are approximating upper and lower bounds for the entropy of hom-shifts.

Entropy Minimality

Entropy Minimality

- If $Y \subset X$ are shift spaces then

$$h_{top}(Y) \leq h_{top}(X).$$

Entropy Minimality

- If $Y \subset X$ are shift spaces then

$$h_{top}(Y) \leq h_{top}(X).$$

- A shift space X is said to be **entropy minimal** if for all shift spaces $Y \subsetneq X$, $h_{top}(Y) < h_{top}(X)$.

Entropy Minimality

- If $Y \subset X$ are shift spaces then

$$h_{top}(Y) \leq h_{top}(X).$$

- A shift space X is said to be **entropy minimal** if for all shift spaces $Y \subsetneq X$, $h_{top}(Y) < h_{top}(X)$.
- That is, if we forbid any pattern from the language $\mathcal{L}(X)$ the entropy will drop.

Entropy Minimality in One Dimension

Entropy Minimality in One Dimension

Let $X = Y \cup Z$ where Y and Z are disjoint nearest neighbour shifts of finite type such that $h_{top}(Y) \geq h_{top}(Z)$.

Entropy Minimality in One Dimension

Let $X = Y \cup Z$ where Y and Z are disjoint nearest neighbour shifts of finite type such that $h_{top}(Y) \geq h_{top}(Z)$.

$$h_{top}(Y) \leq h_{top}(X)$$

Entropy Minimality in One Dimension

Let $X = Y \cup Z$ where Y and Z are disjoint nearest neighbour shifts of finite type such that $h_{top}(Y) \geq h_{top}(Z)$.

$$\begin{aligned} h_{top}(Y) &\leq h_{top}(X) \\ &= \lim_{n \rightarrow \infty} \frac{\log |\mathcal{L}_n(X)|}{n^d} \end{aligned}$$

Entropy Minimality in One Dimension

Let $X = Y \cup Z$ where Y and Z are disjoint nearest neighbour shifts of finite type such that $h_{top}(Y) \geq h_{top}(Z)$.

$$\begin{aligned} h_{top}(Y) &\leq h_{top}(X) \\ &= \lim_{n \rightarrow \infty} \frac{\log |\mathcal{L}_n(X)|}{n^d} \\ &\leq \lim_{n \rightarrow \infty} \frac{\log(|\mathcal{L}_n(Y)| + |\mathcal{L}_n(Z)|)}{n^d} \end{aligned}$$

Entropy Minimality in One Dimension

Let $X = Y \cup Z$ where Y and Z are disjoint nearest neighbour shifts of finite type such that $h_{top}(Y) \geq h_{top}(Z)$.

$$\begin{aligned} h_{top}(Y) &\leq h_{top}(X) \\ &= \lim_{n \rightarrow \infty} \frac{\log |\mathcal{L}_n(X)|}{n^d} \\ &\leq \lim_{n \rightarrow \infty} \frac{\log(|\mathcal{L}_n(Y)| + |\mathcal{L}_n(Z)|)}{n^d} \\ &\leq \lim_{n \rightarrow \infty} \frac{\log(2 \max(|\mathcal{L}_n(Y)|, |\mathcal{L}_n(Z)|))}{n^d} \end{aligned}$$

Entropy Minimality in One Dimension

Let $X = Y \cup Z$ where Y and Z are disjoint nearest neighbour shifts of finite type such that $h_{top}(Y) \geq h_{top}(Z)$.

$$\begin{aligned} h_{top}(Y) &\leq h_{top}(X) \\ &= \lim_{n \rightarrow \infty} \frac{\log |\mathcal{L}_n(X)|}{n^d} \\ &\leq \lim_{n \rightarrow \infty} \frac{\log(|\mathcal{L}_n(Y)| + |\mathcal{L}_n(Z)|)}{n^d} \\ &\leq \lim_{n \rightarrow \infty} \frac{\log(2 \max(|\mathcal{L}_n(Y)|, |\mathcal{L}_n(Z)|))}{n^d} \\ &\leq h_{top}(Y). \end{aligned}$$

Entropy Minimality in One Dimension

Let $X = Y \cup Z$ where Y and Z are disjoint nearest neighbour shifts of finite type such that $h_{top}(Y) \geq h_{top}(Z)$.

$$\begin{aligned} h_{top}(Y) &\leq h_{top}(X) \\ &= \lim_{n \rightarrow \infty} \frac{\log |\mathcal{L}_n(X)|}{n^d} \\ &\leq \lim_{n \rightarrow \infty} \frac{\log(|\mathcal{L}_n(Y)| + |\mathcal{L}_n(Z)|)}{n^d} \\ &\leq \lim_{n \rightarrow \infty} \frac{\log(2 \max(|\mathcal{L}_n(Y)|, |\mathcal{L}_n(Z)|))}{n^d} \\ &\leq h_{top}(Y). \end{aligned}$$

Thus $h_{top}(X) = h_{top}(Y)$.

Entropy Minimality in One Dimension

Entropy Minimality in One Dimension

- Fix $d = 1$.

Entropy Minimality in One Dimension

- Fix $d = 1$.
- We say that a nearest neighbour shift space X is called **irreducible** if $X \neq Y \cup Z$ for some disjoint shift spaces Y and Z .

Entropy Minimality in One Dimension

- Fix $d = 1$.
- We say that a nearest neighbour shift space X is called **irreducible** if $X \neq Y \cup Z$ for some disjoint shift spaces Y and Z .
- We just proved that if a shift space X which is not irreducible, then it is not entropy minimal.

Entropy Minimality in One Dimension

- Fix $d = 1$.
- We say that a nearest neighbour shift space X is called **irreducible** if $X \neq Y \cup Z$ for some disjoint shift spaces Y and Z .
- We just proved that if a shift space X which is not irreducible, then it is not entropy minimal.
- In fact it can be proved that a shift space is irreducible if and only if it is entropy minimal.

Entropy Minimality in Higher Dimensions

It is undecidable whether or not a nearest neighbour shift of finite type is entropy minimal.

Main Result

Main Result

Theorem (Chandgotia, Meyerovitch '13)

Let C_n be an n -cycle. Then X_{C_n} is entropy minimal.

Main Result

Theorem (Chandgotia, Meyerovitch '13)

Let C_n be an n -cycle. Then X_{C_n} is entropy minimal.

A connected, finite graph \mathcal{H} is called **four-cycle free** if it has no self-loops and C_4 is not a subgraph of \mathcal{H} .

Main Result

Theorem (Chandgotia, Meyerovitch '13)

Let C_n be an n -cycle. Then X_{C_n} is entropy minimal.

A connected, finite graph \mathcal{H} is called **four-cycle free** if it has no self-loops and C_4 is not a subgraph of \mathcal{H} .


Theorem (Chandgotia '14)

If \mathcal{H} is a four-cycle free graph then $X_{\mathcal{H}}$ is entropy minimal.

A Technical Result: Some Notation

A Technical Result: Some Notation


- $A \subset \mathbb{Z}^d$ is a finite subset.



- $-$ Elements of A

A Technical Result: Some Notation


- $A \subset \mathbb{Z}^d$ is a finite subset.
- X is a shift space.



- $-$ Elements of A

A Technical Result: Some Notation

- $A \subset \mathbb{Z}^d$ is a finite subset.
- X is a shift space.
- For a pattern $a \in \mathcal{L}(X) \cap \mathfrak{A}^A$ consider




- –Elements of A

A Technical Result: Some Notation

- $A \subset \mathbb{Z}^d$ is a finite subset.
- X is a shift space.
- For a pattern $a \in \mathcal{L}(X) \cap \mathcal{A}^A$ consider

$$[a]_A := \{x \in X \mid x|_A = a\} \text{ (Cylinder set)}.$$



- –Elements of A

Cylinder set $[4,3,1]_A$

A Technical Result: Measures of Maximal Entropy

A Technical Result: Measures of Maximal Entropy

- As with topological entropy, for a given shift-invariant Borel probability measure μ on a shift space we denote its **measure-theoretic entropy** by h_μ .

A Technical Result: Measures of Maximal Entropy

- As with topological entropy, for a given shift-invariant Borel probability measure μ on a shift space we denote its **measure-theoretic entropy** by h_μ .
- (*Variational Principle*) Given a shift space X

$$\sup_{\text{supp}(\mu) \subset X} h_\mu = h_{\text{top}}(X).$$

A Technical Result: Measures of Maximal Entropy

- As with topological entropy, for a given shift-invariant Borel probability measure μ on a shift space we denote its **measure-theoretic entropy** by h_μ .
- (*Variational Principle*) Given a shift space X

$$\sup_{\text{supp}(\mu) \subset X} h_\mu = h_{\text{top}}(X).$$

- The supremum is achieved; such measures are called **measures of maximal entropy**.

A Technical Result: Measures of Maximal Entropy

- As with topological entropy, for a given shift-invariant Borel probability measure μ on a shift space we denote its **measure-theoretic entropy** by h_μ .
- (*Variational Principle*) Given a shift space X

$$\sup_{\text{supp}(\mu) \subset X} h_\mu = h_{\text{top}}(X).$$

- The supremum is achieved; such measures are called **measures of maximal entropy**.
- If μ is a measure of maximal entropy for X then $h_\mu = h_{\text{top}}(X)$.

A Technical Result: Measures of Maximal Entropy

- As with topological entropy, for a given shift-invariant Borel probability measure μ on a shift space we denote its **measure-theoretic entropy** by h_μ .
- (Variational Principle) Given a shift space X

$$\sup_{\text{supp}(\mu) \subset X} h_\mu = h_{\text{top}}(X).$$

- The supremum is achieved; such measures are called **measures of maximal entropy**.
- If μ is a measure of maximal entropy for X then $h_\mu = h_{\text{top}}(X)$.

Theorem

A shift space X is entropy minimal if and only if for every measure of maximal entropy μ , $\text{supp}(\mu) = X$.

A Technical Result: Gibbs Measures

A Technical Result: Gibbs Measures

(Lanford and Ruelle '69-Special case):

A Technical Result: Gibbs Measures

(Lanford and Ruelle '69-Special case): If X is a shift of finite type then for any measure of maximal entropy μ ,

A Technical Result: Gibbs Measures

(*Lanford and Ruelle '69-Special case*): If X is a shift of finite type then for any measure of maximal entropy μ , finite set A ,
 $\partial A \subset B \subset A^c$

A Technical Result: Gibbs Measures

(Lanford and Ruelle '69-Special case): If X is a shift of finite type then for any measure of maximal entropy μ , finite set A , $\partial A \subset B \subset A^c$ and $[b]_B$ of positive measure

A Technical Result: Gibbs Measures

(*Lanford and Ruelle '69-Special case*): If X is a shift of finite type then for any measure of maximal entropy μ , finite set A , $\partial A \subset B \subset A^c$ and $[b]_B$ of positive measure

$$\mu([a]_A \mid [b]_B)$$

is uniform over all patterns $a \in \mathcal{L}(X) \cap \mathfrak{A}^A$ satisfying $ab \in \mathcal{L}(X)$.

A Technical Result: Gibbs Measures

(Lanford and Ruelle '69-Special case): If X is a shift of finite type then for any measure of maximal entropy μ , finite set A , $\partial A \subset B \subset A^c$ and $[b]_B$ of positive measure

$$\mu([a]_A \mid [b]_B)$$

is uniform over all patterns $a \in \mathcal{L}(X) \cap \mathfrak{A}^A$ satisfying $ab \in \mathcal{L}(X)$.

Such measures are called uniform Gibbs measures.

A Technical Result: Gibbs Measures

(Lanford and Ruelle '69-Special case): If X is a shift of finite type then for any measure of maximal entropy μ , finite set A , $\partial A \subset B \subset A^c$ and $[b]_B$ of positive measure

$$\mu([a]_A \mid [b]_B)$$

is uniform over all patterns $a \in \mathcal{L}(X) \cap \mathfrak{A}^A$ satisfying $ab \in \mathcal{L}(X)$.

Such measures are called uniform Gibbs measures.

For example, if μ is the measure of maximal entropy in the hard square model then

A Technical Result: Gibbs Measures

(Lanford and Ruelle '69-Special case): If X is a shift of finite type then for any measure of maximal entropy μ , finite set A , $\partial A \subset B \subset A^c$ and $[b]_B$ of positive measure

$$\mu([a]_A \mid [b]_B)$$

is uniform over all patterns $a \in \mathcal{L}(X) \cap \mathfrak{A}^A$ satisfying $ab \in \mathcal{L}(X)$.

Such measures are called uniform Gibbs measures.

For example, if μ is the measure of maximal entropy in the hard square model then

$$\mu([0]_0 \mid \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}_{\partial 0}) = \mu([1]_0 \mid \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}_{\partial 0}) = \frac{1}{2}.$$

Entropy Minimality of X_{C_3}

The Cayley graph of $\mathbb{Z}/3\mathbb{Z}$ is C_3 .

Let us see why X_{C_3} is entropy minimal.

Height Functions and X_{C_3}

Height Functions and X_{C_3}

- A **height function** is an element of $X_{\mathbb{Z}}$.

Height Functions and X_{C_3}


- A **height function** is an element of $X_{\mathbb{Z}}$.
- For all $h \in X_{\mathbb{Z}}$, $h \bmod 3 \in X_{C_3}$.

Height Functions and X_{C_3}

- A **height function** is an element of $X_{\mathbb{Z}}$.
- For all $h \in X_{\mathbb{Z}}$, $h \bmod 3 \in X_{C_3}$.
- Conversely, given a configuration in X_{C_3} there exists a unique (up to an additive constant) height function corresponding to it.

Height Functions and X_{C_3}


- A **height function** is an element of $X_{\mathbb{Z}}$.
- For all $h \in X_{\mathbb{Z}}$, $h \bmod 3 \in X_{C_3}$.
- Conversely, given a configuration in X_{C_3} there exists a unique (up to an additive constant) height function corresponding to it.



Height Function

Height Functions and X_{C_3}

- A **height function** is an element of $X_{\mathbb{Z}}$.
- For all $h \in X_{\mathbb{Z}}$, $h \bmod 3 \in X_{C_3}$.
- Conversely, given a configuration in X_{C_3} there exists a unique (up to an additive constant) height function corresponding to it.



Height Function

Pattern in X_{C_3}

Slopes

Slopes

- Given a height function h the **slope** in the direction e_i is given by

$$sl_{e_i}(h) := \lim_{n \rightarrow \infty} \frac{h(ne_i) - h(0)}{n}.$$

Slopes

- Given a height function h the **slope** in the direction e_i is given by

$$sl_{e_i}(h) := \lim_{n \rightarrow \infty} \frac{h(ne_i) - h(0)}{n}.$$

- It follows that $|sl_{e_i}(h)| \leq 1$ whenever it exists.

Slopes

- Given a height function h the **slope** in the direction e_i is given by

$$sl_{e_i}(h) := \lim_{n \rightarrow \infty} \frac{h(ne_i) - h(0)}{n}.$$

- It follows that $|sl_{e_i}(h)| \leq 1$ whenever it exists.
- Given a \mathbb{Z}^d -ergodic measure μ on X_{C_3} , the ergodic theorem implies that in every direction the slope exists and is a constant μ -almost everywhere.

Slopes

- Given a height function h the **slope** in the direction e_i is given by

$$sl_{e_i}(h) := \lim_{n \rightarrow \infty} \frac{h(ne_i) - h(0)}{n}.$$

- It follows that $|sl_{e_i}(h)| \leq 1$ whenever it exists.
- Given a \mathbb{Z}^d -ergodic measure μ on X_{C_3} , the ergodic theorem implies that in every direction the slope exists and is a constant μ -almost everywhere.
- The slope may be different in different directions.

Steep Slopes and 'Rigidity'

Steep Slopes and 'Rigidity'

- Suppose μ is a \mathbb{Z}^d -ergodic measure on X_{C_3} such that $|sl_{e_i}(h)| = 1$ almost everywhere.

Steep Slopes and 'Rigidity'

- Suppose μ is a \mathbb{Z}^d -ergodic measure on X_{C_3} such that $|sl_{e_i}(h)| = 1$ almost everywhere.
- Then

$$|h(j + ne_i) - h(j)| = n$$

almost everywhere for all $j \in \mathbb{Z}^d$.

Steep Slopes and 'Rigidity'

- Suppose μ is a \mathbb{Z}^d -ergodic measure on X_{C_3} such that $|sl_{e_i}(h)| = 1$ almost everywhere.
- Then

$$|h(j + ne_i) - h(j)| = n$$

almost everywhere for all $j \in \mathbb{Z}^d$.

- Thus for a height function h in the support of the measure, $h(j + ne_i)$ and $h(j)$ determine the values $h(j + te_1)$ for all $1 \leq t \leq n - 1$.

Steep Slopes and 'Rigidity'

- Suppose μ is a \mathbb{Z}^d -ergodic measure on X_{C_3} such that $|sl_{e_i}(h)| = 1$ almost everywhere.
- Then

$$|h(j + ne_i) - h(j)| = n$$

almost everywhere for all $j \in \mathbb{Z}^d$.

- Thus for a height function h in the support of the measure, $h(j + ne_i)$ and $h(j)$ determine the values $h(j + te_1)$ for all $1 \leq t \leq n - 1$.
- Let $X_{\text{frozen}} \subset X_{C_3}$ be the space of such configurations. Then $h_{\text{top}}(X_{\text{frozen}}) = 0$.

Steep Slopes and ‘Rigidity’

- Suppose μ is a \mathbb{Z}^d -ergodic measure on X_{C_3} such that $|sl_{e_i}(h)| = 1$ almost everywhere.
- Then

$$|h(j + ne_i) - h(j)| = n$$

almost everywhere for all $j \in \mathbb{Z}^d$.

- Thus for a height function h in the support of the measure, $h(j + ne_i)$ and $h(j)$ determine the values $h(j + te_1)$ for all $1 \leq t \leq n - 1$.
- Let $X_{frozen} \subset X_{C_3}$ be the space of such configurations. Then $h_{top}(X_{frozen}) = 0$.
- Thus slope 1 or -1 is ‘improbable’.

Gentle Slopes and 'Mixing'

Gentle Slopes and 'Mixing'

- Let h_1 be a height function on a ball D_n and

Gentle Slopes and 'Mixing'


- Let h_1 be a height function on a ball D_n and
- h_2 be a height function on \mathbb{Z}^d with slope s strictly between 1 and -1 in all directions.

Gentle Slopes and 'Mixing'

- Let h_1 be a height function on a ball D_n and
- h_2 be a height function on \mathbb{Z}^d with slope s strictly between 1 and -1 in all directions.
- Then we can choose an $N_n^s \in \mathbb{N}$ and a height function h which is equal to h_1 on D_n and h_2 on $D_{N_n^s}^c$.


Gentle Slopes and 'Mixing'

- Let h_1 be a height function on a ball D_n and
- h_2 be a height function on \mathbb{Z}^d with slope s strictly between 1 and -1 in all directions.
- Then we can choose an $N_n^s \in \mathbb{N}$ and a height function h which is equal to h_1 on D_n and h_2 on $D_{N_n^s}^c$.




Gentle Slopes and 'Mixing'

- Let h_1 be a height function on a ball D_n and
- h_2 be a height function on \mathbb{Z}^d with slope s strictly between 1 and -1 in all directions.
- Then we can choose an $N_n^s \in \mathbb{N}$ and a height function h which is equal to h_1 on D_n and h_2 on $D_{N_n^s}^c$.




Gentle Slopes and 'Mixing'

- Let h_1 be a height function on a ball D_n and
- h_2 be a height function on \mathbb{Z}^d with slope s strictly between 1 and -1 in all directions.
- Then we can choose an $N_n^s \in \mathbb{N}$ and a height function h which is equal to h_1 on D_n and h_2 on $D_{N_n^s}^c$.



Gentle Slopes and 'Mixing'

- Let h_1 be a height function on a ball D_n and
- h_2 be a height function on \mathbb{Z}^d with slope s strictly between 1 and -1 in all directions.
- Then we can choose an $N_n^s \in \mathbb{N}$ and a height function h which is equal to h_1 on D_n and h_2 on $D_{N_n^s}^c$.



Gentle Slopes and 'Mixing'

Thus if μ is a uniform Gibbs measure with slope between -1 and 1 then $\text{supp}(\mu) = X_{C_3}$.

Measures of Maximal Entropy

Measures of Maximal Entropy

- Since slopes 1 or -1 are 'improbable', the slope for any measure of maximal entropy is strictly between -1 and 1 .

Measures of Maximal Entropy

- Since slopes 1 or -1 are 'improbable', the slope for any measure of maximal entropy is strictly between -1 and 1 .
- For such 'gentle' slopes we just proved that these measures are fully supported.

Measures of Maximal Entropy

- Since slopes 1 or -1 are 'improbable', the slope for any measure of maximal entropy is strictly between -1 and 1 .
- For such 'gentle' slopes we just proved that these measures are fully supported.
- A shift space is entropy minimal if and only if every measure of maximal entropy is fully supported.

Measures of Maximal Entropy

- Since slopes 1 or -1 are 'improbable', the slope for any measure of maximal entropy is strictly between -1 and 1 .
- For such 'gentle' slopes we just proved that these measures are fully supported.
- A shift space is entropy minimal if and only if every measure of maximal entropy is fully supported.
- Therefore X_{C_3} is entropy minimal.

Measures of Maximal Entropy

- Since slopes 1 or -1 are 'improbable', the slope for any measure of maximal entropy is strictly between -1 and 1 .
- For such 'gentle' slopes we just proved that these measures are fully supported.
- A shift space is entropy minimal if and only if every measure of maximal entropy is fully supported.
- Therefore X_{C_3} is entropy minimal.
- What if C_3 is replaced by some other four-cycle free graph \mathcal{H} ?

Four-Cycle Free Graphs

Notice that we were using the following commutative diagram:

Four-Cycle Free Graphs

Notice that we were using the following commutative diagram:

$$\begin{array}{ccc} & \mathbb{Z} & \\ \textit{height function} \nearrow & \downarrow \text{mod } 3 & \\ \mathbb{Z}^d & \longrightarrow & C_3. \end{array}$$

Four-Cycle Free Graphs

Notice that we were using the following commutative diagram:

$$\begin{array}{ccc} & \mathbb{Z} & \\ \textit{height function} \nearrow & \downarrow \text{mod } 3 & \\ \mathbb{Z}^d & \longrightarrow & C_3. \end{array}$$

If C_3 is replaced by a connected four-cycle free graph \mathcal{H}

Four-Cycle Free Graphs

Notice that we were using the following commutative diagram:

$$\begin{array}{ccc} & & \mathbb{Z} \\ \textit{height function} \nearrow & & \downarrow \text{mod } 3 \\ \mathbb{Z}^d & \longrightarrow & C_3. \end{array}$$

If C_3 is replaced by a connected four-cycle free graph \mathcal{H} then \mathbb{Z} is replaced by the universal cover of \mathcal{H} .


Four-Cycle Free Graphs

Notice that we were using the following commutative diagram:

$$\begin{array}{ccc} & \mathbb{Z} & \\ \text{height function} \nearrow & \downarrow \text{mod } 3 & \\ \mathbb{Z}^d & \longrightarrow & C_3. \end{array}$$

If C_3 is replaced by a connected four-cycle free graph \mathcal{H} then \mathbb{Z} is replaced by the universal cover of \mathcal{H} .

Conjecture: For $d = 2$, $X_{\mathcal{H}}$ is entropy minimal for all connected graphs \mathcal{H} .



Thank You!