

# Four-Cycle Free Graphs and Entropy Minimality

Nishant Chandgotia

University of British Columbia

July, 2014

# Outline

- Entropy Minimality and Hom Shifts
- Mixing Conditions and Entropy Minimality
- Rigidity and Flexibility in the Space of 3-Colourings.

# Hom shifts

Given a finite undirected graph  $\mathcal{H}$  without multiple edges,

# Hom shifts

Given a finite undirected graph  $\mathcal{H}$  without multiple edges, we can define a shift space  $X_{\mathcal{H}} \subset \mathcal{H}^{\mathbb{Z}^d}$

## Hom shifts

Given a finite undirected graph  $\mathcal{H}$  without multiple edges, we can define a shift space  $X_{\mathcal{H}} \subset \mathcal{H}^{\mathbb{Z}^d}$  such that symbols on adjacent vertices of  $\mathbb{Z}^d$  form an edge in the graph  $\mathcal{H}$ .

## Hom shifts

Given a finite undirected graph  $\mathcal{H}$  without multiple edges, we can define a shift space  $X_{\mathcal{H}} \subset \mathcal{H}^{\mathbb{Z}^d}$  such that symbols on adjacent vertices of  $\mathbb{Z}^d$  form an edge in the graph  $\mathcal{H}$ .

Here every direction has the same constraint.

# Hom shifts

Given a finite undirected graph  $\mathcal{H}$  without multiple edges, we can define a shift space  $X_{\mathcal{H}} \subset \mathcal{H}^{\mathbb{Z}^d}$  such that symbols on adjacent vertices of  $\mathbb{Z}^d$  form an edge in the graph  $\mathcal{H}$ .

Here every direction has the same constraint.


**Examples:**

# Hom shifts

Given a finite undirected graph  $\mathcal{H}$  without multiple edges, we can define a shift space  $X_{\mathcal{H}} \subset \mathcal{H}^{\mathbb{Z}^d}$  such that symbols on adjacent vertices of  $\mathbb{Z}^d$  form an edge in the graph  $\mathcal{H}$ .

Here every direction has the same constraint.

**Examples:**(Hard Square model)



Graph  $\mathcal{H}$

1	0	0	0	0
0	0	0	0	0
1	0	1	0	0
0	0	0	1	0
0	1	0	0	0

A Pattern




# Hom shifts


Given a finite undirected graph  $\mathcal{H}$  without multiple edges, we can define a shift space  $X_{\mathcal{H}} \subset \mathcal{H}^{\mathbb{Z}^d}$  such that symbols on adjacent vertices of  $\mathbb{Z}^d$  form an edge in the graph  $\mathcal{H}$ .

Here every direction has the same constraint.

**Examples:**(3-colourings)



Graph  $\mathcal{H}$



A Pattern

# Entropy

# Entropy

The **language**  $\mathcal{B}(X)$  of a shift space is the set of patterns appearing in elements of  $X$ .

# Entropy

The **language**  $\mathcal{B}(X)$  of a shift space is the set of patterns appearing in elements of  $X$ .

The **topological entropy** of  $X$  is defined as

# Entropy

The **language**  $\mathcal{B}(X)$  of a shift space is the set of patterns appearing in elements of  $X$ .

The **topological entropy** of  $X$  is defined as

$$h_{top}(X) := \lim_{n \rightarrow \infty} \frac{\log |\mathcal{B}(X) \cap \mathcal{A}^{\{1,2,\dots,n\}^d}|}{n^d}.$$

## Entropy Minimality

# Entropy Minimality

If  $Y \subset X$  are shift spaces then

$$h_{top}(Y) \leq h_{top}(X).$$

# Entropy Minimality

If  $Y \subset X$  are shift spaces then

$$h_{top}(Y) \leq h_{top}(X).$$

A shift space  $X$  is said to be **entropy minimal** if for all shift spaces  $Y \subsetneq X$ ,



# Entropy Minimality

If  $Y \subset X$  are shift spaces then

$$h_{top}(Y) \leq h_{top}(X).$$

A shift space  $X$  is said to be **entropy minimal** if for all shift spaces  $Y \subsetneq X$ ,  $h_{top}(Y) < h_{top}(X)$ ,

# Entropy Minimality

If  $Y \subset X$  are shift spaces then

$$h_{top}(Y) \leq h_{top}(X).$$

A shift space  $X$  is said to be **entropy minimal** if for all shift spaces  $Y \subsetneq X$ ,  $h_{top}(Y) < h_{top}(X)$ , that is, if we forbid any pattern from the language  $\mathcal{B}(X)$  the entropy will drop.

# Entropy Minimality

If  $Y \subset X$  are shift spaces then

$$h_{top}(Y) \leq h_{top}(X).$$

A shift space  $X$  is said to be **entropy minimal** if for all shift spaces  $Y \subsetneq X$ ,  $h_{top}(Y) < h_{top}(X)$ , that is, if we forbid any pattern from the language  $\mathcal{B}(X)$  the entropy will drop.

If  $h_{top}(X) = 0$  then  $X$  is entropy minimal if and only if it is minimal.

# Entropy Minimality

If  $Y \subset X$  are shift spaces then

$$h_{top}(Y) \leq h_{top}(X).$$

A shift space  $X$  is said to be **entropy minimal** if for all shift spaces  $Y \subsetneq X$ ,  $h_{top}(Y) < h_{top}(X)$ , that is, if we forbid any pattern from the language  $\mathcal{B}(X)$  the entropy will drop.

If  $h_{top}(X) = 0$  then  $X$  is entropy minimal if and only if it is minimal.

(*Quas and Trow '00*) Every shift space  $X$  contains an entropy minimal shift space  $Y \subset X$  such that  $h_{top}(X) = h_{top}(Y)$ .

## Main Result

# Main Result

Theorem (Chandgotia, Meyerovitch '13)

*Let  $C_n$  be an  $n$ -cycle for some integer  $n \neq 4$ . Then  $X_{C_n}$  is entropy minimal.*

# Main Result

Theorem (Chandgotia, Meyerovitch '13)

*Let  $C_n$  be an  $n$ -cycle for some integer  $n \neq 4$ . Then  $X_{C_n}$  is entropy minimal.*

Theorem (Chandgotia '14; In preparation)

*If  $\mathcal{H}$  is a connected graph without 4-cycles then  $X_{\mathcal{H}}$  is entropy minimal.*

# Main Result

Theorem (Chandgotia, Meyerovitch '13)

*Let  $C_n$  be an  $n$ -cycle for some integer  $n \neq 4$ . Then  $X_{C_n}$  is entropy minimal.*

Theorem (Chandgotia '14; In preparation)

*If  $\mathcal{H}$  is a connected graph without 4-cycles then  $X_{\mathcal{H}}$  is entropy minimal.*

**Remark:** We will concentrate on  $X_{C_3}$ , the space of all 3-colourings.



## Previous Results

## Previous Results

(Coven and Smítal '93) If a shift space is entropy minimal then it is topologically transitive.

## Previous Results

(*Coven and Smítal '93*) If a shift space is entropy minimal then it is topologically transitive.

(*Boyle, Pavlov and Schraudner '09*) There exists a block-gluing shift space which is not entropy minimal.

## Previous Results

(Coven and Smítal '93) If a shift space is entropy minimal then it is topologically transitive.

(Boyle, Pavlov and Schraudner '09) There exists a block-gluing shift space which is not entropy minimal.

(Schraudner '09) Every strongly irreducible shift space is entropy minimal.

## Previous Results


(Coven and Smítal '93) If a shift space is entropy minimal then it is topologically transitive.

(Boyle, Pavlov and Schraudner '09) There exists a block-gluing shift space which is not entropy minimal.

(Schraudner '09) Every strongly irreducible shift space is entropy minimal.

(Lightwood and Schraudner '12) A shift of finite type is entropy minimal if and only if the set of all 'non-universal' boundary patterns is 'poor'.

# Mixing Conditions and Entropy Minimality



A Few Key Ideas: Height Functions and  $X_{C_3}$

## A Few Key Ideas: Height Functions and $X_{C_3}$

A **height function** is a function  $h : \mathbb{Z}^d \longrightarrow \mathbb{Z}$  such that

$$|h(v) - h(w)| = 1$$

for all adjacent vertices  $v$  and  $w$ .



## A Few Key Ideas: Height Functions and $X_{C_3}$

A **height function** is a function  $h : \mathbb{Z}^d \longrightarrow \mathbb{Z}$  such that

$$|h(v) - h(w)| = 1$$

for all adjacent vertices  $v$  and  $w$ .

If  $h$  is a height function then  $h \bmod 3$  is an element of  $X_{C_3}$ .

## A Few Key Ideas: Height Functions and $X_{C_3}$

A **height function** is a function  $h : \mathbb{Z}^d \longrightarrow \mathbb{Z}$  such that

$$|h(v) - h(w)| = 1$$

for all adjacent vertices  $v$  and  $w$ .

If  $h$  is a height function then  $h \bmod 3$  is an element of  $X_{C_3}$ . This sets up a correspondence between height functions and elements of  $X_{C_3}$ .


## A Few Key Ideas: Height Functions and $X_{C_3}$

A **height function** is a function  $h : \mathbb{Z}^d \rightarrow \mathbb{Z}$  such that

$$|h(v) - h(w)| = 1$$

for all adjacent vertices  $v$  and  $w$ .

If  $h$  is a height function then  $h \bmod 3$  is an element of  $X_{C_3}$ . This sets up a correspondence between height functions and elements of  $X_{C_3}$ .



Height Function


## A Few Key Ideas: Height Functions and $X_{C_3}$

A **height function** is a function  $h : \mathbb{Z}^d \rightarrow \mathbb{Z}$  such that

$$|h(v) - h(w)| = 1$$

for all adjacent vertices  $v$  and  $w$ .

If  $h$  is a height function then  $h \bmod 3$  is an element of  $X_{C_3}$ . This sets up a correspondence between height functions and elements of  $X_{C_3}$ .



Height Function

Pattern in  $X_{C_3}$

## A Few Key Ideas: Slopes

## A Few Key Ideas: Slopes

Given a height function  $h$

## A Few Key Ideas: Slopes

Given a height function  $h$  the **slope** in the direction  $e_i$  is given by

## A Few Key Ideas: Slopes

Given a height function  $h$  the **slope** in the direction  $e_i$  is given by

$$\lim_{n \rightarrow \infty} \frac{h(ne_i) - h(0)}{n}.$$



## A Few Key Ideas: Slopes

Given a height function  $h$  the **slope** in the direction  $e_i$  is given by

$$\lim_{n \rightarrow \infty} \frac{h(ne_i) - h(0)}{n}.$$

The slope is always between  $-1$  and  $1$  in every direction.

## A Few Key Ideas: Slopes

Given a height function  $h$  the **slope** in the direction  $e_i$  is given by

$$\lim_{n \rightarrow \infty} \frac{h(ne_i) - h(0)}{n}.$$

The slope is always between  $-1$  and  $1$  in every direction.

Given an ergodic measure  $\mu$  on  $X_{C_3}$ ,

## A Few Key Ideas: Slopes

Given a height function  $h$  the **slope** in the direction  $e_i$  is given by

$$\lim_{n \rightarrow \infty} \frac{h(ne_i) - h(0)}{n}.$$

The slope is always between  $-1$  and  $1$  in every direction.

Given an ergodic measure  $\mu$  on  $X_{C_3}$ , the slope is a constant  $\mu$ -almost everywhere in every direction.

## A Few Key Ideas: Slopes

Given a height function  $h$  the **slope** in the direction  $e_i$  is given by

$$\lim_{n \rightarrow \infty} \frac{h(ne_i) - h(0)}{n}.$$

The slope is always between  $-1$  and  $1$  in every direction.

Given an ergodic measure  $\mu$  on  $X_{C_3}$ , the slope is a constant  $\mu$ -almost everywhere in every direction.

Note that the slope may be different in different directions.

## A Few Key Ideas: Slopes

Given a height function  $h$  the **slope** in the direction  $e_i$  is given by

$$\lim_{n \rightarrow \infty} \frac{h(ne_i) - h(0)}{n}.$$

The slope is always between  $-1$  and  $1$  in every direction.

Given an ergodic measure  $\mu$  on  $X_{C_3}$ , the slope is a constant  $\mu$ -almost everywhere in every direction.

Note that the slope may be different in different directions.

A Few Key Ideas: Steep Slopes and Rigidity

## A Few Key Ideas: Steep Slopes and Rigidity

If the slope of a height function is 1 or  $-1$  in some direction

## A Few Key Ideas: Steep Slopes and Rigidity

If the slope of a height function is 1 or  $-1$  in some direction then it cannot be changed anywhere to obtain another height function.



## A Few Key Ideas: Steep Slopes and Rigidity

If the slope of a height function is 1 or  $-1$  in some direction then it cannot be changed anywhere to obtain another height function.

0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2

## A Few Key Ideas: Steep Slopes and Rigidity

If the slope of a height function is 1 or  $-1$  in some direction then it cannot be changed anywhere to obtain another height function.

0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2

Let  $X_{\text{frozen}}$  be the space of such configurations. Then 
$$h_{\text{top}}(X_{\text{frozen}}) = 0.$$

## A Few Key Ideas: Steep Slopes and Rigidity

If the slope of a height function is 1 or  $-1$  in some direction then it cannot be changed anywhere to obtain another height function.

0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2

Let  $X_{\text{frozen}}$  be the space of such configurations. Then  $h_{\text{top}}(X_{\text{frozen}}) = 0$ . Thus slope 1 or  $-1$  is 'improbable'.

A Few Key Ideas: Gentle Slopes and Flexibility

## A Few Key Ideas: Gentle Slopes and Flexibility

Given any height function  $h_1$  on a ball  $D_n$  in  $\mathbb{Z}^d$

## A Few Key Ideas: Gentle Slopes and Flexibility

Given any height function  $h_1$  on a ball  $D_n$  in  $\mathbb{Z}^d$  and a height function  $h_2$  on  $\mathbb{Z}^d$  with slope  $s$  strictly between 1 and  $-1$  in all directions

## A Few Key Ideas: Gentle Slopes and Flexibility

Given any height function  $h_1$  on a ball  $D_n$  in  $\mathbb{Z}^d$  and a height function  $h_2$  on  $\mathbb{Z}^d$  with slope  $s$  strictly between 1 and  $-1$  in all directions we can choose an  $N_n^s \in \mathbb{N}$  and a height function  $h$

## A Few Key Ideas: Gentle Slopes and Flexibility

Given any height function  $h_1$  on a ball  $D_n$  in  $\mathbb{Z}^d$  and a height function  $h_2$  on  $\mathbb{Z}^d$  with slope  $s$  strictly between 1 and  $-1$  in all directions we can choose an  $N_n^s \in \mathbb{N}$  and a height function  $h$  which is equal to  $h_1$  on  $D_n$  and  $h_2$  on  $D_{N_n^s}^c$ .




## A Few Key Ideas: Gentle Slopes and Flexibility

Given any height function  $h_1$  on a ball  $D_n$  in  $\mathbb{Z}^d$  and a height function  $h_2$  on  $\mathbb{Z}^d$  with slope  $s$  strictly between 1 and  $-1$  in all directions we can choose an  $N_n^s \in \mathbb{N}$  and a height function  $h$  which is equal to  $h_1$  on  $D_n$  and  $h_2$  on  $D_{N_n^s}^c$ .


## A Few Key Ideas: Gentle Slopes and Flexibility

Given any height function  $h_1$  on a ball  $D_n$  in  $\mathbb{Z}^d$  and a height function  $h_2$  on  $\mathbb{Z}^d$  with slope  $s$  strictly between 1 and  $-1$  in all directions we can choose an  $N_n^s \in \mathbb{N}$  and a height function  $h$  which is equal to  $h_1$  on  $D_n$  and  $h_2$  on  $D_{N_n^s}^c$ .




## A Few Key Ideas: Gentle Slopes and Flexibility

Given any height function  $h_1$  on a ball  $D_n$  in  $\mathbb{Z}^d$  and a height function  $h_2$  on  $\mathbb{Z}^d$  with slope  $s$  strictly between 1 and  $-1$  in all directions we can choose an  $N_n^s \in \mathbb{N}$  and a height function  $h$  which is equal to  $h_1$  on  $D_n$  and  $h_2$  on  $D_{N_n^s}^c$ .




## A Few Key Ideas: Gentle Slopes and Flexibility

Given any height function  $h_1$  on a ball  $D_n$  in  $\mathbb{Z}^d$  and a height function  $h_2$  on  $\mathbb{Z}^d$  with slope  $s$  strictly between 1 and  $-1$  in all directions we can choose an  $N_n^s \in \mathbb{N}$  and a height function  $h$  which is equal to  $h_1$  on  $D_n$  and  $h_2$  on  $D_{N_n^s}^c$ .



## A Few Key Ideas: Gentle Slopes and Flexibility

Given any height function  $h_1$  on a ball  $D_n$  in  $\mathbb{Z}^d$  and a height function  $h_2$  on  $\mathbb{Z}^d$  with slope  $s$  strictly between 1 and  $-1$  in all directions we can choose an  $N_n^s \in \mathbb{N}$  and a height function  $h$  which is equal to  $h_1$  on  $D_n$  and  $h_2$  on  $D_{N_n^s}^c$ .



## A Few Key Ideas: Measures of Maximal Entropy

## A Few Key Ideas: Measures of Maximal Entropy

Since slopes 1 or  $-1$  are 'improbable', the slope of any measure of maximal entropy is strictly between  $-1$  and  $1$ .

## A Few Key Ideas: Measures of Maximal Entropy

Since slopes 1 or  $-1$  are 'improbable', the slope of any measure of maximal entropy is strictly between  $-1$  and  $1$ . Such measures are fully supported.



## A Few Key Ideas: Measures of Maximal Entropy

Since slopes 1 or  $-1$  are 'improbable', the slope of any measure of maximal entropy is strictly between  $-1$  and  $1$ . Such measures are fully supported. A shift space is entropy minimal if and only if every measure of maximal entropy is fully supported.

## A Few Key Ideas: Measures of Maximal Entropy

Since slopes 1 or  $-1$  are 'improbable', the slope of any measure of maximal entropy is strictly between  $-1$  and  $1$ . Such measures are fully supported. A shift space is entropy minimal if and only if every measure of maximal entropy is fully supported. Therefore  $X_{C_3}$  is entropy minimal.

## A Few Key Ideas: Measures of Maximal Entropy

Since slopes 1 or  $-1$  are 'improbable', the slope of any measure of maximal entropy is strictly between  $-1$  and  $1$ . Such measures are fully supported. A shift space is entropy minimal if and only if every measure of maximal entropy is fully supported. Therefore  $X_{C_3}$  is entropy minimal.

What if  $C_3$  is replaced by another connected four-cycle free graph  $\mathcal{H}$ ?

## A Few Key Ideas: Measures of Maximal Entropy

Since slopes 1 or  $-1$  are 'improbable', the slope of any measure of maximal entropy is strictly between  $-1$  and  $1$ . Such measures are fully supported. A shift space is entropy minimal if and only if every measure of maximal entropy is fully supported. Therefore  $X_{C_3}$  is entropy minimal.

What if  $C_3$  is replaced by another connected four-cycle free graph  $\mathcal{H}$ ? Notice that we were using the following commutative diagram:

## A Few Key Ideas: Measures of Maximal Entropy

Since slopes 1 or  $-1$  are 'improbable', the slope of any measure of maximal entropy is strictly between  $-1$  and  $1$ . Such measures are fully supported. A shift space is entropy minimal if and only if every measure of maximal entropy is fully supported. Therefore  $X_{C_3}$  is entropy minimal.

What if  $C_3$  is replaced by another connected four-cycle free graph  $\mathcal{H}$ ? Notice that we were using the following commutative diagram:

$$\begin{array}{ccc} & & \mathbb{Z} \\ \text{height function} \nearrow & & \downarrow \text{mod } 3 \\ \mathbb{Z}^d & \longrightarrow & C_3. \end{array}$$

## A Few Key Ideas: Measures of Maximal Entropy

Since slopes 1 or  $-1$  are ‘improbable’, the slope of any measure of maximal entropy is strictly between  $-1$  and  $1$ . Such measures are fully supported. A shift space is entropy minimal if and only if every measure of maximal entropy is fully supported. Therefore  $X_{C_3}$  is entropy minimal.

What if  $C_3$  is replaced by another connected four-cycle free graph  $\mathcal{H}$ ? Notice that we were using the following commutative diagram:

$$\begin{array}{ccc} & & \mathbb{Z} \\ \text{height function} \nearrow & & \downarrow \text{mod } 3 \\ \mathbb{Z}^d & \longrightarrow & C_3. \end{array}$$

$\mathbb{Z}$  is replaced by the universal cover of  $\mathcal{H}$ .




**Conjecture:** If  $\mathcal{H}$  is any connected graph then  $X_{\mathcal{H}}$  is entropy minimal. (d=2)



**Conjecture:** If  $\mathcal{H}$  is any connected graph then  $X_{\mathcal{H}}$  is entropy minimal. (d=2)

**Question:** What shift spaces are conjugate to  $X_{\mathcal{H}}$  for some graph  $\mathcal{H}$ ?



Thank You!