

Graph Foldings and Markov Random Fields

Nishant Chandgotia

University of British Columbia

August, 2014

Outline

- Markov random fields and Gibbs measures with nearest neighbour interactions
- Review of previous results
- The pivot property
- Graph folding and Hammersly-Clifford spaces

Some Notation and Setting

Consider a countable locally-finite undirected graph $\mathcal{G} = (\mathcal{V}_{\mathcal{G}}, \mathcal{E}_{\mathcal{G}})$ and \mathfrak{A} , a finite set.

Some Notation and Setting

Consider a countable locally-finite undirected graph $\mathcal{G} = (\mathcal{V}_{\mathcal{G}}, \mathcal{E}_{\mathcal{G}})$ and \mathcal{A} , a finite set. Suppose $X \subset \mathcal{A}^{\mathcal{V}_{\mathcal{G}}}$ is a closed set.

Some Notation and Setting

Consider a countable locally-finite undirected graph $\mathcal{G} = (\mathcal{V}_{\mathcal{G}}, \mathcal{E}_{\mathcal{G}})$ and \mathfrak{A} , a finite set. Suppose $X \subset \mathfrak{A}^{\mathcal{V}_{\mathcal{G}}}$ is a closed set. For a finite set $A \subset \mathcal{V}_{\mathcal{G}}$ and a pattern $a : A \longrightarrow \mathfrak{A}$, $[a]_A$ denotes a **cylinder set** in X and is given by

$$[a]_A = \{x \in X \mid x|_A = a\}.$$

Some Notation and Setting

Consider a countable locally-finite undirected graph $\mathcal{G} = (\mathcal{V}_{\mathcal{G}}, \mathcal{E}_{\mathcal{G}})$ and \mathfrak{A} , a finite set. Suppose $X \subset \mathfrak{A}^{\mathcal{V}_{\mathcal{G}}}$ is a closed set. For a finite set $A \subset \mathcal{V}_{\mathcal{G}}$ and a pattern $a : A \longrightarrow \mathfrak{A}$, $[a]_A$ denotes a **cylinder set** in X and is given by

$$[a]_A = \{x \in X \mid x|_A = a\}.$$

The **boundary** of a set A is denoted by ∂A and is given by

$$\partial A = \{v \in A^c \mid v \sim w \in A\}.$$


Some Notation and Setting

Consider a countable locally-finite undirected graph $\mathcal{G} = (\mathcal{V}_{\mathcal{G}}, \mathcal{E}_{\mathcal{G}})$ and \mathfrak{A} , a finite set. Suppose $X \subset \mathfrak{A}^{\mathcal{V}_{\mathcal{G}}}$ is a closed set. For a finite set $A \subset \mathcal{V}_{\mathcal{G}}$ and a pattern $a : A \rightarrow \mathfrak{A}$, $[a]_A$ denotes a **cylinder set** in X and is given by

$$[a]_A = \{x \in X \mid x|_A = a\}.$$

The **boundary** of a set A is denoted by ∂A and is given by

$$\partial A = \{v \in A^c \mid v \sim w \in A\}.$$




Some Notation and Setting

Consider a countable locally-finite undirected graph $\mathcal{G} = (\mathcal{V}_{\mathcal{G}}, \mathcal{E}_{\mathcal{G}})$ and \mathfrak{A} , a finite set. Suppose $X \subset \mathfrak{A}^{\mathcal{V}_{\mathcal{G}}}$ is a closed set. For a finite set $A \subset \mathcal{V}_{\mathcal{G}}$ and a pattern $a : A \rightarrow \mathfrak{A}$, $[a]_A$ denotes a **cylinder set** in X and is given by

$$[a]_A = \{x \in X \mid x|_A = a\}.$$

The **boundary** of a set A is denoted by ∂A and is given by

$$\partial A = \{v \in A^c \mid v \sim w \in A\}.$$



- –Elements of A


Some Notation and Setting

Consider a countable locally-finite undirected graph $\mathcal{G} = (\mathcal{V}_{\mathcal{G}}, \mathcal{E}_{\mathcal{G}})$ and \mathfrak{A} , a finite set. Suppose $X \subset \mathfrak{A}^{\mathcal{V}_{\mathcal{G}}}$ is a closed set. For a finite set $A \subset \mathcal{V}_{\mathcal{G}}$ and a pattern $a : A \longrightarrow \mathfrak{A}$, $[a]_A$ denotes a **cylinder set** in X and is given by

$$[a]_A = \{x \in X \mid x|_A = a\}.$$

The **boundary** of a set A is denoted by ∂A and is given by

$$\partial A = \{v \in A^c \mid v \sim w \in A\}.$$



- –Elements of A
- –Elements of the boundary of A


Some Notation and Setting

Consider a countable locally-finite undirected graph $\mathcal{G} = (\mathcal{V}_{\mathcal{G}}, \mathcal{E}_{\mathcal{G}})$ and \mathfrak{A} , a finite set. Suppose $X \subset \mathfrak{A}^{\mathcal{V}_{\mathcal{G}}}$ is a closed set. For a finite set $A \subset \mathcal{V}_{\mathcal{G}}$ and a pattern $a : A \longrightarrow \mathfrak{A}$, $[a]_A$ denotes a **cylinder set** in X and is given by

$$[a]_A = \{x \in X \mid x|_A = a\}.$$

The **boundary** of a set A is denoted by ∂A and is given by

$$\partial A = \{v \in A^c \mid v \sim w \in A\}.$$



- –Elements of A
- –Elements of the boundary of A

Homomorphism Spaces

For simplification, we will often refer to homomorphism spaces.

Homomorphism Spaces

For simplification, we will often refer to homomorphism spaces.
Consider a finite undirected graph $\mathcal{H} = (\mathcal{V}_{\mathcal{H}}, \mathcal{E}_{\mathcal{H}})$

Homomorphism Spaces

For simplification, we will often refer to homomorphism spaces. Consider a finite undirected graph $\mathcal{H} = (\mathcal{V}_{\mathcal{H}}, \mathcal{E}_{\mathcal{H}})$ and fix a bipartite graph \mathcal{G} .

Homomorphism Spaces

For simplification, we will often refer to homomorphism spaces. Consider a finite undirected graph $\mathcal{H} = (\mathcal{V}_{\mathcal{H}}, \mathcal{E}_{\mathcal{H}})$ and fix a bipartite graph \mathcal{G} . Let $X = \text{Hom}(\mathcal{G}, \mathcal{H})$ represent the space of all graph homomorphisms from \mathcal{G} to \mathcal{H} .

Homomorphism Spaces

For simplification, we will often refer to homomorphism spaces. Consider a finite undirected graph $\mathcal{H} = (\mathcal{V}_{\mathcal{H}}, \mathcal{E}_{\mathcal{H}})$ and fix a bipartite graph \mathcal{G} . Let $X = \text{Hom}(\mathcal{G}, \mathcal{H})$ represent the space of all graph homomorphisms from \mathcal{G} to \mathcal{H} .

Examples:

Homomorphism Spaces


For simplification, we will often refer to homomorphism spaces. Consider a finite undirected graph $\mathcal{H} = (\mathcal{V}_{\mathcal{H}}, \mathcal{E}_{\mathcal{H}})$ and fix a bipartite graph \mathcal{G} . Let $X = \text{Hom}(\mathcal{G}, \mathcal{H})$ represent the space of all graph homomorphisms from \mathcal{G} to \mathcal{H} .

Examples:(Hard Square model)

Homomorphism Spaces

For simplification, we will often refer to homomorphism spaces. Consider a finite undirected graph $\mathcal{H} = (\mathcal{V}_{\mathcal{H}}, \mathcal{E}_{\mathcal{H}})$ and fix a bipartite graph \mathcal{G} . Let $X = \text{Hom}(\mathcal{G}, \mathcal{H})$ represent the space of all graph homomorphisms from \mathcal{G} to \mathcal{H} .

Examples:(Hard Square model)



Graph \mathcal{H}

1	0	0	0	0
0	0	0	0	0
1	0	1	0	0
0	0	0	1	0
0	1	0	0	0

An element of
 $\text{Hom}(\mathcal{G}, \mathcal{H})$

Homomorphism Spaces


For simplification, we will often refer to homomorphism spaces. Consider a finite undirected graph $\mathcal{H} = (\mathcal{V}_{\mathcal{H}}, \mathcal{E}_{\mathcal{H}})$ and fix a bipartite graph \mathcal{G} . Let $X = \text{Hom}(\mathcal{G}, \mathcal{H})$ represent the space of all graph homomorphisms from \mathcal{G} to \mathcal{H} .

Examples:(3-colourings of a graph)


Homomorphism Spaces

For simplification, we will often refer to homomorphism spaces. Consider a finite undirected graph $\mathcal{H} = (\mathcal{V}_{\mathcal{H}}, \mathcal{E}_{\mathcal{H}})$ and fix a bipartite graph \mathcal{G} . Let $X = \text{Hom}(\mathcal{G}, \mathcal{H})$ represent the space of all graph homomorphisms from \mathcal{G} to \mathcal{H} .

Examples:(3-colourings of a graph)



Graph \mathcal{H}



An element of
 $\text{Hom}(\mathcal{G}, \mathcal{H})$

Safe Symbol


Safe Symbol

The space $Hom(\mathcal{G}, \mathcal{H})$ is said to have a **safe symbol** \star if there exists a vertex $\star \in \mathcal{V}_{\mathcal{H}}$ such that for all vertices $v \in \mathcal{V}_{\mathcal{H}}$ the vertex $v \sim \star$.

Safe Symbol

The space $\text{Hom}(\mathcal{G}, \mathcal{H})$ is said to have a **safe symbol** \star if there exists a vertex $\star \in \mathcal{V}_{\mathcal{H}}$ such that for all vertices $v \in \mathcal{V}_{\mathcal{H}}$ the vertex $v \sim \star$.


For instance, 0 is a safe symbol for the hard square model



Safe Symbol

The space $\text{Hom}(\mathcal{G}, \mathcal{H})$ is said to have a **safe symbol** \star if there exists a vertex $\star \in \mathcal{V}_{\mathcal{H}}$ such that for all vertices $v \in \mathcal{V}_{\mathcal{H}}$ the vertex $v \sim \star$.

For instance, 0 is a safe symbol for the hard square model but the space of 3-colourings of a graph does not have any safe symbol.



Markov Random Fields

A **Markov random field** is a probability measure μ on $\mathcal{A}^{\mathcal{V}_G}$

Markov Random Fields

A **Markov random field** is a probability measure μ on $\mathcal{A}^{\mathcal{V}_G}$ with the property that for all finite $A, B \subset \mathcal{V}_G$

Markov Random Fields

A **Markov random field** is a probability measure μ on $\mathcal{A}^{\mathcal{V}_{\mathcal{G}}}$ with the property that for all finite $A, B \subset \mathcal{V}_{\mathcal{G}}$ such that $\partial A \subset B \subset A^c$ and

Markov Random Fields

A **Markov random field** is a probability measure μ on $\mathcal{A}^{\mathcal{V}_G}$ with the property that for all finite $A, B \subset \mathcal{V}_G$ such that $\partial A \subset B \subset A^c$ and $a \in \mathfrak{A}^A, b \in \mathfrak{A}^B$ satisfying $\mu([b]_B) > 0$

Markov Random Fields


A **Markov random field** is a probability measure μ on $\mathcal{A}^{\mathcal{V}_G}$ with the property that for all finite $A, B \subset \mathcal{V}_G$ such that $\partial A \subset B \subset A^c$ and $a \in \mathfrak{A}^A, b \in \mathfrak{A}^B$ satisfying $\mu([b]_B) > 0$

$$\mu([a]_A \mid [b]_B) = \mu([a]_A \mid [b]_{\partial A}).$$

Markov Random Fields

A **Markov random field** is a probability measure μ on $\mathcal{A}^{\mathcal{V}_G}$ with the property that for all finite $A, B \subset \mathcal{V}_G$ such that $\partial A \subset B \subset A^c$ and $a \in \mathcal{A}^A, b \in \mathcal{A}^B$ satisfying $\mu([b]_B) > 0$

$$\mu([a]_A \mid [b]_B) = \mu([a]_A \mid [b]_{\partial A}).$$




- –Elements of A
- –Elements of B
- –Elements of the boundary of A

Markov Random Fields

A **Markov random field** is a probability measure μ on $\mathcal{A}^{\mathcal{V}_G}$ with the property that for all finite $A, B \subset \mathcal{V}_G$ such that $\partial A \subset B \subset A^c$ and $a \in \mathfrak{A}^A, b \in \mathfrak{A}^B$ satisfying $\mu([b]_B) > 0$

$$\mu([a]_A \mid [b]_B) = \mu([a]_A \mid [b]_{\partial A}).$$




- –Elements of A
- –Elements of B
- –Elements of the boundary of A

The set of conditional measures $\mu([\cdot]_A \mid [b]_{\partial A})$ for all $A \subset \mathcal{V}_G$ finite and $b \in \mathfrak{A}^{\partial A}$ is called **specification** for the measure μ .

Markov Random Fields

A **Markov random field** is a probability measure μ on $\mathcal{A}^{\mathcal{V}_G}$ with the property that for all finite $A, B \subset \mathcal{V}_G$ such that $\partial A \subset B \subset A^c$ and $a \in \mathfrak{A}^A, b \in \mathfrak{A}^B$ satisfying $\mu([b]_B) > 0$

$$\mu([a]_A \mid [b]_B) = \mu([a]_A \mid [b]_{\partial A}).$$



- –Elements of A
- –Elements of B
- –Elements of the boundary of A

The set of conditional measures $\mu([\cdot]_A \mid [b]_{\partial A})$ for all $A \subset \mathcal{V}_G$ finite and $b \in \mathfrak{A}^{\partial A}$ is called **specification** for the measure μ . It might not have any finite parametrisation.

Gibbs Measures

Given $X \subset \mathfrak{A}^{\mathcal{V}_G}$

Gibbs Measures

Given $X \subset \mathfrak{A}^{\mathcal{V}_G}$ we define a nearest neighbour interaction on X

Gibbs Measures

Given $X \in \mathbb{A}^{\mathcal{V}_G}$ we define a **nearest neighbour interaction** on X as a function $V : \{[a]_A \mid A \subset \mathcal{V}_G \text{ finite}\} \longrightarrow \mathbb{R}$

Gibbs Measures

Given $X \in \mathcal{A}^{\mathcal{V}_{\mathcal{G}}}$ we define a **nearest neighbour interaction** on X as a function $V : \{[a]_A \mid A \subset \mathcal{V}_{\mathcal{G}} \text{ finite}\} \rightarrow \mathbb{R}$ which is supported on patterns on the edges and the vertices of \mathcal{G} .

Gibbs Measures

Given $X \in \mathfrak{A}^{\mathcal{V}_{\mathcal{G}}}$ we define a **nearest neighbour interaction** on X as a function $V : \{[a]_A \mid A \subset \mathcal{V}_{\mathcal{G}} \text{ finite}\} \longrightarrow \mathbb{R}$ which is supported on patterns on the edges and the vertices of \mathcal{G} .

A **Gibbs state with a nearest neighbor interaction V** is a Markov random field μ such that for all $x \in \text{supp}(\mu)$ and $A, B \subset \mathcal{V}_{\mathcal{G}}$ finite satisfying $\partial A \subset B \subset A^c$

Gibbs Measures

Given $X \subset \mathcal{A}^{\mathcal{V}_{\mathcal{G}}}$ we define a **nearest neighbour interaction** on X as a function $V : \{[a]_A \mid A \subset \mathcal{V}_{\mathcal{G}} \text{ finite}\} \rightarrow \mathbb{R}$ which is supported on patterns on the edges and the vertices of \mathcal{G} .

A **Gibbs state with a nearest neighbor interaction V** is a Markov random field μ such that for all $x \in \text{supp}(\mu)$ and $A, B \subset \mathcal{V}_{\mathcal{G}}$ finite satisfying $\partial A \subset B \subset A^c$

$$\mu([x]_A \mid [x]_B) = \frac{\prod_{C \subset A \cup \partial A} e^{V([x]_C)}}{Z_{A,x|\partial A}}$$

where $Z_{A,x|\partial A}$ is the uniquely determined normalising factor dependent upon A and $x|_{\partial A}$.

Gibbs Measures

Given $X \subset \mathcal{A}^{\mathcal{V}_{\mathcal{G}}}$ we define a **nearest neighbour interaction** on X as a function $V : \{[a]_A \mid A \subset \mathcal{V}_{\mathcal{G}} \text{ finite}\} \rightarrow \mathbb{R}$ which is supported on patterns on the edges and the vertices of \mathcal{G} .

A **Gibbs state with a nearest neighbor interaction V** is a Markov random field μ such that for all $x \in \text{supp}(\mu)$ and $A, B \subset \mathcal{V}_{\mathcal{G}}$ finite satisfying $\partial A \subset B \subset A^c$

$$\mu([x]_A \mid [x]_B) = \frac{\prod_{C \subset A \cup \partial A} e^{V([x]_C)}}{Z_{A,x|\partial A}}$$

where $Z_{A,x|\partial A}$ is the uniquely determined normalising factor dependent upon A and $x|_{\partial A}$.

If $\mathcal{G} = \mathbb{Z}^d$ the specification of a shift-invariant Gibbs measure with a nearest neighbour interaction has a finite description:

Gibbs Measures

Given $X \subset \mathfrak{A}^{\mathcal{V}_{\mathcal{G}}}$ we define a **nearest neighbour interaction** on X as a function $V : \{[a]_A \mid A \subset \mathcal{V}_{\mathcal{G}} \text{ finite}\} \longrightarrow \mathbb{R}$ which is supported on patterns on the edges and the vertices of \mathcal{G} .

A **Gibbs state with a nearest neighbor interaction V** is a Markov random field μ such that for all $x \in \text{supp}(\mu)$ and $A, B \subset \mathcal{V}_{\mathcal{G}}$ finite satisfying $\partial A \subset B \subset A^c$

$$\mu([x]_A \mid [x]_B) = \frac{\prod_{C \subset A \cup \partial A} e^{V([x]_C)}}{Z_{A,x|\partial A}}$$

where $Z_{A,x|\partial A}$ is the uniquely determined normalising factor dependent upon A and $x|_{\partial A}$.

If $\mathcal{G} = \mathbb{Z}^d$ the specification of a shift-invariant Gibbs measure with a nearest neighbour interaction has a finite description: all we need is

Gibbs Measures

Given $X \in \mathcal{A}^{\mathcal{V}_{\mathcal{G}}}$ we define a **nearest neighbour interaction** on X as a function $V : \{[a]_A \mid A \subset \mathcal{V}_{\mathcal{G}} \text{ finite}\} \rightarrow \mathbb{R}$ which is supported on patterns on the edges and the vertices of \mathcal{G} .

A **Gibbs state with a nearest neighbor interaction V** is a Markov random field μ such that for all $x \in \text{supp}(\mu)$ and $A, B \subset \mathcal{V}_{\mathcal{G}}$ finite satisfying $\partial A \subset B \subset A^c$


$$\mu([x]_A \mid [x]_B) = \frac{\prod_{C \subset A \cup \partial A} e^{V([x]_C)}}{Z_{A,x|\partial A}}$$

where $Z_{A,x|\partial A}$ is the uniquely determined normalising factor dependent upon A and $x|_{\partial A}$.

If $\mathcal{G} = \mathbb{Z}^d$ the specification of a shift-invariant Gibbs measure with a nearest neighbour interaction has a finite description: all we need is the interaction V .


Gibbs Measures

Example: If the interaction on the hard square model is given by



Gibbs Measures


Example: If the interaction on the hard square model is given by



that is, $V([00]) = V([10]) = V([01]) = V([0]) = 0$ and $V([1]) = 1$ then

Gibbs Measures

Example: If the interaction on the hard square model is given by



that is, $V([00]) = V([10]) = V([01]) = V([0]) = 0$ and $V([1]) = 1$ then

$$\mu([x]_A \mid [x]_B) = \frac{e^{\sum_{C \subset A \cup \partial A} V([x]_C)}}{Z_{A, x|_{\partial A}}} = \frac{e^{\text{number of 1's in } x|_{A \cup \partial A}}}{Z_{A, x|_{\partial A}}}.$$

Question: Under what conditions on the support is a Markov random field Gibbs with some nearest neighbour interaction?

Previous Results

Positive results: (Instances where every Markov random field is Gibbs)

Previous Results

Positive results:(Instances where every Markov random field is Gibbs)

- The support has a safe symbol: Hammersley and Clifford('71)

Previous Results

Positive results: (Instances where every Markov random field is Gibbs)

- The support has a safe symbol: Hammersley and Clifford('71)
- Algebraic conditions on the support: Sturmfels, Gieger and Meek('06)

Previous Results

Positive results:(Instances where every Markov random field is Gibbs)

- The support has a safe symbol: Hammersley and Clifford('71)
- Algebraic conditions on the support: Sturmfels, Gieger and Meek('06)
- Decomposable graphs: Lauritzen('96)

Previous Results

Positive results:(Instances where every Markov random field is Gibbs)

- The support has a safe symbol: Hammersley and Clifford('71)
- Algebraic conditions on the support: Sturmfels, Gieger and Meek('06)
- Decomposable graphs: Lauritzen('96)
- For shift-invariant measures and $\mathcal{G} = \mathbb{Z}$ under some mixing conditions on the support: Georgii('88)

Previous Results

Positive results:(Instances where every Markov random field is Gibbs)

- The support has a safe symbol: Hammersley and Clifford('71)
- Algebraic conditions on the support: Sturmfels, Gieger and Meek('06)
- Decomposable graphs: Lauritzen('96)
- For shift-invariant measures and $\mathcal{G} = \mathbb{Z}$ under some mixing conditions on the support: Georgii('88)
- For shift-invariant measures and $\mathcal{G} = \mathbb{Z}$: Chandgotia, Han, Marcus, Meyerovitch and Pavlov('11)

Previous Results

Positive results:(Instances where every Markov random field is Gibbs)

- The support has a safe symbol: Hammersley and Clifford('71)
- Algebraic conditions on the support: Sturmfels, Gieger and Meek('06)
- Decomposable graphs: Lauritzen('96)
- For shift-invariant measures and $\mathcal{G} = \mathbb{Z}$ under some mixing conditions on the support: Georgii('88)
- For shift-invariant measures and $\mathcal{G} = \mathbb{Z}$: Chandgotia, Han, Marcus, Meyerovitch and Pavlov('11)
- For shift-invariant measures and support $\text{Hom}(\mathbb{Z}^d, \mathcal{H})$ where \mathcal{H} is an n cycle($n \neq 4$): Chandgotia and Meyerovitch('13)

Previous Results

Counterexamples: (Markov random fields which are not Gibbs)

Previous Results

Counterexamples: (Markov random fields which are not Gibbs)

- When \mathcal{G} is a finite graph: Moussouris('74)

Previous Results

Counterexamples: (Markov random fields which are not Gibbs)

- When \mathcal{G} is a finite graph: Moussouris('74)
- When $\mathcal{G} = \mathbb{Z}$ and the measure is not shift-invariant:
Dobruschin('68)

Previous Results

Counterexamples: (Markov random fields which are not Gibbs)

- When \mathcal{G} is a finite graph: Moussouris('74)
- When $\mathcal{G} = \mathbb{Z}$ and the measure is not shift-invariant: Dobruschin('68)
- When the alphabet is countable: Georgii('88)

Previous Results

Counterexamples: (Markov random fields which are not Gibbs)

- When \mathcal{G} is a finite graph: Moussouris('74)
- When $\mathcal{G} = \mathbb{Z}$ and the measure is not shift-invariant: Dobruschin('68)
- When the alphabet is countable: Georgii('88)
- A shift-invariant measure when $\mathcal{G} = \mathbb{Z}^2$: Chandgotia and Meyerovitch('13)

Previous Results

Counterexamples: (Markov random fields which are not Gibbs)

- When \mathcal{G} is a finite graph: Moussouris('74)
- When $\mathcal{G} = \mathbb{Z}$ and the measure is not shift-invariant: Dobruschin('68)
- When the alphabet is countable: Georgii('88)
- A shift-invariant measure when $\mathcal{G} = \mathbb{Z}^2$: Chandgotia and Meyerovitch('13)

Remark:

Previous Results

Counterexamples: (Markov random fields which are not Gibbs)

- When \mathcal{G} is a finite graph: Moussouris('74)
- When $\mathcal{G} = \mathbb{Z}$ and the measure is not shift-invariant: Dobruschin('68)
- When the alphabet is countable: Georgii('88)
- A shift-invariant measure when $\mathcal{G} = \mathbb{Z}^2$: Chandgotia and Meyerovitch('13)

Remark:

The measure we obtained was not Gibbs for any finite range interaction.

Previous Results

Counterexamples: (Markov random fields which are not Gibbs)

- When \mathcal{G} is a finite graph: Moussouris('74)
- When $\mathcal{G} = \mathbb{Z}$ and the measure is not shift-invariant: Dobruschin('68)
- When the alphabet is countable: Georgii('88)
- A shift-invariant measure when $\mathcal{G} = \mathbb{Z}^2$: Chandgotia and Meyerovitch('13)


Remark:

The measure we obtained was not Gibbs for any finite range interaction.

The support of the measure cannot be represented as $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$ for any graph \mathcal{H} .


Consequences of having a Safe Symbol

Suppose $\text{Hom}(\mathcal{G}, \mathcal{H})$ is the hard square model.



Consequences of having a Safe Symbol


Suppose $\text{Hom}(\mathcal{G}, \mathcal{H})$ is the hard square model.



Then for any pair $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$

Consequences of having a Safe Symbol


Suppose $\text{Hom}(\mathcal{G}, \mathcal{H})$ is the hard square model.



Then for any pair $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only at finitely many sites,

Consequences of having a Safe Symbol


Suppose $\text{Hom}(\mathcal{G}, \mathcal{H})$ is the hard square model.



Then for any pair $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only at finitely many sites, there is a chain $x^1 = x, x^2, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$

Consequences of having a Safe Symbol


Suppose $\text{Hom}(\mathcal{G}, \mathcal{H})$ is the hard square model.



Then for any pair $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only at finitely many sites, there is a chain $x^1 = x, x^2, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ such that x^i, x^{i+1} differ only at a single site.

Consequences of having a Safe Symbol

Suppose $\text{Hom}(\mathcal{G}, \mathcal{H})$ is the hard square model.



Then for any pair $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only at finitely many sites, there is a chain $x^1 = x, x^2, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ such that x^i, x^{i+1} differ only at a single site.

1	0	0	0	1
0	0	0	0	0
0	0	0	1	0
0	0	1	0	0
0	0	0	0	0


x

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	1
0	1	0	1	0

y

Consequences of having a Safe Symbol

Suppose $\text{Hom}(\mathcal{G}, \mathcal{H})$ is the hard square model.



Then for any pair $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only at finitely many sites, there is a chain $x^1 = x, x^2, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ such that x^i, x^{i+1} differ only at a single site.

1	0	0	0	1
0	0	0	0	0
0	0	0	1	0
0	0	1	0	0
0	0	0	0	0


x^1

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	1
0	1	0	1	0

y

Consequences of having a Safe Symbol

Suppose $\text{Hom}(\mathcal{G}, \mathcal{H})$ is the hard square model.



Then for any pair $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only at finitely many sites, there is a chain $x^1 = x, x^2, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ such that x^i, x^{i+1} differ only at a single site.

0	0	0	0	1
0	0	0	0	0
0	0	0	1	0
0	0	1	0	0
0	0	0	0	0


x^2

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	1
0	1	0	1	0

y

Consequences of having a Safe Symbol

Suppose $\text{Hom}(\mathcal{G}, \mathcal{H})$ is the hard square model.



Then for any pair $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only at finitely many sites, there is a chain $x^1 = x, x^2, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ such that x^i, x^{i+1} differ only at a single site.

0	0	0	0	0
0	0	0	0	0
0	0	0	1	0
0	0	1	0	0
0	0	0	0	0


x^3

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	1
0	1	0	1	0

y

Consequences of having a Safe Symbol

Suppose $\text{Hom}(\mathcal{G}, \mathcal{H})$ is the hard square model.



Then for any pair $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only at finitely many sites, there is a chain $x^1 = x, x^2, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ such that x^i, x^{i+1} differ only at a single site.

0	0	0	0	0
0	1	0	0	0
0	0	0	1	0
0	0	1	0	0
0	0	0	0	0


x^4

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	1
0	1	0	1	0

y

Consequences of having a Safe Symbol

Suppose $\text{Hom}(\mathcal{G}, \mathcal{H})$ is the hard square model.



Then for any pair $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only at finitely many sites, there is a chain $x^1 = x, x^2, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ such that x^i, x^{i+1} differ only at a single site.

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	1	0	0
0	0	0	0	0


x^5

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	1
0	1	0	1	0

y

Consequences of having a Safe Symbol

Suppose $\text{Hom}(\mathcal{G}, \mathcal{H})$ is the hard square model.



Then for any pair $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only at finitely many sites, there is a chain $x^1 = x, x^2, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ such that x^i, x^{i+1} differ only at a single site.

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0


x^6

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	1
0	1	0	1	0

y

Consequences of having a Safe Symbol

Suppose $\text{Hom}(\mathcal{G}, \mathcal{H})$ is the hard square model.



Then for any pair $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only at finitely many sites, there is a chain $x^1 = x, x^2, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ such that x^i, x^{i+1} differ only at a single site.

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	1
0	0	0	0	0


x^7

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	1
0	1	0	1	0

y

Consequences of having a Safe Symbol

Suppose $\text{Hom}(\mathcal{G}, \mathcal{H})$ is the hard square model.



Then for any pair $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only at finitely many sites, there is a chain $x^1 = x, x^2, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ such that x^i, x^{i+1} differ only at a single site.

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	1
0	1	0	0	0


x^8

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	1
0	1	0	1	0

y

Consequences of having a Safe Symbol

Suppose $\text{Hom}(\mathcal{G}, \mathcal{H})$ is the hard square model.



Then for any pair $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only at finitely many sites, there is a chain $x^1 = x, x^2, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ such that x^i, x^{i+1} differ only at a single site.

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	1
0	1	0	1	0


x^9

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	1
0	1	0	1	0

y

Consequences of having a Safe Symbol

Suppose $\text{Hom}(\mathcal{G}, \mathcal{H})$ is the hard square model.



Then for any pair $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only at finitely many sites, there is a chain $x^1 = x, x^2, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ such that x^i, x^{i+1} differ only at a single site.

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	1
0	1	0	1	0

y

0	0	0	0	0
0	1	0	0	0
0	0	0	0	0
0	0	0	0	1
0	1	0	1	0

y

The Pivot Property

A space $\text{Hom}(\mathcal{G}, \mathcal{H})$ is said to satisfy the **pivot property**

The Pivot Property

A space $\text{Hom}(\mathcal{G}, \mathcal{H})$ is said to satisfy the **pivot property** if for all $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only on finitely many sites

The Pivot Property

A space $\text{Hom}(\mathcal{G}, \mathcal{H})$ is said to satisfy the **pivot property** if for all $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only on finitely many sites there exists a chain of homomorphisms

$$x = x^1, x^2, x^3, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$$

The Pivot Property

A space $\text{Hom}(\mathcal{G}, \mathcal{H})$ is said to satisfy the **pivot property** if for all $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only on finitely many sites there exists a chain of homomorphisms

$$x = x^1, x^2, x^3, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$$

such that x^i, x^{i+1} differ on at most a single site.

The Pivot Property

A space $\text{Hom}(\mathcal{G}, \mathcal{H})$ is said to satisfy the **pivot property** if for all $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only on finitely many sites there exists a chain of homomorphisms

$$x = x^1, x^2, x^3, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$$

such that x^i, x^{i+1} differ on at most a single site.

Examples:

- If $\text{Hom}(\mathcal{G}, \mathcal{H})$ has a safe symbol.

The Pivot Property

A space $\text{Hom}(\mathcal{G}, \mathcal{H})$ is said to satisfy the **pivot property** if for all $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only on finitely many sites there exists a chain of homomorphisms

$$x = x^1, x^2, x^3, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$$

such that x^i, x^{i+1} differ on at most a single site.

Examples:

- If $\text{Hom}(\mathcal{G}, \mathcal{H})$ has a safe symbol.
- If $\mathcal{G} = \mathbb{Z}^d$ and \mathcal{H} is an n -cycle ($n=3$ corresponds to 3-colourings).

The Pivot Property

A space $\text{Hom}(\mathcal{G}, \mathcal{H})$ is said to satisfy the **pivot property** if for all $x, y \in \text{Hom}(\mathcal{G}, \mathcal{H})$ which differ only on finitely many sites there exists a chain of homomorphisms

$$x = x^1, x^2, x^3, \dots, x^n = y \in \text{Hom}(\mathcal{G}, \mathcal{H})$$

such that x^i, x^{i+1} differ on at most a single site.

Examples:

- If $\text{Hom}(\mathcal{G}, \mathcal{H})$ has a safe symbol.
- If $\mathcal{G} = \mathbb{Z}^d$ and \mathcal{H} is an n -cycle ($n=3$ corresponds to 3-colourings).
- If \mathcal{H} is dismantlable (to be defined in the next few slides).

The Pivot Property

Let $\mathcal{G} = \mathbb{Z}^2$ and suppose μ is a shift-invariant Markov random field whose support has the pivot property.

The Pivot Property

Let $\mathcal{G} = \mathbb{Z}^2$ and suppose μ is a shift-invariant Markov random field whose support has the pivot property. Then given $x, y \in \text{supp}(\mu)$ that differ exactly on a finite set F there exists a chain $x = x^1, x^2, \dots, x^n = y \in \text{supp}(\mu)$ where x^i, x^{i+1} differ exactly at a site $m_i \in \mathbb{Z}^2$

The Pivot Property

Let $\mathcal{G} = \mathbb{Z}^2$ and suppose μ is a shift-invariant Markov random field whose support has the pivot property. Then given $x, y \in \text{supp}(\mu)$ that differ exactly on a finite set F there exists a chain $x = x^1, x^2, \dots, x^n = y \in \text{supp}(\mu)$ where x^i, x^{i+1} differ exactly at a site $m_i \in \mathbb{Z}^2$ and consequently

$$\frac{\mu([x]_F \mid [x]_{\partial F})}{\mu([y]_F \mid [x]_{\partial F})}$$

The Pivot Property

Let $\mathcal{G} = \mathbb{Z}^2$ and suppose μ is a shift-invariant Markov random field whose support has the pivot property. Then given $x, y \in \text{supp}(\mu)$ that differ exactly on a finite set F there exists a chain $x = x^1, x^2, \dots, x^n = y \in \text{supp}(\mu)$ where x^i, x^{i+1} differ exactly at a site $m_i \in \mathbb{Z}^2$ and consequently

$$\frac{\mu([x]_F \mid [x]_{\partial F})}{\mu([y]_F \mid [x]_{\partial F})} = \prod_{i=1}^{n-1} \frac{\mu([x^i]_F \mid [x^i]_{\partial F})}{\mu([x^{i+1}]_F \mid [x^i]_{\partial F})}$$

The Pivot Property

Let $\mathcal{G} = \mathbb{Z}^2$ and suppose μ is a shift-invariant Markov random field whose support has the pivot property. Then given $x, y \in \text{supp}(\mu)$ that differ exactly on a finite set F there exists a chain $x = x^1, x^2, \dots, x^n = y \in \text{supp}(\mu)$ where x^i, x^{i+1} differ exactly at a site $m_i \in \mathbb{Z}^2$ and consequently

$$\begin{aligned} \frac{\mu([x]_F \mid [x]_{\partial F})}{\mu([y]_F \mid [x]_{\partial F})} &= \prod_{i=1}^{n-1} \frac{\mu([x^i]_F \mid [x^i]_{\partial F})}{\mu([x^{i+1}]_F \mid [x^i]_{\partial F})} \\ &= \prod_{i=1}^{n-1} \frac{\mu([x^i]_{m_i} \mid [x^i]_{\partial m_i})}{\mu([x^{i+1}]_{m_i} \mid [x^i]_{\partial m_i})}. \end{aligned}$$

The Pivot Property

Let $\mathcal{G} = \mathbb{Z}^2$ and suppose μ is a shift-invariant Markov random field whose support has the pivot property. Then given $x, y \in \text{supp}(\mu)$ that differ exactly on a finite set F there exists a chain $x = x^1, x^2, \dots, x^n = y \in \text{supp}(\mu)$ where x^i, x^{i+1} differ exactly at a site $m_i \in \mathbb{Z}^2$ and consequently

$$\begin{aligned} \frac{\mu([x]_F \mid [x]_{\partial F})}{\mu([y]_F \mid [x]_{\partial F})} &= \prod_{i=1}^{n-1} \frac{\mu([x^i]_F \mid [x^i]_{\partial F})}{\mu([x^{i+1}]_F \mid [x^i]_{\partial F})} \\ &= \prod_{i=1}^{n-1} \frac{\mu([x^i]_{m_i} \mid [x^i]_{\partial m_i})}{\mu([x^{i+1}]_{m_i} \mid [x^i]_{\partial m_i})}. \end{aligned}$$

Since μ is shift-invariant therefore the entire specification is determined by finitely many parameters viz.

The Pivot Property

Let $\mathcal{G} = \mathbb{Z}^2$ and suppose μ is a shift-invariant Markov random field whose support has the pivot property. Then given $x, y \in \text{supp}(\mu)$ that differ exactly on a finite set F there exists a chain $x = x^1, x^2, \dots, x^n = y \in \text{supp}(\mu)$ where x^i, x^{i+1} differ exactly at a site $m_i \in \mathbb{Z}^2$ and consequently

$$\begin{aligned} \frac{\mu([x]_F \mid [x]_{\partial F})}{\mu([y]_F \mid [x]_{\partial F})} &= \prod_{i=1}^{n-1} \frac{\mu([x^i]_F \mid [x^i]_{\partial F})}{\mu([x^{i+1}]_F \mid [x^i]_{\partial F})} \\ &= \prod_{i=1}^{n-1} \frac{\mu([x^i]_{m_i} \mid [x^i]_{\partial m_i})}{\mu([x^{i+1}]_{m_i} \mid [x^i]_{\partial m_i})}. \end{aligned}$$

Since μ is shift-invariant therefore the entire specification is determined by finitely many parameters viz. $\frac{\mu([x]_{0 \cup \partial 0})}{\mu([y]_{0 \cup \partial 0})}$ for configurations x, y which differ only at 0, the origin.

The Pivot Property

Then ratios of the form $\frac{\mu([x]_{0 \cup \partial 0})}{\mu([y]_{0 \cup \partial 0})}$

The Pivot Property

Then ratios of the form $\frac{\mu([x]_{0 \cup \partial 0})}{\mu([y]_{0 \cup \partial 0})}$ where x and y differ exactly on the origin

The Pivot Property

Then ratios of the form $\frac{\mu([x]_{0 \cup \partial 0})}{\mu([y]_{0 \cup \partial 0})}$ where x and y differ exactly on the origin determine whether μ is Gibbs or not.

3-colourings of \mathbb{Z}^2

Let \mathcal{H} be a 3-cycle with vertices 0, 1 and 2.

3-colourings of \mathbb{Z}^2

Let \mathcal{H} be a 3-cycle with vertices 0, 1 and 2. Then $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$ is the space of 3-colourings of \mathbb{Z}^2 where the colours are given by 0, 1 and 2.

3-colourings of \mathbb{Z}^2

Let \mathcal{H} be a 3-cycle with vertices 0, 1 and 2. Then $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$ is the space of 3-colourings of \mathbb{Z}^2 where the colours are given by 0, 1 and 2. If pairs $[x]_{0 \cup \partial 0}, [y]_{0 \cup \partial 0}$ differ exactly at the origin

3-colourings of \mathbb{Z}^2

Let \mathcal{H} be a 3-cycle with vertices 0, 1 and 2. Then $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$ is the space of 3-colourings of \mathbb{Z}^2 where the colours are given by 0, 1 and 2. If pairs $[x]_{0 \cup \partial 0}, [y]_{0 \cup \partial 0}$ differ exactly at the origin then $x|_{\partial 0}$ and $y|_{\partial 0}$

3-colourings of \mathbb{Z}^2

Let \mathcal{H} be a 3-cycle with vertices 0, 1 and 2. Then $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$ is the space of 3-colourings of \mathbb{Z}^2 where the colours are given by 0, 1 and 2. If pairs $[x]_{0 \cup \partial 0}, [y]_{0 \cup \partial 0}$ differ exactly at the origin then $x|_{\partial 0}$ and $y|_{\partial 0}$ are monochromatic.

3-colourings of \mathbb{Z}^2

Let \mathcal{H} be a 3-cycle with vertices 0, 1 and 2. Then $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$ is the space of 3-colourings of \mathbb{Z}^2 where the colours are given by 0, 1 and 2. If pairs $[x]_{0 \cup \partial 0}, [y]_{0 \cup \partial 0}$ differ exactly at the origin then $x|_{\partial 0}$ and $y|_{\partial 0}$ are monochromatic.

Thus a specification supported on the 3-coloured chessboard is

$$\text{determined the quantities } v_1 = \frac{\mu\left(\begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix}\right)}{\mu\left(\begin{bmatrix} 1 & 2 & 1 \\ 1 & 1 & 1 \end{bmatrix}\right)},$$

3-colourings of \mathbb{Z}^2

Let \mathcal{H} be a 3-cycle with vertices 0, 1 and 2. Then $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$ is the space of 3-colourings of \mathbb{Z}^2 where the colours are given by 0, 1 and 2. If pairs $[x]_{0 \cup \partial 0}, [y]_{0 \cup \partial 0}$ differ exactly at the origin then $x|_{\partial 0}$ and $y|_{\partial 0}$ are monochromatic.

Thus a specification supported on the 3-coloured chessboard is

$$\text{determined the quantities } v_1 = \frac{\mu\left(\begin{bmatrix} 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 \end{bmatrix}\right)}{\mu\left(\begin{bmatrix} 1 & 2 & 1 \\ 1 & 1 \end{bmatrix}\right)}, \quad v_2 = \frac{\mu\left(\begin{bmatrix} 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 \end{bmatrix}\right)}{\mu\left(\begin{bmatrix} 2 & 0 & 2 \\ 2 & 2 \end{bmatrix}\right)}$$

3-colourings of \mathbb{Z}^2

Let \mathcal{H} be a 3-cycle with vertices 0, 1 and 2. Then $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$ is the space of 3-colourings of \mathbb{Z}^2 where the colours are given by 0, 1 and 2. If pairs $[x]_{0 \cup \partial 0}, [y]_{0 \cup \partial 0}$ differ exactly at the origin then $x|_{\partial 0}$ and $y|_{\partial 0}$ are monochromatic.

Thus a specification supported on the 3-coloured chessboard is

determined the quantities $v_1 = \frac{\mu\left(\begin{bmatrix} 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 \end{bmatrix}\right)}{\mu\left(\begin{bmatrix} 1 & 2 \\ 1 & 2 & 1 \\ 1 & 1 \end{bmatrix}\right)}, v_2 = \frac{\mu\left(\begin{bmatrix} 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 \end{bmatrix}\right)}{\mu\left(\begin{bmatrix} 2 & 0 \\ 2 & 0 & 2 \\ 2 & 2 \end{bmatrix}\right)}$ and

$$v_3 = \frac{\mu\left(\begin{bmatrix} 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 \end{bmatrix}\right)}{\mu\left(\begin{bmatrix} 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 \end{bmatrix}\right)}.$$

3-colourings of \mathbb{Z}^2

Let \mathcal{H} be a 3-cycle with vertices 0, 1 and 2. Then $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$ is the space of 3-colourings of \mathbb{Z}^2 where the colours are given by 0, 1 and 2. If pairs $[x]_{0 \cup \partial 0}, [y]_{0 \cup \partial 0}$ differ exactly at the origin then $x|_{\partial 0}$ and $y|_{\partial 0}$ are monochromatic.

Thus a specification supported on the 3-coloured chessboard is

determined the quantities $v_1 = \frac{\mu\left(\begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix}\right)}{\mu\left(\begin{bmatrix} 1 & 2 & 1 \\ 1 & 1 & 1 \end{bmatrix}\right)}, v_2 = \frac{\mu\left(\begin{bmatrix} 2 & 2 & 2 \\ 2 & 2 & 2 \end{bmatrix}\right)}{\mu\left(\begin{bmatrix} 2 & 0 & 2 \\ 2 & 2 & 2 \end{bmatrix}\right)}$ and

$v_3 = \frac{\mu\left(\begin{bmatrix} 0 & 2 & 0 \\ 0 & 0 & 0 \end{bmatrix}\right)}{\mu\left(\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}\right)}$. If μ is a Gibbs measure with some nearest neighbour interaction V then

$$\begin{aligned}
v_1 &= \exp(V(01) + V(10) + V(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix}) + V(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix}) + V(0) \\
&\quad - V(21) - V(12) - V(\begin{smallmatrix} 2 \\ 1 \end{smallmatrix}) - V(\begin{smallmatrix} 1 \\ 2 \end{smallmatrix}) - V(2)), \\
v_2 &= \exp(V(12) + V(21) + V(\begin{smallmatrix} 2 \\ 1 \end{smallmatrix}) + V(\begin{smallmatrix} 1 \\ 2 \end{smallmatrix}) + V(1) \\
&\quad - V(02) - V(20) - V(\begin{smallmatrix} 0 \\ 2 \end{smallmatrix}) - V(\begin{smallmatrix} 2 \\ 0 \end{smallmatrix}) - V(0)), \\
v_3 &= \exp(V(02) + V(20) + V(\begin{smallmatrix} 2 \\ 0 \end{smallmatrix}) + V(\begin{smallmatrix} 0 \\ 2 \end{smallmatrix}) + V(2) \\
&\quad - V(01) - V(10) - V(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix}) - V(\begin{smallmatrix} 1 \\ 0 \end{smallmatrix}) - V(1)).
\end{aligned}$$

$$\begin{aligned}
v_1 &= \exp(V(01) + V(10) + V(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix}) + V(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix}) + V(0) \\
&\quad - V(21) - V(12) - V(\begin{smallmatrix} 2 \\ 1 \end{smallmatrix}) - V(\begin{smallmatrix} 1 \\ 2 \end{smallmatrix}) - V(2)), \\
v_2 &= \exp(V(12) + V(21) + V(\begin{smallmatrix} 2 \\ 1 \end{smallmatrix}) + V(\begin{smallmatrix} 1 \\ 2 \end{smallmatrix}) + V(1) \\
&\quad - V(02) - V(20) - V(\begin{smallmatrix} 0 \\ 2 \end{smallmatrix}) - V(\begin{smallmatrix} 2 \\ 0 \end{smallmatrix}) - V(0)), \\
v_3 &= \exp(V(02) + V(20) + V(\begin{smallmatrix} 2 \\ 0 \end{smallmatrix}) + V(\begin{smallmatrix} 0 \\ 2 \end{smallmatrix}) + V(2) \\
&\quad - V(01) - V(10) - V(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix}) - V(\begin{smallmatrix} 1 \\ 0 \end{smallmatrix}) - V(1)).
\end{aligned}$$

μ is Gibbs if and only if $v_1 v_2 v_3 = 1$.

$$\begin{aligned}
v_1 &= \exp(V(01) + V(10) + V(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix}) + V(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix}) + V(0) \\
&\quad - V(21) - V(12) - V(\begin{smallmatrix} 2 \\ 1 \end{smallmatrix}) - V(\begin{smallmatrix} 1 \\ 2 \end{smallmatrix}) - V(2)), \\
v_2 &= \exp(V(12) + V(21) + V(\begin{smallmatrix} 2 \\ 1 \end{smallmatrix}) + V(\begin{smallmatrix} 1 \\ 2 \end{smallmatrix}) + V(1) \\
&\quad - V(02) - V(20) - V(\begin{smallmatrix} 0 \\ 2 \end{smallmatrix}) - V(\begin{smallmatrix} 2 \\ 0 \end{smallmatrix}) - V(0)), \\
v_3 &= \exp(V(02) + V(20) + V(\begin{smallmatrix} 2 \\ 0 \end{smallmatrix}) + V(\begin{smallmatrix} 0 \\ 2 \end{smallmatrix}) + V(2) \\
&\quad - V(01) - V(10) - V(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix}) - V(\begin{smallmatrix} 1 \\ 0 \end{smallmatrix}) - V(1)).
\end{aligned}$$

μ is Gibbs if and only if $v_1 v_2 v_3 = 1$. Thus we need more than just the pivot property to prove that it is Gibbs.

$$\begin{aligned}
v_1 &= \exp(V(01) + V(10) + V(\overset{0}{1}) + V(\overset{0}{1}) + V(0) \\
&\quad - V(21) - V(12) - V(\overset{2}{1}) - V(\overset{1}{2}) - V(2)), \\
v_2 &= \exp(V(12) + V(21) + V(\overset{2}{1}) + V(\overset{1}{2}) + V(1) \\
&\quad - V(02) - V(20) - V(\overset{0}{2}) - V(\overset{2}{0}) - V(0)), \\
v_3 &= \exp(V(02) + V(20) + V(\overset{2}{0}) + V(\overset{0}{2}) + V(2) \\
&\quad - V(01) - V(10) - V(\overset{0}{1}) - V(\overset{1}{0}) - V(1)).
\end{aligned}$$

μ is Gibbs if and only if $v_1 v_2 v_3 = 1$. Thus we need more than just the pivot property to prove that it is Gibbs. Note that this is only on the level of the **specifications**,

$$\begin{aligned}
v_1 &= \exp(V(01) + V(10) + V(\overset{0}{1}) + V(\overset{0}{1}) + V(0) \\
&\quad - V(21) - V(12) - V(\overset{2}{1}) - V(\overset{1}{2}) - V(2)), \\
v_2 &= \exp(V(12) + V(21) + V(\overset{2}{1}) + V(\overset{1}{2}) + V(1) \\
&\quad - V(02) - V(20) - V(\overset{0}{2}) - V(\overset{2}{0}) - V(0)), \\
v_3 &= \exp(V(02) + V(20) + V(\overset{2}{0}) + V(\overset{0}{2}) + V(2) \\
&\quad - V(01) - V(10) - V(\overset{0}{1}) - V(\overset{1}{0}) - V(1)).
\end{aligned}$$

μ is Gibbs if and only if $v_1 v_2 v_3 = 1$. Thus we need more than just the pivot property to prove that it is Gibbs. Note that this is only on the level of the **specifications**, not the measures themselves

$$\begin{aligned}
v_1 &= \exp(V(01) + V(10) + V(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix}) + V(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix}) + V(0) \\
&\quad - V(21) - V(12) - V(\begin{smallmatrix} 2 \\ 1 \end{smallmatrix}) - V(\begin{smallmatrix} 1 \\ 2 \end{smallmatrix}) - V(2)), \\
v_2 &= \exp(V(12) + V(21) + V(\begin{smallmatrix} 2 \\ 1 \end{smallmatrix}) + V(\begin{smallmatrix} 1 \\ 2 \end{smallmatrix}) + V(1) \\
&\quad - V(02) - V(20) - V(\begin{smallmatrix} 0 \\ 2 \end{smallmatrix}) - V(\begin{smallmatrix} 2 \\ 0 \end{smallmatrix}) - V(0)), \\
v_3 &= \exp(V(02) + V(20) + V(\begin{smallmatrix} 2 \\ 0 \end{smallmatrix}) + V(\begin{smallmatrix} 0 \\ 2 \end{smallmatrix}) + V(2) \\
&\quad - V(01) - V(10) - V(\begin{smallmatrix} 0 \\ 1 \end{smallmatrix}) - V(\begin{smallmatrix} 1 \\ 0 \end{smallmatrix}) - V(1)).
\end{aligned}$$

μ is Gibbs if and only if $v_1 v_2 v_3 = 1$. Thus we need more than just the pivot property to prove that it is Gibbs. Note that this is only on the level of the **specifications**, not the measures themselves and under the assumption of shift-invariance.

Graph Folding(Nowakowski and Winkler-'83)

Consider an undirected finite graph \mathcal{H} .

Graph Folding(Nowakowski and Winkler-'83)

Consider an undirected finite graph \mathcal{H} . Let $N(w)$ denote the *neighbourhood* of w in \mathcal{H} ,

Graph Folding(Nowakowski and Winkler-'83)

Consider an undirected finite graph \mathcal{H} . Let $N(w)$ denote the *neighbourhood* of w in \mathcal{H} , that is,

$$N(w) = \{v \in \mathcal{H} \mid v \sim w\}.$$

Graph Folding(Nowakowski and Winkler-'83)

Consider an undirected finite graph \mathcal{H} . Let $N(w)$ denote the *neighbourhood* of w in \mathcal{H} , that is,

$$N(w) = \{v \in \mathcal{H} \mid v \sim w\}.$$

We will say that \mathcal{H} can be **folded** to a graph $\mathcal{H} \setminus \{v\}$ if there exists a vertex $w \in \mathcal{H}$ such that $N(v) \subset N(w)$.

Graph Folding(Nowakowski and Winkler-'83)

Consider an undirected finite graph \mathcal{H} . Let $N(w)$ denote the *neighbourhood* of w in \mathcal{H} , that is,

$$N(w) = \{v \in \mathcal{H} \mid v \sim w\}.$$

We will say that \mathcal{H} can be **folded** to a graph $\mathcal{H} \setminus \{v\}$ if there exists a vertex $w \in \mathcal{H}$ such that $N(v) \subset N(w)$. \mathcal{H} is called **dismantleable** if \mathcal{H} can be folded all the way to a single vertex(with or without a loop).

Graph Folding(Nowakowski and Winkler-'83)

Consider an undirected finite graph \mathcal{H} . Let $N(w)$ denote the *neighbourhood* of w in \mathcal{H} , that is,

$$N(w) = \{v \in \mathcal{H} \mid v \sim w\}.$$

We will say that \mathcal{H} can be **folded** to a graph $\mathcal{H} \setminus \{v\}$ if there exists a vertex $w \in \mathcal{H}$ such that $N(v) \subset N(w)$. \mathcal{H} is called **dismantleable** if \mathcal{H} can be folded all the way to a single vertex(with or without a loop).

Examples:

Graph Folding(Nowakowski and Winkler-'83)

Consider an undirected finite graph \mathcal{H} . Let $N(w)$ denote the *neighbourhood* of w in \mathcal{H} , that is,

$$N(w) = \{v \in \mathcal{H} \mid v \sim w\}.$$

We will say that \mathcal{H} can be **folded** to a graph $\mathcal{H} \setminus \{v\}$ if there exists a vertex $w \in \mathcal{H}$ such that $N(v) \subset N(w)$. \mathcal{H} is called **dismantleable** if \mathcal{H} can be folded all the way to a single vertex(with or without a loop).

Examples:

Any graph \mathcal{H} such that $\text{Hom}(\mathcal{G}, \mathcal{H})$ has a safe symbol \star .

Graph Folding(Nowakowski and Winkler-'83)

Consider an undirected finite graph \mathcal{H} . Let $N(w)$ denote the *neighbourhood* of w in \mathcal{H} , that is,

$$N(w) = \{v \in \mathcal{H} \mid v \sim w\}.$$

We will say that \mathcal{H} can be **folded** to a graph $\mathcal{H} \setminus \{v\}$ if there exists a vertex $w \in \mathcal{H}$ such that $N(v) \subset N(w)$. \mathcal{H} is called **dismantleable** if \mathcal{H} can be folded all the way to a single vertex(with or without a loop).

Examples:

Any graph \mathcal{H} such that $\text{Hom}(\mathcal{G}, \mathcal{H})$ has a safe symbol \star .

For any vertex $v \in \mathcal{V}_{\mathcal{H}}$, $N(v) \subset N(\star) = \mathcal{V}_{\mathcal{H}}$

Graph Folding(Nowakowski and Winkler-'83)

Consider an undirected finite graph \mathcal{H} . Let $N(w)$ denote the *neighbourhood* of w in \mathcal{H} , that is,

$$N(w) = \{v \in \mathcal{H} \mid v \sim w\}.$$

We will say that \mathcal{H} can be **folded** to a graph $\mathcal{H} \setminus \{v\}$ if there exists a vertex $w \in \mathcal{H}$ such that $N(v) \subset N(w)$. \mathcal{H} is called **dismantleable** if \mathcal{H} can be folded all the way to a single vertex(with or without a loop).

Examples:

Any graph \mathcal{H} such that $\text{Hom}(\mathcal{G}, \mathcal{H})$ has a safe symbol \star .


For any vertex $v \in \mathcal{V}_{\mathcal{H}}$, $N(v) \subset N(\star) = \mathcal{V}_{\mathcal{H}}$ and thus any vertex v can be folded into \star .

Dismantleable Graphs

The following graph \mathcal{H} is dismantlable:


Dismantleable Graphs

The following graph \mathcal{H} is dismantlable:



Dismantleable Graphs


The following graph \mathcal{H} is dismantlable:



by the folding sequence: 3 folds into 1,

Dismantleable Graphs


The following graph \mathcal{H} is dismantlable:



by the folding sequence: 3 folds into 1, 2 folds into 1

Dismantleable Graphs


The following graph \mathcal{H} is dismantlable:



by the folding sequence: 3 folds into 1, 2 folds into 1 and finally 4 folds into 1.

Dismantleable Graphs


The following graph \mathcal{H} is dismantlable:




by the folding sequence: 3 folds into 1, 2 folds into 1 and finally 4 folds into 1. Note, $\text{Hom}(\mathcal{G}, \mathcal{H})$ does not have a safe symbol.

Dismantleable Graphs

The following graph \mathcal{H} is dismantlable:




by the folding sequence: 3 folds into 1, 2 folds into 1 and finally 4 folds into 1. Note, $\text{Hom}(\mathcal{G}, \mathcal{H})$ does not have a safe symbol. The following graph is not dismantlable:




Dismantleable Graphs

The following graph \mathcal{H} is dismantlable:



by the folding sequence: 3 folds into 1, 2 folds into 1 and finally 4 folds into 1. Note, $\text{Hom}(\mathcal{G}, \mathcal{H})$ does not have a safe symbol. The following graph is not dismantlable:



No vertex can be folded into the other. Such a graph is said to be **stiff**.

Hammersley-Clifford Spaces

A space $\text{Hom}(\mathcal{G}, \mathcal{H})$ is called a **Hammersley-Clifford space**

Hammersley-Clifford Spaces

A space $Hom(\mathcal{G}, \mathcal{H})$ is called a **Hammersley-Clifford space** if all Markov specifications on $Hom(\mathcal{G}, \mathcal{H})$ are Gibbs for some nearest neighbour interaction.

Hammersley-Clifford Spaces

A space $Hom(\mathcal{G}, \mathcal{H})$ is called a **Hammersley-Clifford space** if all Markov specifications on $Hom(\mathcal{G}, \mathcal{H})$ are Gibbs for some nearest neighbour interaction.

Remark :

Hammersley-Clifford Spaces

A space $Hom(\mathcal{G}, \mathcal{H})$ is called a **Hammersley-Clifford space** if all Markov specifications on $Hom(\mathcal{G}, \mathcal{H})$ are Gibbs for some nearest neighbour interaction.

Remark : If $Hom(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space

Hammersley-Clifford Spaces

A space $Hom(\mathcal{G}, \mathcal{H})$ is called a **Hammersley-Clifford space** if all Markov specifications on $Hom(\mathcal{G}, \mathcal{H})$ are Gibbs for some nearest neighbour interaction.

Remark : If $Hom(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space then for all Markov random fields μ such that $supp(\mu) = Hom(\mathcal{G}, \mathcal{H})$,

Hammersley-Clifford Spaces

A space $Hom(\mathcal{G}, \mathcal{H})$ is called a **Hammersley-Clifford space** if all Markov specifications on $Hom(\mathcal{G}, \mathcal{H})$ are Gibbs for some nearest neighbour interaction.

Remark : If $Hom(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space then for all Markov random fields μ such that $supp(\mu) = Hom(\mathcal{G}, \mathcal{H})$, μ is a Gibbs measure with some nearest neighbour interaction.

Hammersley-Clifford Spaces

A space $Hom(\mathcal{G}, \mathcal{H})$ is called a **Hammersley-Clifford space** if all Markov specifications on $Hom(\mathcal{G}, \mathcal{H})$ are Gibbs for some nearest neighbour interaction.

Remark : If $Hom(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space then for all Markov random fields μ such that $supp(\mu) = Hom(\mathcal{G}, \mathcal{H})$, μ is a Gibbs measure with some nearest neighbour interaction. The converse need not be true.

Hammersley-Clifford Spaces

A space $Hom(\mathcal{G}, \mathcal{H})$ is called a **Hammersley-Clifford space** if all Markov specifications on $Hom(\mathcal{G}, \mathcal{H})$ are Gibbs for some nearest neighbour interaction.

Remark : If $Hom(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space then for all Markov random fields μ such that $supp(\mu) = Hom(\mathcal{G}, \mathcal{H})$, μ is a Gibbs measure with some nearest neighbour interaction. The converse need not be true.

Examples:

Hammersley-Clifford Spaces

A space $Hom(\mathcal{G}, \mathcal{H})$ is called a **Hammersley-Clifford space** if all Markov specifications on $Hom(\mathcal{G}, \mathcal{H})$ are Gibbs for some nearest neighbour interaction.

Remark : If $Hom(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space then for all Markov random fields μ such that $supp(\mu) = Hom(\mathcal{G}, \mathcal{H})$, μ is a Gibbs measure with some nearest neighbour interaction. The converse need not be true.

Examples:

- ① $Hom(\mathcal{G}, \mathcal{H})$ such that it has a safe symbol,

Hammersley-Clifford Spaces

A space $Hom(\mathcal{G}, \mathcal{H})$ is called a **Hammersley-Clifford space** if all Markov specifications on $Hom(\mathcal{G}, \mathcal{H})$ are Gibbs for some nearest neighbour interaction.

Remark : If $Hom(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space then for all Markov random fields μ such that $supp(\mu) = Hom(\mathcal{G}, \mathcal{H})$, μ is a Gibbs measure with some nearest neighbour interaction. The converse need not be true.

Examples:

- ① $Hom(\mathcal{G}, \mathcal{H})$ such that it has a safe symbol,
- ② $Hom(\mathcal{G}, \mathcal{H})$ for \mathcal{H} being a single vertex or an edge.

Hammersley-Clifford Spaces

A space $Hom(\mathcal{G}, \mathcal{H})$ is called a **Hammersley-Clifford space** if all Markov specifications on $Hom(\mathcal{G}, \mathcal{H})$ are Gibbs for some nearest neighbour interaction.

Remark : If $Hom(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space then for all Markov random fields μ such that $supp(\mu) = Hom(\mathcal{G}, \mathcal{H})$, μ is a Gibbs measure with some nearest neighbour interaction. The converse need not be true.

Examples:

- ① $Hom(\mathcal{G}, \mathcal{H})$ such that it has a safe symbol,
- ② $Hom(\mathcal{G}, \mathcal{H})$ for \mathcal{H} being a single vertex or an edge.
- ③ $Hom(\mathbb{Z}^d, \mathcal{H})$ where \mathcal{H} is a n-cycle with $n \neq 4$.

Graph Folding and Hammersley-Clifford Spaces

Theorem (Chandgotia-'14)

Let \mathcal{G} be bipartite graph

Graph Folding and Hammersley-Clifford Spaces

Theorem (Chandgotia-'14)

Let \mathcal{G} be bipartite graph and \mathcal{H} be a graph with a fold $\mathcal{H} \setminus \{v\}$.

Graph Folding and Hammersley-Clifford Spaces

Theorem (Chandgotia-'14)

Let \mathcal{G} be bipartite graph and \mathcal{H} be a graph with a fold $\mathcal{H} \setminus \{v\}$. Then the space $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space if and only if $\text{Hom}(\mathcal{G}, \mathcal{H} \setminus \{v\})$ is a Hammersley-Clifford space as well.

Graph Folding and Hammersley-Clifford Spaces

Theorem (Chandgotia-'14)

Let \mathcal{G} be bipartite graph and \mathcal{H} be a graph with a fold $\mathcal{H} \setminus \{v\}$. Then the space $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space if and only if $\text{Hom}(\mathcal{G}, \mathcal{H} \setminus \{v\})$ is a Hammersley-Clifford space as well.

Remark:

Graph Folding and Hammersley-Clifford Spaces

Theorem (Chandgotia-'14)

Let \mathcal{G} be bipartite graph and \mathcal{H} be a graph with a fold $\mathcal{H} \setminus \{v\}$. Then the space $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space if and only if $\text{Hom}(\mathcal{G}, \mathcal{H} \setminus \{v\})$ is a Hammersley-Clifford space as well.

Remark:

- There is a corresponding version of the theorem where the specifications and interactions are assumed to be invariant under some automorphism of the graph \mathcal{G} e.g. translations in the case of \mathbb{Z}^d .

Graph Folding and Hammersley-Clifford Spaces

Theorem (Chandgotia-'14)

Let \mathcal{G} be bipartite graph and \mathcal{H} be a graph with a fold $\mathcal{H} \setminus \{v\}$. Then the space $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space if and only if $\text{Hom}(\mathcal{G}, \mathcal{H} \setminus \{v\})$ is a Hammersley-Clifford space as well.

Remark:

- There is a corresponding version of the theorem where the specifications and interactions are assumed to be invariant under some automorphism of the graph \mathcal{G} e.g. translations in the case of \mathbb{Z}^d .
- This result is true for a more general notion of folding on closed spaces of configurations, not just restricted to homomorphism spaces.

An idea of the proof:

For simplicity we will assume that

An idea of the proof:

For simplicity we will assume that $\mathcal{G} = \mathbb{Z}^2$ and $v \approx v$.

An idea of the proof:

For simplicity we will assume that $\mathcal{G} = \mathbb{Z}^2$ and $v \approx v$. Suppose $\mathcal{H} \setminus \{v\}$ is a fold of a graph \mathcal{H} where vertex v is folded onto vertex w .

An idea of the proof:

For simplicity we will assume that $\mathcal{G} = \mathbb{Z}^2$ and $v \approx v$. Suppose $\mathcal{H} \setminus \{v\}$ is a fold of a graph \mathcal{H} where vertex v is folded onto vertex w . What does this imply about the spaces $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$?

An idea of the proof:

For simplicity we will assume that $\mathcal{G} = \mathbb{Z}^2$ and $v \approx v$. Suppose $\mathcal{H} \setminus \{v\}$ is a fold of a graph \mathcal{H} where vertex v is folded onto vertex w . What does this imply about the spaces $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$?

- Since $N(v) \subset N(w)$,

An idea of the proof:

For simplicity we will assume that $\mathcal{G} = \mathbb{Z}^2$ and $v \approx v$. Suppose $\mathcal{H} \setminus \{v\}$ is a fold of a graph \mathcal{H} where vertex v is folded onto vertex w . What does this imply about the spaces $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$?

- Since $N(v) \subset N(w)$, given an element $x \in \text{Hom}(\mathbb{Z}^2, \mathcal{H})$ such that $x_0 = v$,

An idea of the proof:

For simplicity we will assume that $\mathcal{G} = \mathbb{Z}^2$ and $v \approx v$. Suppose $\mathcal{H} \setminus \{v\}$ is a fold of a graph \mathcal{H} where vertex v is folded onto vertex w . What does this imply about the spaces $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$?

- Since $N(v) \subset N(w)$, given an element $x \in \text{Hom}(\mathbb{Z}^2, \mathcal{H})$ such that $x_0 = v$, we can replace the v by w at 0

An idea of the proof:


For simplicity we will assume that $\mathcal{G} = \mathbb{Z}^2$ and $v \approx v$. Suppose $\mathcal{H} \setminus \{v\}$ is a fold of a graph \mathcal{H} where vertex v is folded onto vertex w . What does this imply about the spaces $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$?

- Since $N(v) \subset N(w)$, given an element $x \in \text{Hom}(\mathbb{Z}^2, \mathcal{H})$ such that $x_0 = v$, we can replace the v by w at 0 and still be an element of $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$.

An idea of the proof:

For simplicity we will assume that $\mathcal{G} = \mathbb{Z}^2$ and $v \approx v$. Suppose $\mathcal{H} \setminus \{v\}$ is a fold of a graph \mathcal{H} where vertex v is folded onto vertex w . What does this imply about the spaces $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$?


- Since $N(v) \subset N(w)$, given an element $x \in \text{Hom}(\mathbb{Z}^2, \mathcal{H})$ such that $x_0 = v$, we can replace the v by w at 0 and still be an element of $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$.



An idea of the proof:

For simplicity we will assume that $\mathcal{G} = \mathbb{Z}^2$ and $v \approx v$. Suppose $\mathcal{H} \setminus \{v\}$ is a fold of a graph \mathcal{H} where vertex v is folded onto vertex w . What does this imply about the spaces $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$?

- Since $N(v) \subset N(w)$, given an element $x \in \text{Hom}(\mathbb{Z}^2, \mathcal{H})$ such that $x_0 = v$, we can replace the v by w at 0 and still be an element of $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$.




- Also we can surround $x|_{0 \cup \partial 0}$ by w 's

An idea of the proof:

For simplicity we will assume that $\mathcal{G} = \mathbb{Z}^2$ and $v \approx v$. Suppose $\mathcal{H} \setminus \{v\}$ is a fold of a graph \mathcal{H} where vertex v is folded onto vertex w . What does this imply about the spaces $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$?

- Since $N(v) \subset N(w)$, given an element $x \in \text{Hom}(\mathbb{Z}^2, \mathcal{H})$ such that $x_0 = v$, we can replace the v by w at 0 and still be an element of $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$.




- Also we can surround $x|_{0 \cup \partial 0}$ by w 's such that the partial configuration thus obtained is still valid in $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$.

An idea of the proof:

For simplicity we will assume that $\mathcal{G} = \mathbb{Z}^2$ and $v \approx v$. Suppose $\mathcal{H} \setminus \{v\}$ is a fold of a graph \mathcal{H} where vertex v is folded onto vertex w . What does this imply about the spaces $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$?

- Since $N(v) \subset N(w)$, given an element $x \in \text{Hom}(\mathbb{Z}^2, \mathcal{H})$ such that $x_0 = v$, we can replace the v by w at 0 and still be an element of $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$.



- Also we can surround $x|_{0 \cup \partial 0}$ by w 's such that the partial configuration thus obtained is still valid in $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$.

An idea of the proof:

Suppose $\text{Hom}(\mathbb{Z}^2, \mathcal{H} \setminus \{v\})$ is a Hammersley-Clifford space.

An idea of the proof:

Suppose $\text{Hom}(\mathbb{Z}^2, \mathcal{H} \setminus \{v\})$ is a Hammersley-Clifford space. Let μ be a Markov random field on $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$.

An idea of the proof:

Suppose $\text{Hom}(\mathbb{Z}^2, \mathcal{H} \setminus \{v\})$ is a Hammersley-Clifford space. Let μ be a Markov random field on $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$. To prove that it is Gibbs

An idea of the proof:

Suppose $\text{Hom}(\mathbb{Z}^2, \mathcal{H} \setminus \{v\})$ is a Hammersley-Clifford space. Let μ be a Markov random field on $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$. To prove that it is Gibbs we need to build a correspondence between patterns on edges and vertices of \mathbb{Z}^2

An idea of the proof:

Suppose $\text{Hom}(\mathbb{Z}^2, \mathcal{H} \setminus \{v\})$ is a Hammersley-Clifford space. Let μ be a Markov random field on $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$. To prove that it is Gibbs we need to build a correspondence between patterns on edges and vertices of \mathbb{Z}^2 and pairs of elements of $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$ which differ at a single site.

An idea of the proof:

Suppose $\text{Hom}(\mathbb{Z}^2, \mathcal{H} \setminus \{v\})$ is a Hammersley-Clifford space. Let μ be a Markov random field on $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$. To prove that it is Gibbs we need to build a correspondence between patterns on edges and vertices of \mathbb{Z}^2 and pairs of elements of $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$ which differ at a single site.

Since $\text{Hom}(\mathbb{Z}^2, \mathcal{H} \setminus \{v\})$ is a Hammersley-Clifford space

An idea of the proof:

Suppose $\text{Hom}(\mathbb{Z}^2, \mathcal{H} \setminus \{v\})$ is a Hammersley-Clifford space. Let μ be a Markov random field on $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$. To prove that it is Gibbs we need to build a correspondence between patterns on edges and vertices of \mathbb{Z}^2 and pairs of elements of $\text{Hom}(\mathbb{Z}^2, \mathcal{H})$ which differ at a single site.

Since $\text{Hom}(\mathbb{Z}^2, \mathcal{H} \setminus \{v\})$ is a Hammersley-Clifford space we only care about pairs which involve changing a single v to w .

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$.

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$V([xv]) \quad \text{with} \quad \begin{matrix} w \\ x \\ w \end{matrix} v, \begin{matrix} w \\ a \\ w \end{matrix} v$$

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with} \quad \begin{array}{c} w \\ w \end{array} x \begin{array}{c} w \\ w \end{array} v, \quad \begin{array}{c} w \\ w \end{array} a \begin{array}{c} w \\ w \end{array} v \\
 V([vx]) & \text{with} \quad \begin{array}{c} w \\ w \end{array} v \begin{array}{c} w \\ w \end{array} x \begin{array}{c} w \\ w \end{array} w, \quad \begin{array}{c} w \\ w \end{array} v \begin{array}{c} w \\ w \end{array} a \begin{array}{c} w \\ w \end{array} w
 \end{array}$$

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} v \\ w \end{array} \begin{array}{c} a \\ w \end{array} w
 \end{array}$$

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} v \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ v \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ v \end{array} \begin{array}{c} a \\ w \end{array} w
 \end{array}$$

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} v \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ v \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ v \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([v]) & \text{with } \begin{array}{c} a \\ a \end{array} \begin{array}{c} v \\ a \end{array} a, \begin{array}{c} a \\ a \end{array} \begin{array}{c} w \\ a \end{array} a
 \end{array}$$

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} v \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ v \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ v \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([v]) & \text{with } \begin{array}{c} a \\ a \end{array} \begin{array}{c} v \\ a \end{array} a, \begin{array}{c} a \\ a \end{array} \begin{array}{c} w \\ a \end{array} a
 \end{array}$$

We can use the following order to change a single v to w :

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} v \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ v \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ v \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([v]) & \text{with } \begin{array}{c} a \\ a \end{array} \begin{array}{c} v \\ a \end{array} a, \begin{array}{c} a \\ a \end{array} \begin{array}{c} w \\ a \end{array} a
 \end{array}$$

We can use the following order to change a single v to w :

		w		
	w	e	w	
w	b	v	c	w
	w	d	w	
		w		

		w		
	w	e	w	
w	b	v	c	w
	w	d	w	
		w		

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} v \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ v \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ v \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([v]) & \text{with } \begin{array}{c} a \\ a \end{array} \begin{array}{c} v \\ a \end{array} a, \begin{array}{c} a \\ a \end{array} \begin{array}{c} w \\ a \end{array} a
 \end{array}$$

We can use the following order to change a single v to w :

		w		
	w	e	w	
w	b	v	c	w
	w	d	w	
		w		

		w		
	w	e	w	
w	b	v	c	w
	w	d	w	
		w		

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} v \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ v \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ v \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([v]) & \text{with } \begin{array}{c} a \\ a \end{array} \begin{array}{c} v \\ a \end{array} a, \begin{array}{c} a \\ a \end{array} \begin{array}{c} w \\ a \end{array} a
 \end{array}$$

We can use the following order to change a single v to w :

		w		
	w	e	w	
w	b	v	c	w
	w	d	w	
		w		

		w		
	w	a	w	
w	b	v	c	w
	w	d	w	
		w		

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} v \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ v \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ v \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([v]) & \text{with } \begin{array}{c} a \\ a \end{array} \begin{array}{c} v \\ a \end{array} a, \begin{array}{c} a \\ a \end{array} \begin{array}{c} w \\ a \end{array} a
 \end{array}$$

We can use the following order to change a single v to w :

		w		
	w	e	w	
w	b	v	c	w
	w	d	w	
		w		

		w		
	w	a	w	
w	b	v	c	w
	w	d	w	
		w		

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} x \begin{array}{c} w \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} a \begin{array}{c} w \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} x w, \begin{array}{c} w \\ w \end{array} v a w \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} x w, \begin{array}{c} v \\ w \end{array} a w \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ v \end{array} x w, \begin{array}{c} w \\ v \end{array} a w \\
 V([v]) & \text{with } \begin{array}{c} a \\ a \end{array} v a, \begin{array}{c} a \\ a \end{array} w a
 \end{array}$$

We can use the following order to change a single v to w :

		w		
	w	e	w	
w	b	v	c	w
	w	d	w	
		w		

		w		
	w	a	w	
w	b	v	c	w
	w	d	w	
		w		

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \ x \ v, \ w \ a \ v \\ w \end{array} \\
 V([vx]) & \text{with } \begin{array}{c} w \\ v \ x \ w, \ v \ a \ w \\ w \end{array} \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \ x \ w, \ w \ a \ w \\ w \end{array} \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ w \ x \ w, \ w \ a \ w \\ v \end{array} \\
 V([v]) & \text{with } \begin{array}{c} a \\ a \ v \ a, \ a \ w \ a \\ a \end{array}
 \end{array}$$

We can use the following order to change a single v to w :

		w		
	w	e	w	
w	b	v	c	w
	w	d	w	
		w		

		w		
	w	a	w	
w	a	v	c	w
	w	d	w	
		w		

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} x \begin{array}{c} w \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} a \begin{array}{c} w \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} x \begin{array}{c} w \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} a \begin{array}{c} w \\ w \end{array} v \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} x \begin{array}{c} v \\ w \end{array} w, \begin{array}{c} v \\ w \end{array} a \begin{array}{c} v \\ w \end{array} w \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ v \end{array} x \begin{array}{c} w \\ v \end{array} w, \begin{array}{c} w \\ v \end{array} a \begin{array}{c} w \\ v \end{array} w \\
 V([v]) & \text{with } \begin{array}{c} a \\ a \end{array} v \begin{array}{c} a \\ a \end{array} a, \begin{array}{c} a \\ a \end{array} w \begin{array}{c} a \\ a \end{array} a
 \end{array}$$

We can use the following order to change a single v to w :

		w		
	w	e	w	
w	b	v	c	w
	w	d	w	
		w		

		w		
	w	a	w	
w	a	v	c	w
	w	d	w	
		w		

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} x \begin{array}{c} w \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} a \begin{array}{c} w \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} x \begin{array}{c} w \\ w \end{array} w, \begin{array}{c} w \\ w \end{array} a \begin{array}{c} w \\ w \end{array} w \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} x \begin{array}{c} w \\ w \end{array} w, \begin{array}{c} v \\ w \end{array} a \begin{array}{c} w \\ w \end{array} w \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ v \end{array} x \begin{array}{c} w \\ w \end{array} w, \begin{array}{c} w \\ v \end{array} a \begin{array}{c} w \\ w \end{array} w \\
 V([v]) & \text{with } \begin{array}{c} a \\ a \end{array} v \begin{array}{c} a \\ a \end{array} w, \begin{array}{c} a \\ a \end{array} w \begin{array}{c} a \\ a \end{array} w
 \end{array}$$

We can use the following order to change a single v to w :

		w		
	w	e	w	
w	b	v	c	w
	w	d	w	
		w		

		w		
	w	a	w	
w	a	v	a	w
	w	d	w	
		w		

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} v \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ v \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ v \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([v]) & \text{with } \begin{array}{c} a \\ a \end{array} \begin{array}{c} v \\ a \end{array} a, \begin{array}{c} a \\ a \end{array} \begin{array}{c} w \\ a \end{array} a
 \end{array}$$

We can use the following order to change a single v to w :

		w		
	w	e	w	
w	b	v	c	w
	w	d	w	
		w		

		w		
	w	a	w	
w	a	v	a	w
	w	d	w	
		w		

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} v \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ v \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ v \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([v]) & \text{with } \begin{array}{c} a \\ a \end{array} \begin{array}{c} v \\ a \end{array} a, \begin{array}{c} a \\ a \end{array} \begin{array}{c} w \\ a \end{array} a
 \end{array}$$

We can use the following order to change a single v to w :

		w		
	w	e	w	
w	b	v	c	w
	w	d	w	
		w		

		w		
	w	a	w	
w	a	v	a	w
	w	a	w	
		w		

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} v \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ v \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ v \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([v]) & \text{with } \begin{array}{c} a \\ a \end{array} \begin{array}{c} v \\ v \end{array} \begin{array}{c} a \\ a \end{array}, \begin{array}{c} a \\ a \end{array} \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ a \end{array}
 \end{array}$$

We can use the following order to change a single v to w :

		w		
	w	e	w	
w	b	v	c	w
	w	d	w	
		w		

		w		
	w	a	w	
w	a	v	a	w
	w	a	w	
		w		

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} v \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ v \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ v \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([v]) & \text{with } \begin{array}{c} a \\ a \end{array} \begin{array}{c} v \\ v \end{array} \begin{array}{c} a \\ a \end{array}, \begin{array}{c} a \\ a \end{array} \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ a \end{array}
 \end{array}$$

We can use the following order to change a single v to w :

		w		
	w	e	w	
w	b	v	c	w
	w	d	w	
		w		

		w		
	w	a	w	
w	a	w	a	w
	w	a	w	
		w		

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} v \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ v \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ v \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([v]) & \text{with } \begin{array}{c} a \\ a \end{array} \begin{array}{c} v \\ a \end{array} a, \begin{array}{c} a \\ a \end{array} \begin{array}{c} w \\ a \end{array} a
 \end{array}$$

We can use the following order to change a single v to w :

		w		
	w	e	w	
w	b	v	c	w
	w	d	w	
		w		

		w		
	w	a	w	
w	a	w	a	w
	w	a	w	
		w		

An idea of the proof:

Assume $v \approx v$ and choose some $a \sim v$. For all $x \sim v$ make the following identifications

$$\begin{array}{ll}
 V([xv]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} v, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} v \\
 V([vx]) & \text{with } \begin{array}{c} w \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} v \\ x \end{smallmatrix}]) & \text{with } \begin{array}{c} v \\ w \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} v \\ w \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([\begin{smallmatrix} x \\ v \end{smallmatrix}]) & \text{with } \begin{array}{c} w \\ v \end{array} \begin{array}{c} x \\ w \end{array} w, \begin{array}{c} w \\ v \end{array} \begin{array}{c} a \\ w \end{array} w \\
 V([v]) & \text{with } \begin{array}{c} a \\ a \end{array} \begin{array}{c} v \\ a \end{array} a, \begin{array}{c} a \\ a \end{array} \begin{array}{c} w \\ a \end{array} a
 \end{array}$$

We can use the following order to change a single v to w :

		w		
	w	e	w	
w	b	v	c	w
	w	d	w	
		w		

		w		
	w	e	w	
w	b	w	c	w
	w	d	w	
		w		

An idea of the proof:

Thus in the case when $v \approx v$ we can find an interaction which represents changing a single v to w in any configuration.

An idea of the proof:

Thus in the case when $v \approx v$ we can find an interaction which represents changing a single v to w in any configuration. Since $\text{Hom}(\mathbb{Z}^2, \mathcal{H} \setminus \{v\})$ is a Hammersley-Clifford space

An idea of the proof:

Thus in the case when $v \approx v$ we can find an interaction which represents changing a single v to w in any configuration. Since $\text{Hom}(\mathbb{Z}^2, \mathcal{H} \setminus \{v\})$ is a Hammersley-Clifford space we are done for this particular case.

An idea of the proof:

Thus in the case when $v \approx v$ we can find an interaction which represents changing a single v to w in any configuration. Since $\text{Hom}(\mathbb{Z}^2, \mathcal{H} \setminus \{v\})$ is a Hammersley-Clifford space we are done for this particular case.

If $v \sim v$ then the argument is slightly more involved.

Questions:

Questions:

- What happens when \mathcal{G} is not bipartite?

Questions:

- What happens when \mathcal{G} is not bipartite?
- Given a graph \mathcal{G} what are the graphs \mathcal{H} for which $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space?

Questions:

- What happens when \mathcal{G} is not bipartite?
- Given a graph \mathcal{G} what are the graphs \mathcal{H} for which $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space? Equivalently,

Questions:

- What happens when \mathcal{G} is not bipartite?
- Given a graph \mathcal{G} what are the graphs \mathcal{H} for which $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space? Equivalently, for what stiff graphs \mathcal{H} (ones which cannot be folded) is the space $\text{Hom}(\mathcal{G}, \mathcal{H})$ a Hammersley-Clifford space?

Questions:

- What happens when \mathcal{G} is not bipartite?
- Given a graph \mathcal{G} what are the graphs \mathcal{H} for which $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space? Equivalently, for what stiff graphs \mathcal{H} (ones which cannot be folded) is the space $\text{Hom}(\mathcal{G}, \mathcal{H})$ a Hammersley-Clifford space?

If \mathcal{H} is a single vertex with a loop

Questions:

- What happens when \mathcal{G} is not bipartite?
- Given a graph \mathcal{G} what are the graphs \mathcal{H} for which $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space? Equivalently, for what stiff graphs \mathcal{H} (ones which cannot be folded) is the space $\text{Hom}(\mathcal{G}, \mathcal{H})$ a Hammersley-Clifford space?

If \mathcal{H} is a single vertex with a loop or an edge

Questions:

- What happens when \mathcal{G} is not bipartite?
- Given a graph \mathcal{G} what are the graphs \mathcal{H} for which $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space? Equivalently, for what stiff graphs \mathcal{H} (ones which cannot be folded) is the space $\text{Hom}(\mathcal{G}, \mathcal{H})$ a Hammersley-Clifford space?

If \mathcal{H} is a single vertex with a loop or an edge then $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space.

Questions:

- What happens when \mathcal{G} is not bipartite?
- Given a graph \mathcal{G} what are the graphs \mathcal{H} for which $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space? Equivalently, for what stiff graphs \mathcal{H} (ones which cannot be folded) is the space $\text{Hom}(\mathcal{G}, \mathcal{H})$ a Hammersley-Clifford space?

If \mathcal{H} is a single vertex with a loop or an edge then $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space.

Thus if \mathcal{H} is dismantlable graph, a tree or a 4-cycle, then $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space.

Questions:

- What happens when \mathcal{G} is not bipartite?
- Given a graph \mathcal{G} what are the graphs \mathcal{H} for which $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space? Equivalently, for what stiff graphs \mathcal{H} (ones which cannot be folded) is the space $\text{Hom}(\mathcal{G}, \mathcal{H})$ a Hammersley-Clifford space?

If \mathcal{H} is a single vertex with a loop or an edge then $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space.

Thus if \mathcal{H} is dismantlable graph, a tree or a 4-cycle, then $\text{Hom}(\mathcal{G}, \mathcal{H})$ is a Hammersley-Clifford space.

Are there any more such stiff graphs \mathcal{H} in general?

Thank You!

.