

Four-Cycle Free Graphs and Entropy Minimality

Nishant Chandgotia

University of British Columbia

March, 2014

Outline

- Entropy Minimality and Hom Shifts
- Mixing Conditions and Entropy Minimality
- Measures of Maximal Entropy
- Rigidity and Flexibility in the Space of 3-Colourings.

Nearest Neighbour Shifts of Finite Type

Let \mathcal{A} be a finite alphabet set.

Nearest Neighbour Shifts of Finite Type

Let \mathcal{A} be a finite alphabet set. A shape F is any finite subset of \mathbb{Z}^d .

Nearest Neighbour Shifts of Finite Type

Let \mathcal{A} be a finite alphabet set. A shape F is any finite subset of \mathbb{Z}^d . A pattern is a function $f : F \longrightarrow \mathcal{A}$.

Nearest Neighbour Shifts of Finite Type

Let \mathcal{A} be a finite alphabet set. A shape F is any finite subset of \mathbb{Z}^d . A pattern is a function $f : F \longrightarrow \mathcal{A}$. Given \mathcal{F} , a list of patterns,

Nearest Neighbour Shifts of Finite Type

Let \mathcal{A} be a finite alphabet set. A shape F is any finite subset of \mathbb{Z}^d . A pattern is a function $f : F \longrightarrow \mathcal{A}$. Given \mathcal{F} , a list of patterns, a shift space $X_{\mathcal{F}}$ is given by

$$X_{\mathcal{F}} = \{x \in \mathcal{A}^{\mathbb{Z}^d} \mid \text{patterns from } \mathcal{F} \text{ do not occur in } x\}.$$

Nearest Neighbour Shifts of Finite Type

Let \mathcal{A} be a finite alphabet set. A shape F is any finite subset of \mathbb{Z}^d . A pattern is a function $f : F \longrightarrow \mathcal{A}$. Given \mathcal{F} , a list of patterns, a **shift space** $X_{\mathcal{F}}$ is given by

$$X_{\mathcal{F}} = \{x \in \mathcal{A}^{\mathbb{Z}^d} \mid \text{patterns from } \mathcal{F} \text{ do not occur in } x\}.$$

A **nearest neighbour shift of finite type** is a shift space such that the constraint set \mathcal{F} can be given by patterns on 'edges'.

Nearest Neighbour Shifts of Finite Type

Let \mathcal{A} be a finite alphabet set. A shape F is any finite subset of \mathbb{Z}^d . A pattern is a function $f : F \longrightarrow \mathcal{A}$. Given \mathcal{F} , a list of patterns, a **shift space** $X_{\mathcal{F}}$ is given by

$$X_{\mathcal{F}} = \{x \in \mathcal{A}^{\mathbb{Z}^d} \mid \text{patterns from } \mathcal{F} \text{ do not occur in } x\}.$$

A **nearest neighbour shift of finite type** is a shift space such that the constraint set \mathcal{F} can be given by patterns on 'edges'.

The **cylinder set** for the pattern $f \in \mathcal{A}^F$ is given by

Nearest Neighbour Shifts of Finite Type

Let \mathcal{A} be a finite alphabet set. A shape F is any finite subset of \mathbb{Z}^d . A pattern is a function $f : F \rightarrow \mathcal{A}$. Given \mathcal{F} , a list of patterns, a **shift space** $X_{\mathcal{F}}$ is given by

$$X_{\mathcal{F}} = \{x \in \mathcal{A}^{\mathbb{Z}^d} \mid \text{patterns from } \mathcal{F} \text{ do not occur in } x\}.$$

A **nearest neighbour shift of finite type** is a shift space such that the constraint set \mathcal{F} can be given by patterns on 'edges'.

The **cylinder set** for the pattern $f \in \mathcal{A}^F$ is given by

$$[f]_F = \{x \in X \mid x|_F = f\}.$$

Nearest Neighbour Shifts of Finite Type

Let \mathcal{A} be a finite alphabet set. A shape F is any finite subset of \mathbb{Z}^d . A pattern is a function $f : F \rightarrow \mathcal{A}$. Given \mathcal{F} , a list of patterns, a **shift space** $X_{\mathcal{F}}$ is given by

$$X_{\mathcal{F}} = \{x \in \mathcal{A}^{\mathbb{Z}^d} \mid \text{patterns from } \mathcal{F} \text{ do not occur in } x\}.$$

A **nearest neighbour shift of finite type** is a shift space such that the constraint set \mathcal{F} can be given by patterns on 'edges'.

The **cylinder set** for the pattern $f \in \mathcal{A}^F$ is given by

$$[f]_F = \{x \in X \mid x|_F = f\}.$$

\mathbb{Z}^d acts by translations(shifts) on the shift spaces.

Hom shifts

Given a finite undirected graph \mathcal{H} without multiple edges,

Hom shifts

Given a finite undirected graph \mathcal{H} without multiple edges, we can define a shift space $X_{\mathcal{H}} \subset \mathcal{H}^{\mathbb{Z}^d}$

Hom shifts

Given a finite undirected graph \mathcal{H} without multiple edges, we can define a shift space $X_{\mathcal{H}} \subset \mathcal{H}^{\mathbb{Z}^d}$ such that symbols on adjacent vertices form an edge in the graph \mathcal{H} .

Hom shifts

Given a finite undirected graph \mathcal{H} without multiple edges, we can define a shift space $X_{\mathcal{H}} \subset \mathcal{H}^{\mathbb{Z}^d}$ such that symbols on adjacent vertices form an edge in the graph \mathcal{H} .

Here every direction has the same constraint.

Hom shifts

Given a finite undirected graph \mathcal{H} without multiple edges, we can define a shift space $X_{\mathcal{H}} \subset \mathcal{H}^{\mathbb{Z}^d}$ such that symbols on adjacent vertices form an edge in the graph \mathcal{H} .

Here every direction has the same constraint.


Examples:

Hom shifts

Given a finite undirected graph \mathcal{H} without multiple edges, we can define a shift space $X_{\mathcal{H}} \subset \mathcal{H}^{\mathbb{Z}^d}$ such that symbols on adjacent vertices form an edge in the graph \mathcal{H} .

Here every direction has the same constraint.

Examples:(Hard Square model)



Graph \mathcal{H}

1	0	0	0	0
0	0	0	0	0
1	0	1	0	0
0	0	0	1	0
0	1	0	0	0


A Pattern

Hom shifts


Given a finite undirected graph \mathcal{H} without multiple edges, we can define a shift space $X_{\mathcal{H}} \subset \mathcal{H}^{\mathbb{Z}^d}$ such that symbols on adjacent vertices form an edge in the graph \mathcal{H} .

Here every direction has the same constraint.

Examples:(3-colourings)



Graph \mathcal{H}



A Pattern

Entropy

Entropy

The **language** $\mathcal{B}(X)$ of a shift space is the set of patterns appearing in elements of X .

Entropy

The **language** $\mathcal{B}(X)$ of a shift space is the set of patterns appearing in elements of X .

The **topological entropy** of X is defined as

Entropy

The **language** $\mathcal{B}(X)$ of a shift space is the set of patterns appearing in elements of X .

The **topological entropy** of X is defined as

$$h(X) := \lim_{n \rightarrow \infty} \frac{\log |\mathcal{B}(X) \cap \mathcal{A}^{\{1,2,\dots,n\}^d}|}{n^d}.$$

Entropy Minimality

Entropy Minimality

If $X \subset Y$ are shift spaces then

$$h_{top}(X) \leq h_{top}(Y).$$

Entropy Minimality

If $X \subset Y$ are shift spaces then

$$h_{top}(X) \leq h_{top}(Y).$$

A shift space Y is said to be **entropy minimal** if for all shift spaces $X \subsetneq Y$,

Entropy Minimality

If $X \subset Y$ are shift spaces then

$$h_{top}(X) \leq h_{top}(Y).$$

A shift space Y is said to be **entropy minimal** if for all shift spaces $X \subsetneq Y$, $h_{top}(X) < h_{top}(Y)$,

Entropy Minimality

If $X \subset Y$ are shift spaces then

$$h_{top}(X) \leq h_{top}(Y).$$

A shift space Y is said to be **entropy minimal** if for all shift spaces $X \subsetneq Y$, $h_{top}(X) < h_{top}(Y)$, that is, if we forbid any pattern from the language $\mathcal{B}(Y)$ the entropy will drop.

Entropy Minimality

If $X \subset Y$ are shift spaces then

$$h_{top}(X) \leq h_{top}(Y).$$

A shift space Y is said to be **entropy minimal** if for all shift spaces $X \subsetneq Y$, $h_{top}(X) < h_{top}(Y)$, that is, if we forbid any pattern from the language $\mathcal{B}(Y)$ the entropy will drop.

If a shift space X is entropy minimal and $h_{top}(X) = 0$ then it is minimal.

Entropy Minimality

If $X \subset Y$ are shift spaces then

$$h_{top}(X) \leq h_{top}(Y).$$

A shift space Y is said to be **entropy minimal** if for all shift spaces $X \subsetneq Y$, $h_{top}(X) < h_{top}(Y)$, that is, if we forbid any pattern from the language $\mathcal{B}(Y)$ the entropy will drop.

If a shift space X is entropy minimal and $h_{top}(X) = 0$ then it is minimal.

(*Quas and Trow '00*) Every shift space X contains an entropy minimal shift space $Y \subset X$ such that $h_{top}(X) = h_{top}(Y)$.

Main Result

Main Result

Theorem (Chandgotia, Meyerovitch '13)

Let C_n be an n -cycle for some integer $n \neq 4$. Then X_{C_n} is entropy minimal.

Main Result

Theorem (Chandgotia, Meyerovitch '13)

Let C_n be an n -cycle for some integer $n \neq 4$. Then X_{C_n} is entropy minimal.

Theorem (Chandgotia '14; In preperation)

If \mathcal{H} is a connected graph without 4-cycles then $X_{\mathcal{H}}$ is entropy minimal.

Main Result

Theorem (Chandgotia, Meyerovitch '13)

Let C_n be an n -cycle for some integer $n \neq 4$. Then X_{C_n} is entropy minimal.

Theorem (Chandgotia '14; In preperation)

If \mathcal{H} is a connected graph without 4-cycles then $X_{\mathcal{H}}$ is entropy minimal.

Remark: We will concentrate on X_{C_3} , the space of all 3-colourings.

Mixing Conditions and Entropy Minimality

Mixing Conditions and Entropy Minimality


Transitivity:

Mixing Conditions and Entropy Minimality

Transitivity: For all $a, b \in \mathcal{B}(X)$ there exists $N_{a,b} \in \mathbb{Z}^d$ such that

Mixing Conditions and Entropy Minimality


Transitivity: For all $a, b \in \mathcal{B}(X)$ there exists $N_{a,b} \in \mathbb{Z}^d$ such that



is a pattern in $\mathcal{B}(X)$.

Mixing Conditions and Entropy Minimality

Transitivity: For all $a, b \in \mathcal{B}(X)$ there exists $N_{a,b} \in \mathbb{Z}^d$ such that



is a pattern in $\mathcal{B}(X)$.

(Coven and Smítal '93) If a shift space is entropy minimal then it is topologically transitive.

Mixing Conditions and Entropy Minimality


Block-Gluing:

Mixing Conditions and Entropy Minimality

Block-Gluing: There exists a $N \in \mathbb{N}$ such that for all rectangular patterns $a, b \in \mathcal{B}(X)$

Mixing Conditions and Entropy Minimality


Block-Gluing: There exists a $N \in \mathbb{N}$ such that for all rectangular patterns $a, b \in \mathcal{B}(X)$



is a pattern in $\mathcal{B}(X)$.

Mixing Conditions and Entropy Minimality

Block-Gluing: There exists a $N \in \mathbb{N}$ such that for all rectangular patterns $a, b \in \mathcal{B}(X)$




is a pattern in $\mathcal{B}(X)$.

Block-gluing shift spaces are transitive.

Mixing Conditions and Entropy Minimality

Block-Gluing: There exists a $N \in \mathbb{N}$ such that for all rectangular patterns $a, b \in \mathcal{B}(X)$




is a pattern in $\mathcal{B}(X)$.

Block-gluing shift spaces are transitive.

(Boyle, Pavlov and Schraudner '09) There exists a block-gluing shift space which is not entropy minimal.

A Block-Gluing Shift Which Is Not Entropy Minimal

Symbols




A Pattern

1	15	20	49		56	115
119	19	30	17		19	77
					22	40
67	59	46	117	57	46	17
5	8	73				
24	57	79		43	82	108
79	64	52		93	37	9

A Block-Gluing Shift Which Is Not Entropy Minimal

Symbols



A Pattern

1	15	20	49		56	115
119	19	30	17		19	77
					22	40
67	59	46	117	57	46	17
5	8	73				
24	57	79		43	82	108
79	64	52		93	37	9

The symbols with arrows do not contribute any entropy.

Mixing Conditions and Entropy Minimality


Strongly Irreducible:

Mixing Conditions and Entropy Minimality

Strongly Irreducible: There exists an $N \in \mathbb{N}$ such that for all patterns $a, b \in \mathcal{B}(X)$

Mixing Conditions and Entropy Minimality


Strongly Irreducible: There exists an $N \in \mathbb{N}$ such that for all patterns $a, b \in \mathcal{B}(X)$



is a pattern in $\mathcal{B}(X)$.

Mixing Conditions and Entropy Minimality

Strongly Irreducible: There exists an $N \in \mathbb{N}$ such that for all patterns $a, b \in \mathcal{B}(X)$




is a pattern in $\mathcal{B}(X)$.

Strongly irreducible shift spaces are block-gluing and hence transitive.

Mixing Conditions and Entropy Minimality

Strongly Irreducible: There exists an $N \in \mathbb{N}$ such that for all patterns $a, b \in \mathcal{B}(X)$




is a pattern in $\mathcal{B}(X)$.

Strongly irreducible shift spaces are block-gluing and hence transitive.

(Schraudner '09) Every strongly irreducible shift space is entropy minimal.

Mixing Conditions and Entropy Minimality



Measures of Maximal Entropy

Measures of Maximal Entropy

For a given shift-invariant Borel probability measure μ on a shift space we denote its **measure-theoretic entropy** by h_μ .

Measures of Maximal Entropy

For a given shift-invariant Borel probability measure μ on a shift space we denote its **measure-theoretic entropy** by h_μ .

(Variational Principle) Given a shift space X

$$\sup_{\text{supp}(\mu) \subset X} h_\mu = h_{\text{top}}(X).$$

Measures of Maximal Entropy

For a given shift-invariant Borel probability measure μ on a shift space we denote its **measure-theoretic entropy** by h_μ .

(Variational Principle) Given a shift space X

$$\sup_{\text{supp}(\mu) \subset X} h_\mu = h_{\text{top}}(X).$$

Moreover the supremum is achieved.

Measures of Maximal Entropy

For a given shift-invariant Borel probability measure μ on a shift space we denote its **measure-theoretic entropy** by h_μ .

(Variational Principle) Given a shift space X

$$\sup_{\text{supp}(\mu) \subset X} h_\mu = h_{\text{top}}(X).$$

Moreover the supremum is achieved. Such measures are called **measures of maximal entropy**.

Measures of Maximal Entropy

For a given shift-invariant Borel probability measure μ on a shift space we denote its **measure-theoretic entropy** by h_μ .

(Variational Principle) Given a shift space X

$$\sup_{\text{supp}(\mu) \subset X} h_\mu = h_{\text{top}}(X).$$

Moreover the supremum is achieved. Such measures are called **measures of maximal entropy**.

If μ is a measure of maximal entropy for X then $h_\mu = h_{\text{Top}}(X)$.

Measures of Maximal Entropy

For a given shift-invariant Borel probability measure μ on a shift space we denote its **measure-theoretic entropy** by h_μ .

(Variational Principle) Given a shift space X

$$\sup_{\text{supp}(\mu) \subset X} h_\mu = h_{\text{top}}(X).$$

Moreover the supremum is achieved. Such measures are called **measures of maximal entropy**.

If μ is a measure of maximal entropy for X then $h_\mu = h_{\text{Top}}(X)$. Therefore, a shift space X is entropy minimal if and only if for every measure of maximal entropy μ , $\text{supp}(\mu) = X$.

Measures of Maximal Entropy and Gibbs Measures

Measures of Maximal Entropy and Gibbs Measures

(Lanford and Ruelle '69-Special case):

Measures of Maximal Entropy and Gibbs Measures

(Lanford and Ruelle '69-Special case): If X is a shift of finite type then for any measure of maximal entropy μ ,

Measures of Maximal Entropy and Gibbs Measures

(*Lanford and Ruelle '69-Special case*): If X is a shift of finite type then for any measure of maximal entropy μ , finite set A ,
 $\partial A \subset B \subset A^c$

Measures of Maximal Entropy and Gibbs Measures

(*Lanford and Ruelle '69-Special case*): If X is a shift of finite type then for any measure of maximal entropy μ , finite set A , $\partial A \subset B \subset A^c$ and $[b]_B$ of positive measure

Measures of Maximal Entropy and Gibbs Measures

(Lanford and Ruelle '69-Special case): If X is a shift of finite type then for any measure of maximal entropy μ , finite set A , $\partial A \subset B \subset A^c$ and $[b]_B$ of positive measure

$$\mu([a]_A \mid [b]_B)$$

is uniform over all patterns $a \in \mathcal{B}(X) \cap \mathcal{A}^A$ satisfying $ab \in \mathcal{B}(X)$.

Measures of Maximal Entropy and Gibbs Measures

(Lanford and Ruelle '69-Special case): If X is a shift of finite type then for any measure of maximal entropy μ , finite set A , $\partial A \subset B \subset A^c$ and $[b]_B$ of positive measure

$$\mu([a]_A \mid [b]_B)$$

is uniform over all patterns $a \in \mathcal{B}(X) \cap \mathcal{A}^A$ satisfying $ab \in \mathcal{B}(X)$.

Such measures are called uniform Gibbs measures.

Measures of Maximal Entropy and Gibbs Measures

(Lanford and Ruelle '69-Special case): If X is a shift of finite type then for any measure of maximal entropy μ , finite set A , $\partial A \subset B \subset A^c$ and $[b]_B$ of positive measure

$$\mu([a]_A \mid [b]_B)$$

is uniform over all patterns $a \in \mathcal{B}(X) \cap \mathcal{A}^A$ satisfying $ab \in \mathcal{B}(X)$.

Such measures are called uniform Gibbs measures.

For example, if μ is the measure of maximal entropy in the hard square model then

Measures of Maximal Entropy and Gibbs Measures

(Lanford and Ruelle '69-Special case): If X is a shift of finite type then for any measure of maximal entropy μ , finite set A , $\partial A \subset B \subset A^c$ and $[b]_B$ of positive measure

$$\mu([a]_A \mid [b]_B)$$

is uniform over all patterns $a \in \mathcal{B}(X) \cap \mathcal{A}^A$ satisfying $ab \in \mathcal{B}(X)$.

Such measures are called uniform Gibbs measures.

For example, if μ is the measure of maximal entropy in the hard square model then

$$\mu([0]_0 \mid \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}_{\partial 0}) = \mu([1]_0 \mid \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}_{\partial 0}) = \frac{1}{2}.$$

Strongly Irreducible Implies Entropy Minimal

Strongly Irreducible Implies Entropy Minimal

(Meester and Steif '01)

Strongly Irreducible Implies Entropy Minimal

(*Meester and Steif '01*) Every strongly irreducible shift of finite type is entropy minimal.

Strongly Irreducible Implies Entropy Minimal

(Meester and Steif '01) Every strongly irreducible shift of finite type is entropy minimal.

Proof:

Strongly Irreducible Implies Entropy Minimal

(Meester and Steif '01) Every strongly irreducible shift of finite type is entropy minimal.

Proof: Suppose X is a strongly irreducible shift of finite type with a measure of maximal entropy μ .

Strongly Irreducible Implies Entropy Minimal

(Meester and Steif '01) Every strongly irreducible shift of finite type is entropy minimal.

Proof: Suppose X is a strongly irreducible shift of finite type with a measure of maximal entropy μ . We want to prove $\text{supp}(\mu) = X$.

Strongly Irreducible Implies Entropy Minimal

(Meester and Steif '01) Every strongly irreducible shift of finite type is entropy minimal.

Proof: Suppose X is a strongly irreducible shift of finite type with a measure of maximal entropy μ . We want to prove $\text{supp}(\mu) = X$. Consider a pattern $a \in \mathcal{B}(X)$ and $x \in \text{supp}(\mu)$.

Strongly Irreducible Implies Entropy Minimal


(Meester and Steif '01) Every strongly irreducible shift of finite type is entropy minimal.

Proof: Suppose X is a strongly irreducible shift of finite type with a measure of maximal entropy μ . We want to prove $\text{supp}(\mu) = X$. Consider a pattern $a \in \mathcal{B}(X)$ and $x \in \text{supp}(\mu)$. Since X is strongly irreducible

Strongly Irreducible Implies Entropy Minimal

(Meester and Steif '01) Every strongly irreducible shift of finite type is entropy minimal.

Proof: Suppose X is a strongly irreducible shift of finite type with a measure of maximal entropy μ . We want to prove $\text{supp}(\mu) = X$. Consider a pattern $a \in \mathcal{B}(X)$ and $x \in \text{supp}(\mu)$. Since X is strongly irreducible




is 'allowed' in X .

Strongly Irreducible Implies Entropy Minimal

(Meester and Steif '01) Every strongly irreducible shift of finite type is entropy minimal.

Proof: Suppose X is a strongly irreducible shift of finite type with a measure of maximal entropy μ . We want to prove $\text{supp}(\mu) = X$. Consider a pattern $a \in \mathcal{B}(X)$ and $x \in \text{supp}(\mu)$. Since X is strongly irreducible




is 'allowed' in X . Since μ is a measure of maximal entropy it is uniform Gibbs.

Strongly Irreducible Implies Entropy Minimal

(Meester and Steif '01) Every strongly irreducible shift of finite type is entropy minimal.

Proof: Suppose X is a strongly irreducible shift of finite type with a measure of maximal entropy μ . We want to prove $\text{supp}(\mu) = X$. Consider a pattern $a \in \mathcal{B}(X)$ and $x \in \text{supp}(\mu)$. Since X is strongly irreducible




is 'allowed' in X . Since μ is a measure of maximal entropy it is uniform Gibbs. Together with $x \in \text{supp}(\mu)$ we get $\mu([a]) > 0$.

Strongly Irreducible Implies Entropy Minimal

(Meester and Steif '01) Every strongly irreducible shift of finite type is entropy minimal.

Proof: Suppose X is a strongly irreducible shift of finite type with a measure of maximal entropy μ . We want to prove $\text{supp}(\mu) = X$. Consider a pattern $a \in \mathcal{B}(X)$ and $x \in \text{supp}(\mu)$. Since X is strongly irreducible




is 'allowed' in X . Since μ is a measure of maximal entropy it is uniform Gibbs. Together with $x \in \text{supp}(\mu)$ we get $\mu([a]) > 0$. Therefore $\text{supp}(\mu) = X$.

Strongly Irreducible Implies Entropy Minimal

(Meester and Steif '01) Every strongly irreducible shift of finite type is entropy minimal.

Proof: Suppose X is a strongly irreducible shift of finite type with a measure of maximal entropy μ . We want to prove $\text{supp}(\mu) = X$. Consider a pattern $a \in \mathcal{B}(X)$ and $x \in \text{supp}(\mu)$. Since X is strongly irreducible



is 'allowed' in X . Since μ is a measure of maximal entropy it is uniform Gibbs. Together with $x \in \text{supp}(\mu)$ we get $\mu([a]) > 0$. Therefore $\text{supp}(\mu) = X$. This completes the proof.

Mixing Properties of the Space of 3-colourings

Mixing Properties of the Space of 3-colourings

X_{C_3} is transitive but not block-gluing.

Mixing Properties of the Space of 3-colourings

X_{C_3} is transitive but not block-gluing.

1	2	0	2		2	0	2	0
2	0	2	0		1	2	0	2
0	2	0	2		2	0	2	1
1	0	2	0		1	2	1	0
2	1	0	2		2	1	0	2
0	2	1	0		1	0	2	1
<div><div>a</div><div>b</div></div>								

Mixing Properties of the Space of 3-colourings

X_{C_3} is transitive but not block-gluing.

1	2	0	2	1	2	0	2	0
2	0	2	0	2	1	2	0	2
0	2	0	2	1	2	0	2	1
1	0	2	0	2	1	2	1	0
2	1	0	2	1	2	1	0	2
0	2	1	0	2	1	0	2	1

Distance depends on the size of
a and b

Mixing Properties of the Space of 3-colourings

X_{C_3} is transitive but not block-gluing.

1	2	0	2	1	2	0	2	0
2	0	2	0	2	1	2	0	2
0	2	0	2	1	2	0	2	1
1	0	2	0	2	1	2	1	0
2	1	0	2	1	2	1	0	2
0	2	1	0	2	1	0	2	1

Distance depends on the size of
a and b

Yet, X_{C_3} is entropy minimal.

Proof Components: Height Functions and X_{C_3}

Proof Components: Height Functions and X_{C_3}

A **height function** is function $h : \mathbb{Z}^d \longrightarrow \mathbb{Z}$ such that

$$|h(v) - h(w)| = 1$$

for all adjacent vertices v and w .

Proof Components: Height Functions and X_{C_3}

A **height function** is function $h : \mathbb{Z}^d \longrightarrow \mathbb{Z}$ such that

$$|h(v) - h(w)| = 1$$

for all adjacent vertices v and w . It can also be seen as an element of $X_{\mathbb{Z}}$.

Proof Components: Height Functions and X_{C_3}

A **height function** is function $h : \mathbb{Z}^d \longrightarrow \mathbb{Z}$ such that

$$|h(v) - h(w)| = 1$$

for all adjacent vertices v and w . It can also be seen as an element of $X_{\mathbb{Z}}$.

If h is a height function then $h \bmod 3$ is an element of X_{C_3} .

Proof Components: Height Functions and X_{C_3}

A **height function** is function $h : \mathbb{Z}^d \longrightarrow \mathbb{Z}$ such that

$$|h(v) - h(w)| = 1$$

for all adjacent vertices v and w . It can also be seen as an element of $X_{\mathbb{Z}}$.

If h is a height function then $h \bmod 3$ is an element of X_{C_3} . This sets up a correspondence between height functions and elements of X_{C_3} .


Proof Components: Height Functions and X_{C_3}

A **height function** is function $h : \mathbb{Z}^d \longrightarrow \mathbb{Z}$ such that

$$|h(v) - h(w)| = 1$$

for all adjacent vertices v and w . It can also be seen as an element of $X_{\mathbb{Z}}$.

If h is a height function then $h \bmod 3$ is an element of X_{C_3} . This sets up a correspondence between height functions and elements of X_{C_3} .



Height Function


Proof Components: Height Functions and X_{C_3}

A **height function** is function $h : \mathbb{Z}^d \rightarrow \mathbb{Z}$ such that

$$|h(v) - h(w)| = 1$$

for all adjacent vertices v and w . It can also be seen as an element of $X_{\mathbb{Z}}$.

If h is a height function then $h \bmod 3$ is an element of X_{C_3} . This sets up a correspondence between height functions and elements of X_{C_3} .



Height Function

Pattern in X_{C_3}

Proof Components: Slopes

Proof Components: Slopes

Let e_1, e_2, \dots, e_d be the coordinate basis for \mathbb{Z}^d .

Proof Components: Slopes

Let e_1, e_2, \dots, e_d be the coordinate basis for \mathbb{Z}^d . Given a height function h

Proof Components: Slopes

Let e_1, e_2, \dots, e_d be the coordinate basis for \mathbb{Z}^d . Given a height function h the **slope** in the direction e_i is given by

Proof Components: Slopes

Let e_1, e_2, \dots, e_d be the coordinate basis for \mathbb{Z}^d . Given a height function h the **slope** in the direction e_i is given by

$$\lim_{n \rightarrow \infty} \frac{h(ne_i) - h(0)}{n}.$$

Proof Components: Slopes

Let e_1, e_2, \dots, e_d be the coordinate basis for \mathbb{Z}^d . Given a height function h the **slope** in the direction e_i is given by

$$\lim_{n \rightarrow \infty} \frac{h(ne_i) - h(0)}{n}.$$

The slope is always between -1 and 1 in every direction.

Proof Components: Slopes

Let e_1, e_2, \dots, e_d be the coordinate basis for \mathbb{Z}^d . Given a height function h the **slope** in the direction e_i is given by

$$\lim_{n \rightarrow \infty} \frac{h(ne_i) - h(0)}{n}.$$

The slope is always between -1 and 1 in every direction.

Given an ergodic measure μ on X_{C_3} ,

Proof Components: Slopes

Let e_1, e_2, \dots, e_d be the coordinate basis for \mathbb{Z}^d . Given a height function h the **slope** in the direction e_i is given by

$$\lim_{n \rightarrow \infty} \frac{h(ne_i) - h(0)}{n}.$$

The slope is always between -1 and 1 in every direction.

Given an ergodic measure μ on X_{C_3} , the slope is a constant μ -almost everywhere in every direction.

Proof Components: Slopes

Let e_1, e_2, \dots, e_d be the coordinate basis for \mathbb{Z}^d . Given a height function h the **slope** in the direction e_i is given by

$$\lim_{n \rightarrow \infty} \frac{h(ne_i) - h(0)}{n}.$$

The slope is always between -1 and 1 in every direction.

Given an ergodic measure μ on X_{C_3} , the slope is a constant μ -almost everywhere in every direction.

Note that the slope may be different in different directions.

Proof Components: Steep Slopes and Rigidity

Proof Components: Steep Slopes and Rigidity

Let μ be an ergodic measure of maximal entropy for X_{C_3} .

Proof Components: Steep Slopes and Rigidity

Let μ be an ergodic measure of maximal entropy for X_{C_3} . If the slope for μ is 1 or -1 in some direction

Proof Components: Steep Slopes and Rigidity

Let μ be an ergodic measure of maximal entropy for X_{C_3} . If the slope for μ is 1 or -1 in some direction then the height function cannot be changed at any place to obtain another height function.

Proof Components: Steep Slopes and Rigidity

Let μ be an ergodic measure of maximal entropy for X_{C_3} . If the slope for μ is 1 or -1 in some direction then the height function cannot be changed at any place to obtain another height function.

0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2

Proof Components: Steep Slopes and Rigidity

Let μ be an ergodic measure of maximal entropy for X_{C_3} . If the slope for μ is 1 or -1 in some direction then the height function cannot be changed at any place to obtain another height function.

0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2

If X_{frozen} is the space of such elements then

Proof Components: Steep Slopes and Rigidity

Let μ be an ergodic measure of maximal entropy for X_{C_3} . If the slope for μ is 1 or -1 in some direction then the height function cannot be changed at any place to obtain another height function.

0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2

If X_{frozen} is the space of such elements then $h_{Top}(X_{frozen}) = 0$.

Proof Components: Steep Slopes and Rigidity

Let μ be an ergodic measure of maximal entropy for X_{C_3} . If the slope for μ is 1 or -1 in some direction then the height function cannot be changed at any place to obtain another height function.

0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2

If X_{frozen} is the space of such elements then $h_{Top}(X_{frozen}) = 0$. But $h_\mu = h_{Top}(X_{C_3}) > 0$.

Proof Components: Steep Slopes and Rigidity

Let μ be an ergodic measure of maximal entropy for X_{C_3} . If the slope for μ is 1 or -1 in some direction then the height function cannot be changed at any place to obtain another height function.

0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2

If X_{frozen} is the space of such elements then $h_{Top}(X_{frozen}) = 0$. But $h_\mu = h_{Top}(X_{C_3}) > 0$. Thus the slope for μ

Proof Components: Steep Slopes and Rigidity

Let μ be an ergodic measure of maximal entropy for X_{C_3} . If the slope for μ is 1 or -1 in some direction then the height function cannot be changed at any place to obtain another height function.

0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2
2	0	1	2	0	1	2	0	1	2	0	1
0	1	2	0	1	2	0	1	2	0	1	2

If X_{frozen} is the space of such elements then $h_{Top}(X_{frozen}) = 0$. But $h_{\mu} = h_{Top}(X_{C_3}) > 0$. Thus the slope for μ is strictly between -1 and 1 in every direction.

Proof Components: Gentle Slopes and Flexibility

Proof Components: Gentle Slopes and Flexibility

Given any height function h_1 on a ball D_n in \mathbb{Z}^d

Proof Components: Gentle Slopes and Flexibility

Given any height function h_1 on a ball D_n in \mathbb{Z}^d and a height function h_2 on \mathbb{Z}^d with slope strictly between 1 and -1

Proof Components: Gentle Slopes and Flexibility

Given any height function h_1 on a ball D_n in \mathbb{Z}^d and a height function h_2 on \mathbb{Z}^d with slope strictly between 1 and -1 we can choose an $N \in \mathbb{N}$ and a height function h

Proof Components: Gentle Slopes and Flexibility

Given any height function h_1 on a ball D_n in \mathbb{Z}^d and a height function h_2 on \mathbb{Z}^d with slope strictly between 1 and -1 we can choose an $N \in \mathbb{N}$ and a height function h which is equal to h_1 on D_n and h_2 on D_N^c .

Proof Components: Gentle Slopes and Flexibility


Given any height function h_1 on a ball D_n in \mathbb{Z}^d and a height function h_2 on \mathbb{Z}^d with slope strictly between 1 and -1 we can choose an $N \in \mathbb{N}$ and a height function h which is equal to h_1 on D_n and h_2 on D_N^c . The choice of N depends only on n

Proof Components: Gentle Slopes and Flexibility

Given any height function h_1 on a ball D_n in \mathbb{Z}^d and a height function h_2 on \mathbb{Z}^d with slope strictly between 1 and -1 we can choose an $N \in \mathbb{N}$ and a height function h which is equal to h_1 on D_n and h_2 on D_N^c . The choice of N depends only on n and the slope of h_2 .


Proof Components: Gentle Slopes and Flexibility

Given any height function h_1 on a ball D_n in \mathbb{Z}^d and a height function h_2 on \mathbb{Z}^d with slope strictly between 1 and -1 we can choose an $N \in \mathbb{N}$ and a height function h which is equal to h_1 on D_n and h_2 on D_N^c . The choice of N depends only on n and the slope of h_2 .




Proof Components: Gentle Slopes and Flexibility

Given any height function h_1 on a ball D_n in \mathbb{Z}^d and a height function h_2 on \mathbb{Z}^d with slope strictly between 1 and -1 we can choose an $N \in \mathbb{N}$ and a height function h which is equal to h_1 on D_n and h_2 on D_N^c . The choice of N depends only on n and the slope of h_2 .




Proof Components: Gentle Slopes and Flexibility

Given any height function h_1 on a ball D_n in \mathbb{Z}^d and a height function h_2 on \mathbb{Z}^d with slope strictly between 1 and -1 we can choose an $N \in \mathbb{N}$ and a height function h which is equal to h_1 on D_n and h_2 on D_N^c . The choice of N depends only on n and the slope of h_2 .



Proof Components: Gentle Slopes and Flexibility

Given any height function h_1 on a ball D_n in \mathbb{Z}^d and a height function h_2 on \mathbb{Z}^d with slope strictly between 1 and -1 we can choose an $N \in \mathbb{N}$ and a height function h which is equal to h_1 on D_n and h_2 on D_N^c . The choice of N depends only on n and the slope of h_2 .



Proof Components: Wrapping it up

Proof Components: Wrapping it up

Height functions to 3-colourings:

Proof Components: Wrapping it up

Height functions to 3-colourings: Suppose μ is an ergodic measure of maximal entropy for X_{C_3} .

Proof Components: Wrapping it up

Height functions to 3-colourings: Suppose μ is an ergodic measure of maximal entropy for X_{C_3} . Then the slope for μ in every direction lies strictly between

Proof Components: Wrapping it up

Height functions to 3-colourings: Suppose μ is an ergodic measure of maximal entropy for X_{C_3} . Then the slope for μ in every direction lies strictly between -1 and 1 .

Proof Components: Wrapping it up


Height functions to 3-colourings: Suppose μ is an ergodic measure of maximal entropy for X_{C_3} . Then the slope for μ in every direction lies strictly between -1 and 1 . Consider $x \in \text{supp}(\mu)$ with such a slope

Proof Components: Wrapping it up

Height functions to 3-colourings: Suppose μ is an ergodic measure of maximal entropy for X_{C_3} . Then the slope for μ in every direction lies strictly between -1 and 1 . Consider $x \in \text{supp}(\mu)$ with such a slope and any pattern $a \in \mathcal{B}_{X_{C_3}}$ on D_n for some n .

Proof Components: Wrapping it up


Height functions to 3-colourings: Suppose μ is an ergodic measure of maximal entropy for X_{C_3} . Then the slope for μ in every direction lies strictly between -1 and 1 . Consider $x \in \text{supp}(\mu)$ with such a slope and any pattern $a \in \mathcal{B}_{X_{C_3}}$ on D_n for some n . Then



is 'allowed' in X .

Proof Components: Wrapping it up


Height functions to 3-colourings: Suppose μ is an ergodic measure of maximal entropy for X_{C_3} . Then the slope for μ in every direction lies strictly between -1 and 1 . Consider $x \in \text{supp}(\mu)$ with such a slope and any pattern $a \in \mathcal{B}_{X_{C_3}}$ on D_n for some n . Then



is 'allowed' in X . But μ is a uniform Gibbs measure.

Proof Components: Wrapping it up


Height functions to 3-colourings: Suppose μ is an ergodic measure of maximal entropy for X_{C_3} . Then the slope for μ in every direction lies strictly between -1 and 1 . Consider $x \in \text{supp}(\mu)$ with such a slope and any pattern $a \in \mathcal{B}_{X_{C_3}}$ on D_n for some n . Then



is 'allowed' in X . But μ is a uniform Gibbs measure. Together with $x \in \text{supp}(\mu)$ we get $\mu([a]_{D_n}) > 0$.

Proof Components: Wrapping it up

Height functions to 3-colourings: Suppose μ is an ergodic measure of maximal entropy for X_{C_3} . Then the slope for μ in every direction lies strictly between -1 and 1 . Consider $x \in \text{supp}(\mu)$ with such a slope and any pattern $a \in \mathcal{B}_{X_{C_3}}$ on D_n for some n . Then



is 'allowed' in X . But μ is a uniform Gibbs measure. Together with $x \in \text{supp}(\mu)$ we get $\mu([a]_{D_n}) > 0$. Thus $\text{supp}(\mu) = X_{C_3}$.

Therefore X_{C_3} is entropy minimal.

Therefore X_{C_3} is entropy minimal.

What if C_3 is replaced by another connected four-cycle free graph?

Therefore X_{C_3} is entropy minimal.

What if C_3 is replaced by another connected four-cycle free graph?
Notice that we were using the following commutative diagram:

Therefore X_{C_3} is entropy minimal.

What if C_3 is replaced by another connected four-cycle free graph?
Notice that we were using the following commutative diagram:

$$\begin{array}{ccc} & & \mathbb{Z} \\ \text{height function} \nearrow & & \downarrow \text{mod } 3 \\ \mathbb{Z}^d & \longrightarrow & C_3. \end{array}$$

Therefore X_{C_3} is entropy minimal.

What if C_3 is replaced by another connected four-cycle free graph?
Notice that we were using the following commutative diagram:

$$\begin{array}{ccc} & & \mathbb{Z} \\ \text{height function} \nearrow & & \downarrow \text{mod } 3 \\ \mathbb{Z}^d & \longrightarrow & C_3. \end{array}$$

If C_3 is replaced by a connected four-cycle free graph \mathcal{H}

Therefore X_{C_3} is entropy minimal.

What if C_3 is replaced by another connected four-cycle free graph? Notice that we were using the following commutative diagram:


$$\begin{array}{ccc} & & \mathbb{Z} \\ \text{height function} \nearrow & & \downarrow \text{mod } 3 \\ \mathbb{Z}^d & \longrightarrow & C_3. \end{array}$$

If C_3 is replaced by a connected four-cycle free graph \mathcal{H} then \mathbb{Z} is replaced by the universal cover of \mathcal{H} .

Conjecture: If \mathcal{H} is any connected graph then $X_{\mathcal{H}}$ is entropy minimal. (d=2)

Conjecture: If \mathcal{H} is any connected graph then $X_{\mathcal{H}}$ is entropy minimal. (d=2)

Question: What shift spaces are conjugate to $X_{\mathcal{H}}$ for some graph \mathcal{H} ?



Thank You!