

## Case discussion: Two years follow-up of RASL procedure for chronic scapholunate dissociation

By: Dr Hazem Al Hasan - Sr Specialist in DMC/ Syria/ abstract No. 54734, Dr Abdulkhaliq Al Qahtani- Consultant Orthopedics- DMC- KSA, Dr Fatema Abdulmohsen- R4 Saudi board resident

### Case discussion:

30 years old Saudi male, medically free, History of trauma to his left hand during playing football, fell down on pronated wrist 2 months prior to his presentation to OPD  
He heard click in left wrist, ROM not affected but he complains of pain with lifting heavy objects but unable to do pushups in the gym  
Initial x-rays and MRI:



### Diagnostic wrist arthroscopy:

Patient underwent diagnostic wrist arthroscopy and found to have:

- RADIOCARPAL ARTHRITIS & DETACHED LIGAMENTS
- TFCC INJURY WITH SYNOVITIS
- SCAPHOLUNATE ARTHRITIS
- MIDCARPAL SYNOVITIS & DEGENERATIVE CHANGES

### Therefore, plan was done to take the patient for open reduction and association of scaphoid and lunate (RASL procedure):

Z SHAPE DORSAL INCISION DONE, DISSECTION BETWEEN 3RD AND 4TH EXTENSOR COMPARTMENT DONE FOLLOWED BY OPENING OF THE CAPSULE AND EXPLORATION OF SCAPHOLUNATE LEGMENT FOUND TO BE RUPTURE AND IRREPARABLE, SO, DECORTICATION DONE, SCAPHOLUNATE FIXED TOGETHER AFTER RESTORATION OF THE CARPAL BONE ALLIGMENT USING JOYSTICK K.WIE AND FIXATION DONE USING HERBERT SCREW

### Post op x-rays:



### x-rays 1 year post op:



Clinically, patient improved his ROM and hand grip:



One-year post op, patients' grip is almost equal bilaterally:



### Conclusion:

scapholunate dissociation is the most common cause of wrist instability. Reduction and association of scaphoid and lunate (RASL procedure) is found to be safe and effective treatment of chronic static scapholunate dissociation. It realigns the scaphoid and lunate, restore the function, reduce the pain and appears to be robust over time