







**Takeaway:-** COVID-19 confirmed cases varied significantly across WHO regions, with the Americas reporting the highest number of confirmed cases.




**Takeaway:** COVID-19 deaths varied substantially across WHO regions, with some regions experiencing a much higher mortality burden than others.




**Takeaway:** COVID-19 recoveries varied across WHO regions, reflecting differences in the reported recovery burden among regions.



**Takeaway:-** Active COVID-19 cases varied across WHO regions, indicating differences in the ongoing reported case burden.



**Takeaway:-** Confirmed COVID-19 cases were concentrated among a smaller group of countries, showing substantial differences in reported case counts.




**Takeaway:-** COVID-19 deaths were concentrated among a smaller number of countries, with substantial differences in reported mortality.

# Covid-19 Mortality & Recovery Analysis

Top 10 Countries by COVID-19 Deaths


Sum of Deaths by Country/Region



**Takeaway:-** COVID-19 deaths were concentrated among a smaller number of countries, with substantial differences in reported mortality.

Sum of Recovered

by WHO Region



**Takeaway:-** Reported recoveries varied substantially across WHO regions, reflecting differences in the scale of reported cases and outcomes.

# Key Insights & Recommendations

## MORTALITY CONCENTRATION

COVID-19 deaths were highly concentrated in a small number of countries, indicating significant differences in reported mortality.

The United States recorded the highest reported deaths among the countries shown.

## RECOVERY PATTERN

Reported recoveries varied substantially across WHO regions, reflecting differences in the scale of reported cases and outcomes.

The Americas and Europe recorded the highest reported recovery counts among the WHO regions.

## KEY RECOMMENDATION

Focus on high-burden countries and regions for deeper analysis, resource planning, and targeted public-health interventions.

Prioritize data-driven monitoring and targeted resource allocation in high-burden areas.