

§2.2: Sandwiched Singularities, Extended Graphs, and Decorated Germs

Nishtha Tikalal

August 23, 2026

1. NOTES FROM REMARK 6

1.1. The Idea

In geometry, two spaces can look identical from a “flexible” topological point of view, yet remain completely different from a “rigid” algebraic or analytic point of view.

Remark 2.6 establishes a crucial conceptual bridge: when studying the **contact geometry** (the flexible boundary data) of a surface singularity, we can completely ignore the complicated **analytic structure** (the rigid exact equations) and focus purely on the discrete **combinatorics** of the resolution graph.

2. ANALYTIC VS. TOPOLOGICAL/CONTACT

Property	Analytic Type	Topological / Contact Type
Perspective	Rigid algebraic geometry (complex calculus, exact equations).	Flexible geometry (rubber-sheet topology, contact structures).
Data Kept	Continuous parameters, precise complex coordinates, angles of intersection.	Discrete data: graphs, vertex weights, intersection counts.
Uniqueness	Not unique. Many different complex equations yield the same graph.	Unique. The contact manifold (Y, ξ) is locked in by the graph.

Table 1: Comparison of Analytic and Contact Viewpoints.

3. KEY CONCEPTS

3.1. The Resolution Graph G is a “Shared Skeleton”

Suppose you have two completely different complex surface singularities, $(X, 0)$ and $(X', 0)$, defined by distinct algebraic equations.

- By a theorem of Neumann, if their 3-dimensional boundaries (links) look topologically identical, they **must share the exact same minimal resolution graph G** .
- However, Laufer proved that if any vertex in G splits into more than 3 branches (valency > 3), the exact equations cannot be uniquely determined by the graph alone. Thus, X and X' are analytically distinct, but they share the same topological skeleton.

3.2. What Information is “Lost”?

Imagine drawing two lines intersecting at the origin on a piece of paper.

- **Analytic view:** It matters intensely whether the lines intersect at a 90° angle or a 10° angle. Changing the angle changes the continuous algebraic equations of the lines.
- **Topological view:** Both scenarios are just “two lines intersecting at a point.” The exact angle does not matter.

By looking only at the resolution graph, we **lose** the exact angles and analytic equations of the curve germ $\mathcal{C}$, but we **keep** the structural intersection data.

3.3. Why Contact Topology

The main punchline of the remark is that the **contact 3-manifold** (Y, ξ) **does not care about the lost analytic details**.

The contact structure is entirely determined by:

1. The discrete weights of the components.
2. The **pairwise orders of tangency** (how tightly clustered the branches are, combinatorially).

Therefore, if we want to prove something about the contact manifold or its Stein fillings (like in Theorem 7.8), we can freely pick whichever analytic representative is easiest to work with, or study the topology directly via the decorated graph G' .

4. SUMMARY

- **Lemma 2.5** is a tool that counts how many topological graph decorations (G') are mathematically possible.
- **Remark 2.6** is the justification for *why* we are allowed to use those graph decorations to study contact topology, explicitly guaranteeing that our smooth topological invariants won’t break down just because the underlying algebraic equations change.

REFERENCES

- [1] R. E. Gompf and A. I. Stipsicz, *4-Manifolds and Kirby Calculus*, Graduate Studies in Mathematics, vol. 20, American Mathematical Society, Providence, RI, 1999.
- [2] O. Plamenevskaya and L. Starkston, *Unexpected Stein fillings, rational surface singularities, and plane curve arrangements*, *Geometry & Topology*, 27(3), 2023, pp. 1083–1166. Preprint available at arXiv:2006.06631 [math.GT], <https://arxiv.org/abs/2006.06631>.
- [3] G.-M. Greuel, C. Lossen, and E. Shustin, *Introduction to Singularities and Deformations*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2007.
- [4] T. de Jong and D. van Straten, *Deformation theory of sandwiched singularities*, *Duke Mathematical Journal*, 95(3), 1998, pp. 451–522.