

# Introduction to probability for CS - Assignment 4

Michael art

March 12, 2024

## 1 Exercise 6

### 1.1 Q 3 b:

3. Find  $E(X)$  and  $V(X)$  if  $X$  is distributed according to the following probability functions:

(a)

$$p(x) = \begin{cases} \frac{e^{-\lambda} \lambda^x}{x!(1-e^{-\lambda})}, & x = 1, 2, \dots, (\lambda > 0), \\ 0, & \text{otherwise.} \end{cases}$$

(b)

$$p(x) = \begin{cases} \frac{(1-p)^x}{-x \ln p}, & x = 1, 2, \dots, (0 < p < 1), \\ 0, & \text{otherwise.} \end{cases}$$

**Solution:**

(b) (i)

$$E(X) = \sum_{x=0}^{\infty} p(x) \cdot x = \sum_{x=0}^{\infty} \frac{(1-p)^x}{-x \ln p} \cdot x = \sum_{x=1}^{\infty} \frac{(1-p)^x}{-x \ln p} \cdot x = -\frac{1}{\ln p} \sum_{x=1}^{\infty} (1-p)^x = \boxed{\frac{p-1}{p \ln p}}$$

(ii) (1)

$$\begin{aligned} E(X^2) &= \sum_{x=0}^{\infty} p(x) \cdot x^2 = \sum_{x=0}^{\infty} \frac{(1-p)^x}{-x \ln p} \cdot x^2 = \sum_{x=1}^{\infty} \frac{(1-p)^x}{-x \ln p} \cdot x^2 = \\ &= -\frac{1}{\ln p} \sum_{x=1}^{\infty} x(1-p)^x = -\frac{1}{\ln p} \cdot \frac{1-p}{p^2} = \frac{p-1}{p^2 \ln p} \end{aligned}$$

(2)

$$E(X)^2 = \left( \frac{p-1}{p \ln p} \right)^2 = \frac{p^2 - 2p + 1}{p^2 \ln^2 p}$$

In total:

$$V(X) = E(X^2) - E(X)^2 = \frac{p-1}{p^2 \ln p} - \frac{(p-1)^2}{p^2 \ln^2 p} = (p-1) \left( \frac{1}{p^2 \ln p} - \frac{p-1}{p^2 \ln^2 p} \right) = \boxed{(p-1) \frac{\ln p - p + 1}{p^2 \ln^2 p}}$$

### 1.2 Q 8:

8. Find the variance of the number of letters sent to the right destination in the absent-minded secretary problem.

**Solution:**

Let us define an indicator random variable:

$$X_i = \begin{cases} 1, & \text{if the } i\text{'th letter reached its destination} \\ 0, & \text{otherwise} \end{cases}$$

Where  $X = \sum_{i=1}^n X_i$ , therefore:

$$\begin{aligned} V(X) &= \text{cov}(X, X) = \text{cov}\left(\sum_{i=1}^n X_i, \sum_{i=1}^n X_i\right) = n \cdot V(X_1) + n(n-1) \cdot \text{cov}(X_1, X_2) = \\ &= n\left(E(X_1^2) - E(X_1)^2\right) + n(n-1)\left(E(X_1 X_2) - E(X_1)E(X_2)\right) = \\ &= n\left(\frac{1}{n} - \frac{1}{n^2}\right) + n(n-1)\left(P(X_1 = 1, X_2 = 1) - \frac{1}{n^2}\right) = 1 - \frac{1}{n} + n(n-1)\left(\frac{1}{n(n-1)} - \frac{1}{n^2}\right) = \\ &= 1 - \frac{1}{n} + 1 - 1 + \frac{1}{n} = \boxed{1} \end{aligned}$$

**2 Exercise 7****2.1 Q 1:**

1. The probability function of  $(X, Y)$  is given by the following table:

$y \backslash x$	1	2	3
1	1/12	1/6	1/12
2	1/6	1/4	1/12
3	1/12	1/12	0

Find:

- (a) the probability for  $X$  to be odd.
- (b) the probability for  $XY$  to be odd.
- (c)  $P(X + Y \geq 4)$ .
- (d)  $E(X + Y)$ .
- (e)  $V(X + Y)$ .

**Solution:**

(a)

$$P(X \text{ is odd}) = P(x = 1) + P(x = 3) = \frac{1}{12} + \frac{1}{6} + \frac{1}{12} + \frac{1}{12} + \frac{1}{12} = \boxed{\frac{1}{2}}$$

(b)

$$P(XY \text{ is odd}) = P(1, 1) + P(1, 3) + P(3, 1) + P(3, 3) = \frac{1}{12} + \frac{1}{12} + \frac{1}{12} + 0 = \boxed{\frac{1}{4}}$$

(c)

$$\begin{aligned}
P(X+Y \geq 4) &= P(1,3) + P(2,2) + P(2,3) + P(3,1) + P(3,2) + P(3,3) = \\
&= \frac{1}{12} + \frac{1}{4} + \frac{1}{12} + \frac{1}{12} + \frac{1}{12} + 0 = \boxed{\frac{7}{12}}
\end{aligned}$$

(d)

$$E(X+Y) = E(X) + E(Y) = 2\left(1 \cdot \left(\frac{1}{12} + \frac{1}{6} + \frac{1}{12}\right) + 2 \cdot \left(\frac{1}{6} + \frac{1}{4} + \frac{1}{12}\right) + 3 \cdot \left(\frac{1}{12} + \frac{1}{12}\right)\right) = \boxed{\frac{11}{3}}$$

(e)

$$V(X+Y) = E((X+Y)^2) - E(X+Y)^2 = 4 \cdot \frac{1}{12} + 9 \cdot \frac{1}{3} + 16 \cdot \frac{5}{12} + 25 \cdot \frac{1}{6} - \left(\frac{11}{3}\right)^2 = \boxed{\frac{13}{18}}$$

## 2.2 Q 5:

5. A die is rolled  $n$  times. Let  $X_1$  and  $X_2$  be the number of 1-s and of 2-s, respectively. Find:

- (a) the probability function of  $(X_1, X_2)$ .
- (b)  $\text{Cov}(X_1, X_2)$ .

**Solution:**

(a)

$$\begin{aligned}
P(X_1 = i, X_2 = j) &= \binom{n}{i} \left(\frac{1}{6}\right)^i \cdot \binom{n-i}{j} \left(\frac{1}{6}\right)^j \cdot \left(\frac{4}{6}\right)^{(n-i-j)} = \\
&= \boxed{\binom{n}{i, j, n-i-j} \left(\frac{1}{6}\right)^{(i+j)} \cdot \left(\frac{4}{6}\right)^{(n-i-j)}}
\end{aligned}$$

(b) (i)

$$X_1, X_2 \sim B\left(n, \frac{1}{6}\right) \implies E(X_1) = E(X_2) = n \cdot \frac{1}{6} = \frac{n}{6}$$

- (ii) Let  $X_{1,i}$  be an indicator random variable, where  $X_{1,i} = 1$  if the outcome on the  $i$ 'th position is 1 and 0 otherwise. Let  $X_{2,i}$  be an indicator random variable, where  $X_{2,i} = 1$  if the outcome on the  $i$ 'th position is 2 and 0 otherwise. We know that  $X_1 = \sum_{i=1}^n X_{1,i}$  and  $X_2 = \sum_{i=1}^n X_{2,i}$ . Therefore:

$$\begin{aligned}
E(X_1 X_2) &= E\left(\sum_{i=1}^n X_{1,i} \cdot \sum_{j=1}^n X_{2,i}\right) = \sum_{i,j=1}^n E(X_{1,i} X_{2,j}) = \\
&= \sum_{i \neq j}^n E(X_{1,i}) E(X_{2,j}) + \sum_{i=1}^n E(X_{1,i} X_{2,i}) \quad \begin{array}{l} \nearrow 0 \\ \nwarrow 0 \end{array} = \frac{n}{6} \frac{n-1}{6} = \frac{n(n-1)}{36}
\end{aligned}$$

$$\text{In total: } \text{Cov}(X_1, X_2) = E(X_1 X_2) - E(X_1) E(X_2) = \frac{n(n-1)}{36} - \frac{n^2}{36} = \boxed{-\frac{n}{36}}$$

### 3 Exercise 8:

#### 3.1 Q 2 a:

2. Let  $X_i$ ,  $i = 1, 2, \dots, m$ , be independent random variables distributed according to one of the following possibilities:

- (a)  $X_i \sim P(\lambda_i)$ .
- (b)  $X_i \sim B(n_i, p)$ .
- (c)  $X_i \sim G(p)$ .

Let  $Y = \sum_{i=1}^m X_i$ . Prove that, correspondingly:

- (a)  $Y \sim P(\sum_{i=1}^m \lambda_i)$ .
- (b)  $Y \sim B(\sum_{i=1}^m n_i, p)$ .
- (c)  $Y \sim \overline{B}(m, p)$ .

**Solution:** Recall the multinomial theorem:

$$(x_1 + x_2 + \dots + x_m)^n = \sum_{k_1+k_2+\dots+k_m=n; k_1, k_2, \dots, k_m \geq 0} \binom{n}{k_1, k_2, \dots, k_m} \prod_{i=1}^m x_i^{k_i}$$

$$X_i \sim P(\lambda_i) \implies P(X_i = k) = \frac{\lambda_i^k e^{-\lambda_i}}{k!}$$

$$P(Y = k) = P(X_1 + X_2 + \dots + X_m = k)$$

Since  $X_i$ 's are independent we get:

$$\begin{aligned} P(Y = k) &= P(X_1 + X_2 + \dots + X_m = k) = \sum_{k_1+\dots+k_m=k} P(X_1 = k_1)P(X_2 = k_2) \cdot \dots \cdot P(X_m = k_m) = \\ &= \sum_{k_1+\dots+k_m=k} \frac{\lambda_1^{k_1} e^{-\lambda_1}}{k_1!} \cdot \dots \cdot \frac{\lambda_m^{k_m} e^{-\lambda_m}}{k_m!} = e^{-(\lambda_1+\lambda_2+\dots+\lambda_m)} \cdot \left( \sum_{k_1+\dots+k_m=k} \frac{\lambda_1^{k_1}}{k_1!} \cdot \dots \cdot \frac{\lambda_m^{k_m}}{k_m!} \right) \cdot \frac{k!}{k!} = \\ &= \frac{e^{-(\lambda_1+\lambda_2+\dots+\lambda_m)}}{k!} \cdot \left( \sum_{k_1+\dots+k_m=k} \frac{k!}{k_1! \cdot k_2! \cdot \dots \cdot k_m!} \cdot (\lambda_1^{k_1} \cdot \lambda_2^{k_2} \cdot \dots \cdot \lambda_m^{k_m}) \right) = \\ &= \frac{e^{-(\lambda_1+\lambda_2+\dots+\lambda_m)}}{k!} \cdot \left( \sum_{k_1+\dots+k_m=k} \binom{k}{k_1, k_2, \dots, k_m} \cdot \prod_{i=1}^m \lambda_i^{k_i} \right) = \frac{e^{-(\lambda_1+\lambda_2+\dots+\lambda_m)}}{k!} \cdot (\lambda_1 + \lambda_2 + \dots + \lambda_m)^k = \\ &= \frac{e^{-(\sum_{i=1}^m \lambda_i)} \cdot (\sum_{i=1}^m \lambda_i)^k}{k!} \implies Y \sim P\left(\sum_{i=1}^m \lambda_i\right) \blacksquare \end{aligned}$$

#### 3.2 Q 4:

4. A die is rolled twice. Let  $X_i$  denote the outcome of the  $i$ th roll, and put  $S = X_1 + X_2$  and  $D = |X_1 - X_2|$ .

- (a) Show that  $E(SD) = E(S)E(D)$ .
- (b) Are  $S$  and  $D$  independent?

**Solution:**

(a)

$$E(S) = E(X_1 + X_2) = E(X_1) + E(X_2) = 3.5 + 3.5 = 7$$

$$E(D) = E(|X_1 - X_2|) = \sum_{i=0}^5 P(|X_1 - X_2| = i) \cdot i = 0 \cdot \frac{6}{36} + 1 \cdot \frac{10}{36} + 2 \cdot \frac{8}{36} + 3 \cdot \frac{6}{36} + 4 \cdot \frac{4}{36} + 5 \cdot \frac{2}{36} = \frac{35}{18}$$

$$E(SD) = E((X_1 + X_2)|X_1 - X_2|) = E(X_1 \cdot |X_1 - X_2|) + E(X_2 \cdot |X_1 - X_2|) = 2 \cdot E(X_1 \cdot |X_1 - X_2|)$$

The probability function  $P(X_1, |X_1 - X_2|)$  is given by the following table:

$X_1 \backslash  X_1 - X_2 $	0	1	2	3	4	5
1	1/36	1/36	1/36	1/36	1/36	1/36
2	1/36	1/18	1/36	1/36	1/36	0
3	1/36	1/18	1/18	1/36	0	0
4	1/36	1/18	1/18	1/36	0	0
5	1/36	1/18	1/36	1/36	1/36	0
6	1/36	1/36	1/36	1/36	1/36	1/36

Therefore:

$$E(SD) = 2 \cdot \sum_{i=1}^6 \sum_{j=0}^5 i \cdot j \cdot P(X_1 = i, |X_1 - X_2| = j) = \frac{245}{18} = 7 \cdot \frac{35}{18} = E(S) \cdot E(D) \blacksquare$$

(b)

$$P(X_1 + X_2 = 7, |X_1 - X_2| = 0) = 0 \neq \frac{6}{36} \cdot \frac{6}{36} = P(X_1 + X_2 = 7) \cdot P(|X_1 - X_2| = 0)$$

Therefore  $S$  and  $D$  are not independent.

### 3.3 Q 7:

7. A tetrahedron, with the numbers 1, 2, 3, and 4 marked on its faces, is tossed twice. Let  $X$  be the smaller of the two outcomes and  $Y$  the larger. Find  $\rho(X, Y)$ .

**Solution:**

$$1. P(X = k) = \frac{9-2k}{16}$$

$$5. P(Y = k) = \frac{2k-1}{16}$$

$$2. E(X) = \sum_{k=1}^4 k \cdot \frac{9-2k}{16} = \frac{15}{8}$$

$$6. E(Y) = \sum_{k=1}^4 k \cdot \frac{2k-1}{16} = \frac{25}{8}$$

$$3. E(X^2) = \sum_{k=1}^4 k^2 \cdot \frac{9-2k}{16} = \frac{35}{8}$$

$$7. E(Y^2) = \sum_{k=1}^4 k^2 \cdot \frac{2k-1}{16} = \frac{85}{8}$$

$$4. V(X) = \frac{35}{8} - \left(\frac{15}{8}\right)^2 = \frac{55}{64}$$

$$8. V(Y) = \frac{85}{8} - \left(\frac{25}{8}\right)^2 = \frac{55}{64}$$

The probability function  $P(X, Y)$  is given by the following table:

$X \backslash Y$	1	2	3	4
1	1/16	1/8	1/8	1/8
2	0	1/16	1/8	1/8
3	0	0	1/16	1/8
4	0	0	0	1/16

$$\begin{aligned}
E(XY) &= P(1,1) + 2P(1,2) + 3P(1,3) + 4\left(P(1,4) + P(2,2)\right) + \\
&\quad + 6P(2,3) + 8P(2,4) + 9P(3,3) + 12P(3,4) + 16P(4,4) = \\
&= \frac{1}{16} + \frac{2}{8} + \frac{3}{8} + \frac{4}{8} + \frac{4}{16} + \frac{6}{8} + \frac{8}{8} + \frac{9}{16} + \frac{12}{8} + \frac{16}{16} = \frac{25}{4}
\end{aligned}$$

In total we get:

$$\rho(X, Y) = \frac{\text{cov}(X, Y)}{\sqrt{(V(X)V(Y))}} = \frac{E(XY) - E(X)E(Y)}{\sqrt{(V(X)V(Y))}} = \frac{\frac{25}{4} - \frac{15}{8} \cdot \frac{25}{8}}{\sqrt{\frac{55}{64} \cdot \frac{55}{64}}} = \frac{\frac{25}{64}}{\frac{55}{64}} = \boxed{\frac{5}{11}}$$