

Distribution function of the total stopping time of the Collatz (3x+1) problem

Tomohiro Nishiyama
Email: htam0ybboh@gmail.com

Let $a := \ln \frac{3}{2}$ and $b := -\ln 2$. We approximate the total stopping time distribution of the Collatz problem by the Brownian motion with drift and a variance parameter σ . We define scaled parameters by

$$v := \frac{a+b}{2\sigma}, \quad s := \frac{a-b}{2\sigma},$$

and let

$$\alpha := \frac{9}{2}s^2 - (3s-v)\sigma, \quad \beta := 2s^2.$$

The total stopping time distribution of the Brownian motion for numbers $n \in [n_s, n_e]$ is given by

$$C(T; n_s, n_e) = \phi(T; n_e) - \phi(T; n_s), \quad (1)$$

$$\phi(T; n) := \sqrt{2}s\sigma \exp(g(x, T)) \left[\left(-\frac{s}{\sqrt{\beta}} - \frac{3s-v}{2\sqrt{\alpha}} \right) \operatorname{erfcx}(X) + \left(\frac{s}{\sqrt{\beta}} - \frac{3s-v}{2\sqrt{\alpha}} \right) \operatorname{erfcx}(Y) \right], \quad (2)$$

where

$$x := \frac{1}{\sigma} \ln n,$$

$$t := \frac{x + 2sT}{3s - v},$$

$$X := T\sqrt{\frac{\beta}{t}} + \sqrt{\alpha t}, \quad Y := T\sqrt{\frac{\beta}{t}} - \sqrt{\alpha t},$$

$$g(x, T) := (6s^2 - 2s\sigma)T - \alpha t - \frac{\beta}{t}T^2,$$

and $\operatorname{erfcx}(x)$ is the scaled complementary error function defined as $\operatorname{erfcx}(x) := \exp(x^2)\operatorname{erfc}(x)$. We can estimate the max of the total stopping time T_{\max} for numbers $n \in [n_s, n_e]$ by solving the following equation.

$$C(T_{\max}; n_s, n_e) \sim \phi(T_{\max}; n_e) = 1. \quad (3)$$

In our implementation, we put $\sigma = \frac{a-b}{2}$ and we use $\log \phi$ instead of ϕ to avoid overflow.