1

Assignment-2 **CHAPETR-15**

Matrices and Determinants

EE24BTECH11048-NITHIN.K

I. Section-B

1) Let $A = \begin{pmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{pmatrix}$. The only correct state-

ment about the matrix A is

[2004]

- a) $A^2 = I$
- b) A = (-1)I, where I is a unit matrix
- c) A^{-1} does not exist
- d) A is a zero matrix
- 2) Let $A = \begin{pmatrix} 1 & -1 & 1 \\ 2 & 1 & -3 \\ 1 & 1 & 1 \end{pmatrix}$, and $10B = \begin{pmatrix} 4 & 2 & 2 \\ -5 & 0 & \alpha \\ 1 & -2 & 3 \end{pmatrix}$. If B is the inversion of matrix A, then α is
 - a) 5

[2004]

- b) -1
- c) 2
- d) -2
- 3) If $a_1, a_2, a_3, \ldots, a_n, \ldots$ are in G.P, then the value of the determinant [2004] $| log a_n \quad log a_{n+1} \quad log a_{n+2} |$ $\begin{vmatrix} log a_{n+3} & log a_{n+4} & log a_{n+5} \\ log a_{n+6} & log a_{n+7} & log a_{n+8} \end{vmatrix}, \text{ is }$
 - a) -2
 - b) 1
 - c) 2
 - d) 0
- 4) If $A^2 A + I = 0$, then the inverse of A is [2005]
 - a) A+I
 - b) A
 - c) A-I
 - d) I-A
- 5) The system of equations $\alpha x + y + z = \alpha - 1$

- $x + \alpha y + z = \alpha 1$ $x + y + \alpha z = \alpha - 1$ has infinite solutions, if α is [2005]
- a) -2
- b) either -2 or 1
- c) not -2
- d) 1
- 6) If $a^2 + b^2 + c^2 = -2$ and $f(x) = \begin{vmatrix} 1 + a^2x & (1 + b^2)x & (1 + c^2)x \\ (1 + a^2)x & 1 + b^2x & (1 + c^2)x \\ (1 + a^2)x & (1 + b^2)x & 1 + c^2x \end{vmatrix}$ then $f(x) \text{ is a polynomial of degree} \qquad [2005]$
 - a) 1
 - b) 0
 - c) 3
 - d) 2
- 7) If $a_1, a_2, a_3, \ldots, a_n, \ldots$ are in G.P,then the determinant [2005] $\Delta = \begin{vmatrix} loga_n & loga_{n+1} & loga_{n+2} \\ loga_{n+3} & loga_{n+4} & loga_{n+5} \\ loga_{n+6} & loga_{n+7} & loga_{n+8} \end{vmatrix}$ is equal to
 - a) 1
 - b) 0
 - c) 4
 - d) 2
- 8) If A and B are square matrices of size n x n such that $A^2 - B^2 = (A - B)(A + B)$, then which of the following will be always true? [2006]
 - a) A = B
 - b) AB = BA
 - c) either of A or B is zero matrix
 - d) either of A or B is identity matrix

- 9) Let $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$ and $B = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$, $a,b \in N$. Then [2006]
 - a) there cannot exist any B such that AB = BA
 - b) there exist more than one but finite number of B's such that AB = BA
 - c) there exists exactly one B such that AB = BA
 - d) there exist infinitely many B's such that AB = BA
- 10) If D= $\begin{vmatrix} 1 & 1 & 1 \\ 1 & 1+x & 1 \\ 1 & 1 & 1+y \end{vmatrix}$ for $x \neq 0$, $y \neq 0$, then D is [2007]
 - a) divisible by x but not y
 - b) divisible by y but not x
 - c) divisible neither by x nor y
 - d) divisible by both x and y
- 11) Let A= $\begin{vmatrix} 5 & 5\alpha & \alpha \\ 0 & \alpha & 5\alpha \\ 0 & 0 & 5 \end{vmatrix}$. If $|A^2| = 25$, then $|\alpha|$ equals [2007]
 - a) $\frac{1}{5}$
 - b) 5
 - c) 5^2
 - d) 1
- 12) Let A be a 2x2 matrix with real entries. Let I be the 2x2 identity matrix. Denote by tr(A), the sum of diagonal entries of A. Assume that $A^2 = I$. [2008] Statement-1: If $A \neq I$ and $A \neq -I$, then det(A)=-1 Statement-2: If $A \neq I$ and $A \neq -I$, then $tr(A) \neq 0$.
 - a) Statement-1 is false, Statement-2 is true
 - b) Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1
 - c) Statement-1 is true, Statement-2 is true; Statement-2 is not a correct explanation for Statement-1
 - d) Statement-1 is true, Statement-2 is false
- 13) Let a,b,c be any real numbers. Suppose that there are real numbers x,y,z not all zero such that x = cy + bz, y = az + cx, and z = bx + ay. Then $a^2 + b^2 + c^2 + 2abc$ is equal to [2008]

- a) 2
- b) -1
- c) 0
- d) 1
- 14) Let A be a square matrix all of whose entries are integers. Then which of the following is true? [2008]
 - a) If $det(A) \neq \pm 1$, then A^{-1} exists but all its entries are not necessarily integers
 - b) If $det(A) \neq \pm 1$, then A^{-1} exists and all its entries are non integers
 - c) If $det(A) = \pm 1$, then A^{-1} exists but all its entries are integers
 - d) If $det(A) = \pm 1$, then A^{-1} need not exists
- 15) Let A be a 2x2 matrix Statement-1:adj(adjA) = AStatement-2:|adjA| = |A| [2009]
 - a) Statement-1 is true, Statement-2 is true; Statement-2 is not a correct explanation for Statement-1
 - b) Statement-1 is true, Statement-2 is false
 - c) Statement-1 is false, Statement-2 is true
 - d) Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1