1

Quiz 9

S Nithish

Abstract—This document contains the solution of the question from NCERT 9th standard chapter 10 exercise 10.4 problem 4

1 Exercise 10.4

 If a line intersects two concentric circles with centre O at A,B,C and D, prove that AB=CD. Let the inner circle be of radius 3 and the outer circle be of radius 5. Then the equation of the two circles are,

$$||\mathbf{x}|| = 9 \tag{1.0.1}$$

$$||\mathbf{x}|| = 25 \tag{1.0.2}$$

Let the equation of the line be,

$$\mathbf{x} = \begin{pmatrix} 0 \\ -2 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 1 \end{pmatrix} \tag{1.0.3}$$

$$\mathbf{x} = \begin{pmatrix} \lambda \\ \lambda - 2 \end{pmatrix} \tag{1.0.4}$$

The points of intersection of the line and the inner circle are,

$$||\mathbf{x}||^2 = \lambda^2 + (\lambda - 2)^2$$
 (1.0.5)

$$||\mathbf{x}||^2 = 9 \tag{1.0.6}$$

$$\lambda^2 + (\lambda - 2)^2 = 9 \tag{1.0.7}$$

$$2\lambda^2 - 4\lambda + 4 = 9 \tag{1.0.8}$$

$$\lambda^2 - 2\lambda - \frac{5}{2} = 0 \tag{1.0.9}$$

$$\implies \lambda_c = 1 + \sqrt{\frac{7}{2}}, \ \lambda_b = 1 - \sqrt{\frac{7}{2}}$$
 (1.0.10)

The points of intersection of the line and the outter circle are,

$$||\mathbf{x}||^{2} = \lambda^{2} + (\lambda - 2)^{2}$$

$$(1.0.11)$$

$$||\mathbf{x}||^{2} = 25 \qquad (1.0.12)$$

$$\lambda^{2} + (\lambda - 2)^{2} = 25 \qquad (1.0.13)$$

$$2\lambda^{2} - 4\lambda + 4 = 25 \qquad (1.0.14)$$

$$\lambda^{2} - 2\lambda - \frac{21}{2} = 0 \qquad (1.0.15)$$

$$\implies \lambda_{d} = 1 + \sqrt{\frac{23}{2}}, \ \lambda_{a} = 1 - \sqrt{\frac{23}{2}}$$

$$\lambda_{a} = 1 - \sqrt{\frac{23}{2}} \implies \mathbf{x_{a}} = \begin{pmatrix} 1 - \sqrt{\frac{23}{2}} \\ -1 - \sqrt{\frac{23}{2}} \end{pmatrix}$$
(1.0.17)
$$\lambda_{b} = 1 - \sqrt{\frac{7}{2}} \implies \mathbf{x_{b}} = \begin{pmatrix} 1 - \sqrt{\frac{7}{2}} \\ -1 - \sqrt{\frac{7}{2}} \end{pmatrix}$$
(1.0.18)
$$\lambda_{c} = 1 + \sqrt{\frac{7}{2}} \implies \mathbf{x_{c}} = \begin{pmatrix} 1 + \sqrt{\frac{7}{2}} \\ -1 + \sqrt{\frac{7}{2}} \end{pmatrix}$$
(1.0.19)
$$\lambda_{d} = 1 + \sqrt{\frac{23}{2}} \implies \mathbf{x_{d}} = \begin{pmatrix} 1 + \sqrt{\frac{23}{2}} \\ -1 + \sqrt{\frac{23}{2}} \end{pmatrix}$$

The distance between points A and B is,

$$AB = \|\mathbf{x_a} - \mathbf{x_b}\| = \left\| \left(\frac{\sqrt{7} - \sqrt{23}}{\sqrt{2}} \frac{\sqrt{7} - \sqrt{23}}{\sqrt{2}} \right) \right\| = \sqrt{23} - \sqrt{7}$$

$$(1.0.21)$$

$$CD = \|\mathbf{x_c} - \mathbf{x_d}\| = \left\| \left(\frac{\sqrt{7} - \sqrt{23}}{\sqrt{2}} \frac{\sqrt{7} - \sqrt{23}}{\sqrt{2}} \right) \right\| = \sqrt{23} - \sqrt{7}$$

$$(1.0.22)$$

$$\implies AB = CD \qquad (1.0.23)$$

(1.0.20)