OurMocks

2027 CYCLE

MOCK · PRACTICE · CRACK

CUET UG

Biology

# Organisms & Populations

Official Syllabus Unit X · Class 12 · Comprehensive NCERT Theory, Diagrams & 15 Solved MCQs

INSIDE
Concept Notes

INSIDE
Solved Examples

INSIDE
Practice MCQs + Solutions

Aligned to the latest official syllabus · Concept-first explanations

www.ourmocks.com

# Syllabus Alignment & Exam Blueprint

| Sub-Topic | Official Syllabus Core Coverage (NCERT Class 12 Unit X) | Exam Focus & Question Types |
|---|---|---|
| **Organism & Abiotic Environment** | Ecological hierarchy (organism, population, community, biome); major abiotic factors: temperature (stenothermal vs eurythermal), water (stenohaline vs euryhaline), light, and soil (edaphic factors); organismic responses: regulate (homeostasis), conform (>99% animals), migrate (Keoladeo Ghana National Park, Bharatpur), and suspend (diapause in zooplankton, hibernation in bears, aestivation in snails/fish). | Definitions of eurythermal/ stenothermal; physiological responses to high altitude (erythropoietin, RBC count); Keoladeo migratory birds. |
| **Adaptations to Habitats** | Morphological, physiological, and behavioral adaptations; desert animals (Kangaroo rat internal fat oxidation, hyper-concentrated urine); desert plants (CAM photosynthetic pathway, sunken stomata, thick cuticle, *Opuntia* phylloclade spines); Allen's rule (shorter extremities in cold climates); blubber in aquatic mammals (seals); behavioral thermal regulation in desert lizards. | Calculation of water balance; identification of Allen's rule; distinguishing physiological adaptations from behavioral thermoregulation. |
| **Population Attributes & Growth Models** | Natality, mortality, sex ratio, and age pyramids (expanding, stable, declining); population density estimation (direct biomass/percent cover vs indirect pugmarks/fecal pellets); vital rates equation $N_{t+1} = N_t + [(B+I) - (D+E)]$; exponential growth ($d\frac{N}{d}t = rN$, $N_t = N_0 e^{rt}$, J-curve, intrinsic rate $r$); logistic growth (Verhulst-Pearl, $d\frac{N}{d}t = rN\left(\frac{K-N}{K}\right)$, carrying capacity $K$, S-curve). | Per capita birth/death rate arithmetic; interpretation of age pyramid shapes; identifying peak growth rate at $\frac{K}{2}$; exponential vs logistic comparison. |
| **Interspecific Population Interactions** | Master matrix of species interactions: Mutualism (+/+), Competition (-/-), Predation (+/-), Parasitism (+/-), Commensalism (+/0), and Amensalism (-/0); Gause's competitive exclusion principle; Connell's barnacles (*Balanus* vs *Chthamalus*); MacArthur's resource partitioning (5 warbler species); keystone predator *Pisaster*; Monarch butterfly cardiac glycoside; brood parasitism (cuckoo & crow); *Ophrys* sexual deceit. | Sign matching (+/+, +/-, -/-, +/0, -/0); case study identification: Abingdon tortoise extinction, prickly pear cactus, cattle egret, *Penicillium* antibiosis. |

### EXAM PATTERN

**CUET UG Scoring Blueprint:** Organisms and Populations accounts for **3 to 4 questions (15 to 20 marks)** in the CUET UG domain paper. NTA frequently tests **exact mathematical formulations** of growth models ($d\frac{N}{d}t = rN\left(\frac{K-N}{K}\right)$ and $N_t = N_0 e^{rt}$), **per capita vital rate calculations**, **age pyramid morphology**, and **case study matching across the six interspecific interactions** (especially *Ophrys*, brood parasitism, and amensalism).

# 1. Ecology, Abiotic Factors & Organismic Adaptations

Ecology is the fundamental biological discipline that investigates the reciprocal relationships and dynamic interactions among living organisms and between organisms and their abiotic (physical and chemical) environment. The ecological hierarchy is organized across four foundational levels of increasing biological complexity: **Organisms → Populations → Biological Communities → Biomes**. At the organismic level, ecology is fundamentally **physiological ecology**, examining how individual species are morphologically, anatomically, biochemically, and behaviorally adapted to survive, thrive, and reproduce under shifting environmental regimes.

## Habitat vs Ecological Niche Architecture

In ecological theory, a fundamental distinction exists between an organism's physical dwelling and its functional status within the biotic community:

- **Habitat (The Ecological Address):** The specific physical geographical locality where an organism lives, encompassing its ambient physical microclimate, topography, and associated biotic community.
- **Ecological Niche (The Ecological Profession):** First conceptualized by Joseph Grinnell (1917) and expanded by Charles Elton and G. Evelyn Hutchinson, the ecological niche encompasses the **entire functional role** played by an organism in its ecosystem: the exact physical and chemical conditions it can tolerate, the specific resources it utilizes, and its interactions with other species.
- **Fundamental vs Realized Niche:**
  - **Fundamental Niche:** The total theoretical range of environmental conditions under which a species can survive and reproduce in the complete absence of interspecific competitors and predators.
  - **Realized Niche:** The actual, restricted portion of the fundamental niche that the species occupies in nature in the presence of competing species and natural enemies.

## Major Planetary Biomes & Environmental Gradients

On a global macro-ecological scale, the continuous rotation of planet Earth around the Sun combined with the tilt of its rotational axis ($23.5^\circ$) causes annual variations in the intensity and duration of solar radiation. These annual temperature cycles, coupled with annual precipitation patterns (rain and snow), generate Earth's major terrestrial **Biomes**:

1. **Tropical Rain Forests:** Characterized by warm temperatures ($20^\circ$ C $-$ $25^\circ$ C) and high annual rainfall ($150 - 400$ cm), fostering multi-layered evergreen canopies and maximum planetary biodiversity.
2. **Deciduous Temperate Forests:** Moderate seasonal temperatures and moderate precipitation ($100 -$ $200$ cm), where broad-leaved trees shed foliage in autumn.
3. **Deserts:** Scorching temperatures and extremely low precipitation ($< 50$ cm), supporting specialized xerophytic flora and burrowing fauna.
4. **Coniferous (Taiga / Boreal) Forests:** Sub-polar forests with cold temperatures ($-1^\circ$ C $-$ $15^\circ$ C) and annual precipitation of $100 - 250$ cm, dominated by evergreen gymnosperms (pines, spruces, firs).
5. **Arctic and Alpine Tundra:** Treeless polar landscapes with freezing temperatures ($-10^\circ$ C $-$ $0^\circ$ C), low precipitation ($< 100$ cm), and permanently frozen subsoil (permafrost).

| Major Biome | Mean Annual Temp ($^\circ$ C) | Mean Annual Precipitation (cm) | Dominant Floral & Faunal Characteristics |
|---|---|---|---|
| Tropical Rain Forest | $20 - 25^\circ$ C | $150 - 400$ cm | Multi-tiered dense canopy; lianas, epiphytic orchids; maximum planetary species richness. |
| Deciduous Forest | $10 - 20^\circ$ C | $90 - 160$ cm | Broad-leaved trees (oak, maple, sal, teak) shedding leaves seasonally in winter/autumn. |
| Desert (Hot & Cold) | $5 - 25^\circ$ C | $< 50$ cm | Xerophytes, succulents, CAM plants (*Opuntia*); nocturnal rodents (kangaroo rats). |
| Coniferous Forest | $-1 - 15^\circ$ C | $100 - 250$ cm | Needle-leaved gymnosperms (pines, firs); acidic podzol soil; moose, wolves, lynx. |
| Arctic Tundra | $-10 - 0^\circ$ C | $< 100$ cm | Permafrost subsoil; dwarf willows, lichens (reindeer moss), mosses; arctic fox, caribou. |

## The Four Major Abiotic Environmental Factors

1. **Temperature (The Primary Ecological Driver):**
   - Temperature is the most ecologically decisive physical environmental factor. It exhibits a global latitudinal gradient, progressively decreasing from the warm equator toward the frigid poles, and an altitudinal gradient, decreasing from plains to mountain summits.
   - Temperature ranges from subzero levels ($< -50^{\circ}$ C) in polar blizzards to $> 50^{\circ}$ C in tropical deserts. Extreme habitats—such as deep-sea hydrothermal vents and hot geothermal springs—harbor specialized hyperthermophiles thriving at $> 100^{\circ}$ C.
   - **Thermal Tolerance Spectra:**
     - **Eurythermal Organisms:** Organisms capable of tolerating, metabolizing, and reproducing across a **wide range** of temperatures (e.g., most mammals, birds, and humans).
     - **Stenothermal Organisms:** Organisms restricted to a **narrow, specific temperature band** (e.g., polar bears, tropical corals, deep-sea fishes).
   - Temperature governs the geographical distribution of organisms primarily by dictating the kinetics of metabolic enzymes, cell membrane fluidity, and basal metabolic rates.
   - **Thermal Stratification in Aquatic Systems:** In deep freshwater lakes during summer, water stratifies into three distinct thermal zones: the warm, circulating surface **Epilimnion**, the middle transition layer characterized by a steep temperature gradient termed the **Thermocline (Metalimnion)**, and the cold, dense, stagnant bottom layer termed the **Hypolimnion**. In temperate lakes, seasonal density changes trigger spring and autumn **turnovers**, circulating oxygen and nutrients between benthic and surface waters.
2. **Water (The Solvent of Life):**
   - Life on Earth originated in primeval seas and remains biochemically unsustainable without water. Plant primary productivity and geographical distribution are decisively water-regulated.
   - **Salinity Dynamics (Parts Per Thousand / ppt):**
     - Inland freshwaters: $< 5$ ppt.
     - Open marine oceans: $30 - 35$ ppt.
     - Hypersaline coastal lagoons: $> 100$ ppt.
   - **Euryhaline** organisms tolerate wide fluctuations in salinity (e.g., salmon and euryhaline crabs), whereas **Stenohaline** organisms are restricted to narrow salinity ranges.
   - **Osmoregulatory Challenges:** Freshwater teleosts inhabit a hypotonic medium and face constant water influx and salt loss; they excrete large volumes of dilute urine and actively absorb salts via gill chloride cells. Marine teleosts inhabit a hypertonic medium and face continuous osmotic dehydration; they continuously drink seawater and actively excrete excess salts across their gills while producing meager concentrated urine.
3. **Light (Photoperiodism & Vertical Zonation):**
   - Solar radiation drives autotrophic photosynthetic energy capture.
   - Understory herbs and shrubs in dense tropical rainforests are sciophytes, adapted to photosynthesize at low light intensities under the shadow of upper canopy trees.
   - **Photoperiodism:** Many plants rely on seasonal changes in photoperiod (day length) as an environmental cue to induce flowering. Animals synchronize their foraging excursions, long-distance seasonal migrations, and reproductive cycles using photoperiodic signals.
   - **Spectral Distribution in Oceans:** Sunlight penetrating ocean waters is filtered selectively. Long wavelengths (red and yellow) are absorbed within the upper 20 meters, whereas short blue-green wavelengths penetrate deepest. Consequently, **Red Algae (Rhodophyceae)** colonize the deepest photic zones because their accessory pigment **phycoerythrin** efficiently absorbs blue-green light.
4. **Soil (Edaphic Factors):**
   - Soil characteristics vary according to climate, parent rock weathering, transport history (alluvial, colluvial, aeolian), and soil age.
   - Physical edaphic properties—grain size distribution, pore volume, mineral composition, pH, and aggregation—dictate soil permeability and water-holding capacity.
   - A complete soil profile comprises distinctive horizons: the organic humus-rich **O horizon**, the biologically active mineral topsoil **A horizon**, the illuviated subsoil **B horizon**, the partially weathered parent regolith **C horizon**, and solid unweathered bedrock (**R horizon**).

- In aquatic ecosystems, bottom sediment characteristics determine the benthic fauna that can anchor and forage on the lake or ocean floor.

## Responses to Abiotic Environmental Fluctuations (Homeostasis)

Organisms maintain internal constancy (**homeostasis**) through four major evolutionary strategies:

1. **Regulate (Regulators):**
   - Organisms maintain internal homeostasis through physiological and behavioral adjustments, ensuring constant body temperature (thermoregulation) and constant blood osmotic concentration (osmoregulation).
   - All birds and mammals, and very few lower vertebrates and invertebrates, are regulators. Humans maintain a constant body core temperature of $37^{\circ}$ C. When ambient temperatures exceed $37^{\circ}$ C, sweating and evaporative cooling dissipate excess heat; in freezing conditions, shivering (involuntary somatic muscle contraction) generates metabolic heat.
   - **Energetics of Thermoregulation (Why Small Animals are Rare in Polar Regions):**
     - Heat loss or heat gain is a physical function of **surface area**.
     - Small animals (such as shrews and hummingbirds) have a **very high surface-area-to-volume ratio**. Consequently, they lose body heat extraordinarily fast in cold climates and must consume vast caloric food supplies to sustain baseline body heat, making small endotherms evolutionarily unviable in polar zones.
2. **Conform (Conformers):**
   - An overwhelming majority (**99% of all animals and virtually all plants**) cannot maintain constant internal osmolarity or core temperature. Their internal parameters change passively with ambient fluctuations.
3. **Migrate:**
   - Organisms temporarily evacuate a stressful geographical habitat to relocate to a hospitable sanctuary, returning once the stressful period terminates.
   - The **Keoladeo National Park in Bharatpur (Rajasthan)** hosts thousands of migratory birds arriving from Siberia and extreme northern regions every winter.
4. **Suspend (Dormancy Strategies):**
   - Organisms escape environmental stress in **time** rather than space:
     - Bacteria, fungi, and lower plants synthesize thick-walled resistant **spores**.
     - Higher plants enter seed and bud **dormancy**.
     - **Hibernation:** Winter sleep undertaken by northern bears to escape freezing food-scarce winters.
     - **Aestivation:** Summer sleep undertaken by terrestrial snails and lungfishes to avoid scorching summer heat and desiccation.
     - **Diapause:** A developmental stage of suspended metabolic activity exhibited by many freshwater zooplankton species under adverse lake conditions.

## Morphological, Physiological & Behavioral Adaptations

- **Desert Plants (Xerophytic Adaptations):**
  - Thick, waxy leaf cuticles minimize transpirational water loss.
  - Sunken stomata arranged in deep crypts reduce convective airflow.
  - **CAM Photosynthesis (Crassulacean Acid Metabolism):** Stomata remain tightly closed during hot daytime hours and open only at night to take up $CO_2$, minimizing evaporative water loss.
  - Desert shrubs like *Opuntia* (prickly pear) reduce leaves to defensive spines; photosynthesis is taken over by flattened green photosynthetic stems (**phylloclades**).
- **Desert Animals (Kangaroo Rat):**
  - The kangaroo rat of North American deserts survives without drinking liquid water: it derives 90% of its water from the **internal oxidation of dietary fats** (metabolic water) and concentrates its urine so intensely that minimal water is excreted.
- **Zoological Ecological Rules:**
  - **Allen's Rule:** Mammals inhabiting colder geographical climates possess **shorter ears and shorter limbs (extremities)** to minimize surface area and heat dissipation.

- **Bergmann's Rule:** Warm-blooded vertebrates inhabiting colder geographical climates tend to have larger total body sizes than related species living in warmer climates (reducing surface-area-to-volume ratio).
- **Gloger's Rule:** Animals inhabiting warm and humid regions develop darker melanin pigmentation than individuals of the same species living in cold and dry climates.

- **Marine Mammals (Blubber):** Polar seals, sea lions, and whales possess a thick subcutaneous insulating layer of fat (**blubber**) underneath their epidermis that reduces body heat dissipation in freezing oceans.
- **High-Altitude Sickness & Acclimatization:**
  - At elevations $> 3,500$ meters (e.g., Rohtang Pass, Leh), low atmospheric barometric pressure reduces the partial pressure of oxygen ($pO_2$), triggering altitude sickness (nausea, fatigue, palpitations).
  - The human body acclimatizes through three simultaneous physiological adjustments:
    1. Kidneys secrete **erythropoietin**, accelerating **red blood cell (RBC) synthesis**.
    2. The respiratory control center increases baseline **breathing rate**.
    3. The body **decreases the binding affinity of hemoglobin for oxygen**, allowing rapid and complete unloading of oxygen to tissue capillaries.

# 2. Population Attributes & Growth Dynamics

In nature, individual organisms rarely exist in total isolation. They congregate in geographical groups of conspecific individuals that interbreed, share or compete for similar resources, and occupy defined spatial territories—termed a **Population**. While natural selection operates on individuals, biological evolution occurs at the **population level**.

## Individual vs Population Attributes

| Attribute Category | Individual Organism Level | Ecological Population Level |
|---|---|---|
| Birth & Death | An individual is **born** or **dies** (singular absolute events). | A population exhibits **birth rates (natality)** and **death rates (mortality)** expressed as per capita metrics. |
| Sex Distribution | An individual is either **male** or **female**. | A population exhibits a **sex ratio** (e.g., 60% females, 40% males). |
| Age Structure | An individual has a specific chronological age. | A population is composed of distinct age classes, represented visually via **Age Pyramids**. |

## Age Pyramids of Human Populations

Plotting the percentage individuals across three ecological age categories—**Pre-reproductive** (0–14 years), **Reproductive** (15–44 years), and **Post-reproductive** (45+ years)—yields three distinctive population profiles:

1. **Expanding Population (Triangular Base):** High proportion of pre-reproductive individuals forming a broad base; rapid positive population growth (typical of developing nations).
2. **Stable Population (Bell-Shaped):** Pre-reproductive and reproductive cohorts are roughly equal in size; zero or near-zero population growth.
3. **Declining Population (Urn-Shaped with Narrow Base):** Pre-reproductive cohort is significantly smaller than the reproductive class; negative population growth (birth rates fall below death rates).





Figure 1: Population growth kinetics — comparative mathematical trajectories of exponential J-shaped growth (unlimited resources) and Verhulst-Pearl logistic S-shaped sigmoid growth (carrying capacity K).

## Population Density & Vital Demographic Processes

Population density ($N$) is the number of individuals of a species per unit area or volume. It fluctuates dynamically in response to four core demographic processes:

$$N_{t+1} = N_t + [(B + I) - (D + E)]$$

- **Natality ($B$):** Number of births in the population during a specified time interval.
- **Mortality ($D$):** Number of deaths in the population during the period.
- **Immigration ($I$):** Number of individuals of the same species that moved into the habitat from elsewhere.
- **Emigration ($E$):** Number of individuals that permanently departed the habitat.
- **Diagnostic Rule:** Under ordinary conditions, births and deaths are the most decisive factors; however, when a new habitat is colonized, **immigration ($I$)** contributes more decisively to population growth than birth rates.

## Mathematical Population Growth Models

1. **Exponential Growth (J-Shaped Trajectory):**
   - Occurs when habitat resources (food and space) are completely unlimited.
   - Rate of population change is given by the differential equation:
   $$\frac{dN}{dt} = (b - d)N = rN$$
   - The parameter $r$ is the **intrinsic rate of natural increase**, a key metric evaluating biotic potential. (Values of $r$: Norway rat = 0.015, flour beetle = 0.12, Indian human population in 1981 = 0.0205).
   - Integrated equation form:
   $$N_t = N_0 e^{rt}$$
   - Produces an explosive, geometric **J-shaped growth curve**.
2. **Logistic Growth (Verhulst-Pearl S-Shaped Sigmoid Curve):**
   - In nature, resources are strictly finite. Every habitat has an upper threshold of resource availability capable of sustaining a maximal population density, beyond which no further growth is possible—designated as the **Carrying Capacity ($K$)**.
   - Verhulst-Pearl Logistic Growth equation:
   $$\frac{dN}{dt} = rN\left(\frac{K - N}{K}\right)$$
   - **Inflection Point & Peak Growth:** The term $\frac{K-N}{K}$ represents environmental resistance. The absolute rate of population growth ($\frac{dN}{dt}$) reaches its maximum at **half the carrying capacity ($N = \frac{K}{2}$)**. As $N$ approaches $K$, growth slows to zero, establishing a stable plateau (asymptote).

## Life History Variations: r-Selected vs K-Selected Strategies

Populations evolve reproductive strategies that maximize their evolutionary Darwinian fitness ($r$) in the specific habitats they occupy:

| Life History Feature | r-Selected Strategists | K-Selected Strategists |
|---|---|---|
| Environmental Stability | Unstable, unpredictable, or ephemeral habitats. | Stable, predictable, crowded habitats near carrying capacity ($K$). |
| Body Size & Longevity | Small body size; short lifespan; rapid maturation. | Large body size; long lifespan; delayed sexual maturation. |
| Breeding Frequency | Semelparous (breed once in lifetime, e.g., Pacific salmon, bamboo). | Iteroparous (breed repeatedly over lifetime, e.g., birds, mammals). |
| Offspring Number & Size | Vast numbers of small-sized offspring (e.g., oysters, pelagic fishes). | Few, large-sized offspring with high energetic investment. |
| Parental Care | Minimal or zero parental care; high juvenile mortality. | Extensive parental care; high juvenile survival rates. |

# 3. Interspecific Population Interactions

In any biological community, no species can survive in isolation. Biological interactions between two distinct interacting species are termed **interspecific interactions**, categorized by beneficial (+), harmful (−), or neutral (0) outcomes:




Figure 2: Taxonomy of interspecific species interactions — ecological matrices of mutualism, competition, predation, parasitism, commensalism, and amensalism across terrestrial and marine ecosystems.

| Interaction Type | Species A | Species B | Ecological Definition & High-Yield Paradigm |
|---|---|---|---|
| Mutualism | + | + | Obligate or facultative cooperation where both interacting partners benefit (Lichens, Mycorrhizae, Fig & Fig Wasp). |
| Competition | − | − | Both species suffer fitness costs due to mutual exploitation of limiting resources (Balanus vs Chthamalus). |
| Predation | + | − | Predator catches, kills, and consumes prey; acts as conduit for energy transfer (Tiger & Deer, Pisaster starfish). |
| Parasitism | + | − | Parasite lives in or on host, deriving nutrients while reducing host survival and fitness (Lice, Ticks, Liver fluke). |

| Interaction Type | Species A | Species B | Ecological Definition & High-Yield Paradigm |
|---|---|---|---|
| Commensalism | + | 0 | One partner benefits while the second partner is completely unaffected (Orchid on Mango tree, Cattle Egret). |
| Amensalism | − | 0 | One species is inhibited or destroyed while the second partner remains neutral (*Penicillium* and Bacteria). |

## Predation & Its Ecological Significance

- **Energy Conduits:** Predators serve as conduits for energy transfer across successive trophic levels in food webs.
- **Controlling Prey Populations:** In the 1920s, the invasive **Prickly Pear Cactus (*Opuntia*)** was introduced into Australia, spreading uncontrollably over millions of hectares of rangeland. It was brought under control only after introducing its natural predator, a cactus-feeding moth (***Cactoblastis cactorum***).
- **Maintaining Community Biodiversity (Keystone Predation):**
  - In a celebrated field experiment conducted by ecologist Robert Paine in the rocky intertidal communities of the American Pacific coast, all predatory starfish of the genus ***Pisaster*** were removed.
  - Within a single year, **more than 10 species of intertidal invertebrates became locally extinct** because unchecked competition among dominant mussel prey excluded other species.
- **Prey Defenses:**
  - **Camouflage (Cryptic Coloration):** Insects and frogs blend into background vegetation to avoid detection.
  - **Chemical Toxins:** The **Monarch butterfly** accumulates a highly toxic **cardiac glycoside** during its caterpillar stage by feeding on poisonous milkweed, making adult butterflies intensely distasteful to predatory birds.
  - **Plant Defenses:** Morphological defenses (thorns of *Acacia*, spines of cactus); chemical repellents (cardiac glycosides in *Calotropis*, nicotine, caffeine, quinine, strychnine, and opium). Nearly 25% of all known insects are phytophagous (feeding on plant sap and tissues), driving an intense chemical arms race where plants synthesize complex secondary metabolites to deter herbivores.

## Interspecific Competition & Coexistence Mechanisms

- **Gause's Competitive Exclusion Principle:** Two closely related species competing for the exact same limiting resource cannot coexist indefinitely; the competitively inferior species will be eliminated. (Illustrated when the **Abingdon tortoise** in the Galapagos Islands became extinct within a decade after domestic goats were introduced, due to the goats' greater browsing efficiency).
- **Connell's Elegant Field Experiments:** On the rocky intertidal coasts of Scotland, the large, competitively dominant barnacle ***Balanus*** occupies the lower intertidal zone, competitively excluding the smaller barnacle ***Chthamalus*** from that zone.
- **Resource Partitioning (Avoiding Exclusion):** If two species compete for the same resource, they avoid exclusion by choosing different times for feeding or different foraging patterns. **Robert MacArthur** demonstrated that five closely related species of **Warblers** successfully coexisted on the same spruce tree by foraging in different micro-habitats and vertical zones of the canopy.

## Parasitism: Ecto- and Endoparasitism & Brood Parasitism

- **Parasitic Adaptations:** Morphological reductions including loss of unnecessary sense organs, development of adhesive organs or suckers, loss of digestive tracts, high reproductive capacity, and intricate multi-host developmental cycles. For instance, the human liver fluke (*Fasciola*) depends on two intermediate hosts: a freshwater snail and a fish.
- **Ectoparasites vs Endoparasites:** Ectoparasites feed on the external surface of host organisms (head lice on humans, ticks on dogs, marine copepods on fish, and the non-chlorophyllous angiosperm ***Cuscuta*** / dodder on hedge plants). Endoparasites inhabit internal organs (gut, liver, erythrocytes, lungs); their morphology is profoundly simplified to maximize nutrient absorption and gamete production.
- **Brood Parasitism:** A specialized behavioral parasitism in birds where the parasite lays eggs in the host's nest. The **Asian Koel (cuckoo)** lays eggs in the nest of the **crow**; the eggs have evolved to closely mimic host eggs in size, color, and speckling to prevent host recognition and nest eviction.

## Commensalism, Mutualism & Co-evolution

- **Commensalism ($\frac{+}{0}$):**
  - Epiphytic orchids growing perched on mango tree branches for physical support and sunlight.
  - Barnacles permanently attached to the dorsal skin of migratory baleen whales.
  - Cattle egrets foraging in pastures, feeding on insects flushed out by the movement of grazing cattle.
  - Sea anemone with venomous stinging tentacles sheltering the clownfish, whose mucus coating protects it from stinging.
- **Mutualism ($\frac{+}{+}$) & Obligate Symbioses:**
  - **Lichens:** Intimate mutualism between a heterotrophic fungal partner (**mycobiont**) providing moisture, minerals, and shelter, and a photosynthetic green algal or cyanobacterial partner (**phycobiont**) providing carbohydrates.
  - **Mycorrhizae:** Symbiotic mutualism between soil fungi (predominantly genus ***Glomus***) and the root systems of higher plants. The fungal hyphae absorb immobile phosphorus from soil and deliver it to the plant, receiving photosynthetic carbohydrates and shelter. *Pinus* seeds cannot germinate or establish seedlings without obligate mycorrhizal colonization!
  - **Plant-Pollinator Mutualisms:** The celebrated mutualism between the **Fig tree** and its obligate pollinator **Fig Wasp** (*Blastophaga*). The female wasp searches for an inflorescence (hypanthodium), lays eggs inside unfertilized ovaries (forming gall flowers), and simultaneously pollinates other female flowers. Emerging wingless males mate with females inside the syconium; newly hatched winged females collect pollen and exit to colonize new figs.
  - **Sexual Deceit in Orchid *Ophrys*:** The Mediterranean orchid ***Ophrys*** employs sexual deceit for pollination. One floral petal mimics the female solitary bee in physical shape, color markings, and chemical sex pheromone bouquet. The male bee attempts copulation (**pseudocopulation**), during which pollen masses (pollinia) are transferred to the bee's head and deposited on subsequent flowers. If the female bee evolves minor morphological or chemical variations, the orchid must co-evolve corresponding adjustments to maintain pollination fidelity.

## Evolutionary Arms Race: Host-Parasite & Plant-Herbivore Dynamics

In co-evolutionary relationships, an evolutionary innovation in one species acts as a potent selective pressure driving reciprocal adaptations in the interacting species:

- **Parasite-Host Evolutionary Trajectory:**
  - If a host species evolves novel cellular or physiological defense mechanisms against a parasite, the parasite must evolve counter-adaptations to neutralize or bypass those host defenses to avoid extinction.
  - **Extreme Morphological Specialization:** In response to selection pressure, endoparasites have undergone profound anatomical reduction: loss of unnecessary locomotory appendages, complete degeneration of digestive systems (absorbing predigested nutrients directly through the tegument), and explosive expansion of reproductive potential (producing millions of eggs per day).
  - **Complex Multi-Host Life Cycles:** Many endoparasites exploit complex life cycles involving one or two intermediate vectors to facilitate transmission between definitive hosts:
    - **Human Liver Fluke (*Clonorchis sinensis*):** Relies on two obligatory intermediate hosts—a freshwater snail (first intermediate host) and a cyprinid fish (second intermediate host)—before infecting the human hepatic bile ducts.
    - **Malarial Parasite (*Plasmodium*):** Alternates between sexual reproduction inside the gut of the female *Anopheles* mosquito vector and asexual schizogony inside human hepatocytes and erythrocytes.
- **Plant-Herbivore Chemical Warfare:**
  - Approximately 25% of all phytophagous insects feed exclusively on plant sap and vegetative tissues. Because plants cannot flee from herbivores, they have evolved elaborate secondary chemical defenses:
    - **Cardiac Glycosides:** The weed *Calotropis* produces highly poisonous cardiac glycosides in its milky latex; grazing cattle and goats completely avoid browsing on it.
    - **Commercial Secondary Metabolites:** Widely extracted compounds—such as **nicotine** (tobacco), **caffeine** (coffee/tea), **quinine** (cinchona bark), **strychnine** (*Strychnos*), and **morphine/opium** (poppy)—are synthesized by plants not for primary cellular metabolism, but as potent neurotoxins and chemical deterrents against herbivorous insects and mammals.

**CUET HIGH-YIELD TRAP: INTERSPECIFIC SIGN CONVENTIONS & NTA TRAPS**

NTA examination papers consistently probe sign conventions in interaction matrices:

1. **Predation ($\overset{+}{-}$) vs Parasitism ($\overset{+}{-}$):** Both interactions share identical mathematical sign designations ($\overset{+}{-}$). However, predators typically kill their prey rapidly and consume multiple prey individuals over their lifetime, whereas parasites are generally smaller than their host, derive nutrition without immediate lethality, and are closely associated with a single host individual.
2. **Commensalism ($\overset{+}{0}$) vs Amensalism ($\overset{-}{0}$):** Do not confuse the zero partner! In commensalism, one partner benefits while the other is neutral ($\overset{+}{0}$); in amensalism, one partner is inhibited/harmed while the second partner remains completely unaffected ($\overset{-}{0}$).

## Comprehensive Matrix of Interspecific Interactions

| Interaction Category | Species 1 ($\overset{\pm}{0}$) | Species 2 ($\overset{\pm}{0}$) | Classic NCERT Example / Paradigm |
|---|---|---|---|
| Mutualism | Benefited (+) | Benefited (+) | *Ophrys* orchid and pollinating bee (sexual deceit); Lichen (fungus + alga). |
| Competition | Inhibited (−) | Inhibited (−) | Abingdon tortoise vs feral goats; *Balanus* vs *Chthamalus* barnacles. |
| Predation | Benefited (+) | Harmed (−) | *Pisaster* starfish preying on mussels; *Cactoblastis* moth feeding on *Opuntia*. |
| Parasitism | Benefited (+) | Harmed (−) | Human liver fluke (snail/fish vectors); Asian koel brood parasitism in crow's nest. |
| Commensalism | Benefited (+) | Neutral (0) | Epiphytic orchid on mango; cattle egret and cattle; clownfish and sea anemone. |
| Amensalism | Harmed (−) | Neutral (0) | *Penicillium* mold secreting penicillin; Black walnut tree producing juglone. |

OurMocks

# Strategic Synthesis: Organisms and Populations Architecture




Figure 3: Master synthesis mindmap — high-yield integrated architecture of Organisms and Populations covering environmental adaptations, population growth dynamics, and interspecific species interactions.

## CUET HIGH-YIELD MEMORY HOOKS & DIAGNOSTIC TRIAD

1. **Organismic Adaptation Triad:** Eurythermal/euryhaline (wide tolerance) vs stenothermal/ stenohaline (narrow tolerance); Allen's rule states colder-climate mammals possess shorter ears and shorter limbs to minimize heat loss; kangaroo rat oxidizes internal fats and produces concentrated urine; high-altitude sickness is compensated by increasing RBC count, increasing breathing rate, and decreasing hemoglobin oxygen-binding affinity.
2. **Population Growth Metrics Triad:** Birth rate and death rate are per capita metrics; exponential growth follows $d\frac{N}{d}t = rN$ ($N_t = N_0 e^{rt}$, J-shaped curve); logistic growth follows $d\frac{N}{d}t = rN\frac{K-N}{K}$ (Verhulst-Pearl S-shaped sigmoid curve); maximum growth rate ($d\frac{N}{d}t$) occurs at half carrying capacity ($N = \frac{K}{2}$); carrying capacity $K$ is the habitat's maximum sustainable density.
3. **Interspecific Interactions Triad:** Predation ($\frac{+}{-}$, Paine's *Pisaster* removal caused 10 invertebrate extinctions); Competition ( $\frac{-}{-}$ , Connell's *Balanus* excludes *Chthamalus*, MacArthur's warblers avoid exclusion via resource partitioning); Parasitism ($\frac{+}{-}$, brood parasitism in cuckoo and crow); Commensalism ($\frac{+}{0}$, orchid on mango, clownfish and sea anemone); Mutualism ($\frac{+}{+}$, fig and fig wasp, lichens, mycorrhizae); Amensalism ( $\frac{-}{0}$ , *Penicillium* producing penicillin inhibiting bacteria).

# Master Diagnostic Cheatsheet & Fatal CUET Traps

## MASTER CONCEPTUAL & DIAGNOSTIC CHEATSHEET — ORGANISMS & POPULATIONS

- **Temperature & Salinity Tolerance:** Eurythermal & Euryhaline (tolerate wide ranges); Stenothermal & Stenohaline (restricted to narrow bands).
- **Ecological Rules:** Allen's Rule (colder climate mammals have shorter ears/limbs to reduce heat loss); Gloger's Rule (warm/humid animals darker); Bergmann's Rule (larger body size in colder regions).
- **High Altitude Adaptations:** Decreased oxygen partial pressure compensated by: (1) increased RBC production (erythropoietin), (2) elevated breathing rate, and (3) **decreased hemoglobin binding affinity** for oxygen (facilitating tissue oxygen release).
- **Population Parameters:** Vital rates are **per capita** (e.g., lotus birth rate $= \frac{8}{20} = 0.4$ offspring/lotus/year); age pyramids: expanding (triangular base), stable (bell-shaped), declining (urn-shaped with narrow base).
- **Population Growth Equations:** Exponential ($d\frac{N}{d}t = rN$, integrated form $N_t = N_0 e^{rt}$, J-shaped); Logistic / Verhulst-Pearl ($d\frac{N}{d}t = rN\left(\frac{K-N}{K}\right)$, S-shaped sigmoid); inflection point with peak growth rate at $N = \frac{K}{2}$.
- **Keystone Predation:** Paine's intertidal removal of starfish ***Pisaster*** caused extinction of $> 10$ invertebrate species due to unchecked competitive exclusion.
- **Competitive Principles:** Gause's Competitive Exclusion Principle (complete competitors cannot coexist on limiting resources); Connell's field experiments (*Balanus* excludes *Chthamalus*); MacArthur's Resource Partitioning (5 warbler species coexist via behavioral/spatial niches).
- **Co-evolution & Specialized Interactions:** Sexual deceit in Mediterranean orchid ***Ophrys*** (pseudocopulation by male bees); Brood parasitism (Koel lays eggs in Crow's nest); Commensalism ($\frac{+}{0}$, orchid on mango, cattle egret and grazing cattle); Amensalism ($\frac{-}{0}$, *Penicillium* secretes penicillin killing bacteria).

## THE SEVEN FATAL CUET TRAPS IN ORGANISMS AND POPULATIONS

1. **The Hemoglobin Affinity Trap:** At high altitudes, the body does NOT increase hemoglobin binding affinity; it **decreases hemoglobin affinity** so oxygen is easily unloaded to tissues!
2. **The Vital Statistics Denominator Trap:** Birth and death rates in ecology are strictly **per capita rates** (offspring per individual per time), NEVER absolute totals.
3. **The Maximum Logistic Rate Trap:** Population growth rate ($d\frac{N}{d}t$) does NOT peak at carrying capacity ($K$); at $K$, growth rate is zero! Maximum rate occurs at **half carrying capacity ($N = \frac{K}{2}$)**.
4. **The J-Shaped vs S-Shaped Equation Trap:** Exponential J-shaped curve is $N_t = N_0 e^{rt}$; Logistic S-shaped curve is $d\frac{N}{d}t = rN\frac{K-N}{K}$. Never confuse the differential and integrated formulas.
5. **The Resource Partitioning Reality Trap:** Gause's principle does NOT mean competition always leads to extinction; species frequently coexist through **resource partitioning** (e.g., MacArthur's warblers).
6. **The Amensalism vs Commensalism Sign Trap:** Commensalism is $\left(\frac{+}{0}\right)$ (orchid on mango); Amensalism is $\left(\frac{-}{0}\right)$ (*Penicillium* secreting penicillin). Never swap the negative and positive signs!
7. **The Brood Parasitism Identification Trap:** Brood parasitism is a specialized form of **Parasitism ($\frac{+}{-}$, where host crow loses fitness)**, NOT commensalism.

# Practice Questions — Part 1 (Q1 to Q5)

**Q1.** Organisms that are capable of tolerating and thriving across a wide range of ambient environmental temperatures are scientifically designated as:

(A) Eurythermal organisms
(B) Stenothermal organisms
(C) Euryhaline organisms
(D) Stenohaline organisms

**Sol. Q1.** Organisms that can tolerate and flourish across a **wide range** of temperatures are designated as **eurythermal** (from Greek *eurys*, meaning wide; e.g., most mammals and birds). In contrast, **stenothermal** organisms are restricted to a narrow temperature band (e.g., polar bears, corals). The terms **euryhaline** and **stenohaline** refer strictly to salinity tolerance, not temperature. **Answer: (A)**

**Q2.** When a person travels from the coastal plains to high-altitude regions (exceeding 3,500 meters), the human body rapidly compensates for altitude sickness through which set of physiological adjustments?

(A) Decreasing total red blood cell count, decreasing breathing rate, and increasing hemoglobin oxygen affinity
(B) Increasing red blood cell production, accelerating breathing rate, and decreasing hemoglobin binding affinity for oxygen
(C) Increasing blood plasma volume, halting renal filtration, and permanently constricting peripheral capillaries
(D) Suppressing erythropoietin secretion from the kidneys and increasing cellular reliance on anaerobic lactic fermentation

**Sol. Q2.** At high altitudes, low atmospheric pressure reduces the partial pressure of oxygen. The human body compensates through three distinct physiological adaptations: (1) Kidneys secrete erythropoietin, **increasing total red blood cell count**; (2) The respiratory center accelerates **breathing rate**; and (3) The body **decreases the binding affinity of hemoglobin for oxygen**, which accelerates the release of oxygen to oxygen-deprived tissues. **Answer: (B)**

**Q3.** Allen's Rule, a well-established macro-ecological generalization observed across endothermic animals, states that mammals inhabiting colder geographical climates tend to possess:

(A) Shorter ears and shorter limbs to minimize body heat dissipation
(B) Larger body surface-to-volume ratios to maximize convective heat loss
(C) Thinner layers of subcutaneous adipose tissue to accelerate cellular metabolism
(D) Extremely long external appendages to promote efficient directional thermoregulation

**Sol. Q3.** **Allen's Rule** states that mammals living in colder climates generally possess **shorter ears and shorter limbs (extremities)** to minimize convective and conductive heat loss through reduced exposed surface area. Thick subcutaneous fat (blubber) is an adaptation of marine mammals, but is not Allen's rule. **Answer: (A)**

**Q4.** In a wild population of lotus plants inhabiting an undisturbed freshwater pond, there were initially 20 plants. Over a one-year reproductive cycle, 8 new lotus plants were born through vegetative reproduction, while zero plants died. What is the per capita birth rate of this population?

(A) 0.8 offspring per lotus per year
(B) 2.5 offspring per lotus per year
(C) 0.2 offspring per lotus per year
(D) 0.4 offspring per lotus per year

**Sol. Q4.** In population ecology, birth rate is calculated as a per capita metric:

$$\text{Birth rate} = \frac{\text{Number of newly born individuals}}{\text{Initial total population}} = \frac{8}{20} = 0.4 \text{ offspring per lotus per year.}$$

Option D is the exact mathematically verified value. **Answer: (D)**

**Q5.** An ecological age pyramid displaying a narrow base with pre-reproductive individuals being significantly fewer than reproductive individuals indicates that the population is:

(A) Rapidly expanding exponentially
(B) Completely stable and stationary
(C) Declining towards negative growth
(D) Undergoing severe competitive exclusion

**Sol. Q5.** An age pyramid characterized by a **narrow base** where the pre-reproductive cohort is smaller than the reproductive age group is termed an **urn-shaped pyramid**. Because fewer young individuals are entering the breeding pool, birth rates fall below death rates, indicating that the population is **declining**. A wide triangular base indicates expanding growth; a bell-shaped pyramid indicates a stable population. **Answer: (C)**

# Practice Questions — Part 2 (Q6 to Q10)

**Q6.** Under conditions of unlimited resources (food and space), a biological population exhibits exponential growth. Which mathematical equation and graphical curve correctly represent this unrestricted growth?

**(A)** Verhulst-Pearl logistic equation $d\frac{N}{d}t = rN\left(\frac{K-N}{K}\right)$, producing an S-shaped sigmoid curve
**(B)** Integrated logistic equation $N_t = N_0 + (b-d)t$, producing a linear horizontal plateau
**(C)** Density-dependent carrying capacity equation $d\frac{N}{d}t = \frac{K}{rN}$, producing a parabolic hyperbola
**(D)** Exponential growth equation $d\frac{N}{d}t = rN$ (integrated form $N_t = N_0 e^{rt}$), producing a steep J-shaped curve

**Sol. Q6.** When resources in the habitat are unlimited, population growth is exponential, following the differential equation $d\frac{N}{d}t = rN$ (where $r = b - d$ is the intrinsic rate of natural increase). Integrating this equation over time yields the exponential formula $N_t = N_0 e^{rt}$, producing a steep **J-shaped curve**. Option A is the logistic differential equation, not exponential. **Answer: (D)**

**Q7.** In the Verhulst-Pearl logistic growth model represented by the equation $d\frac{N}{d}t = rN\left(\frac{K-N}{K}\right)$, the population growth rate reaches its maximum absolute value when population density ($N$) is equal to:

**(A)** Carrying capacity ($N = K$)
**(B)** Zero ($N = 0$)
**(C)** Half of the carrying capacity ($N = \frac{K}{2}$)
**(D)** Double the carrying capacity ($N = 2K$)

**Sol. Q7.** In logistic growth, the growth rate $d\frac{N}{d}t$ is a parabolic function of population density $N$. Growth is zero when $N = 0$ (no individuals) and drops to zero when $N = K$ (because $\frac{K-K}{K} = 0$). The rate of population increase reaches its absolute peak (**inflection point**) midway along the sigmoid curve, exactly at **half the carrying capacity ($N = \frac{K}{2}$)**. **Answer: (C)**

**Q8.** The removal of the predatory starfish *Pisaster* from its natural rocky intertidal habitat in a celebrated field experiment conducted by Robert Paine resulted in:

**(A)** A dramatic increase in the species richness and biodiversity of the intertidal community
**(B)** The rapid extinction of all top carnivorous marine birds feeding in the rocky intertidal zone
**(C)** The extinction of more than 10 species of intertidal invertebrates within a year due to interspecific competition
**(D)** A permanent shift towards mutualistic symbiosis between barnacles and sea anemones

**Sol. Q8.** Ecologist Robert Paine demonstrated that the predatory starfish ***Pisaster*** acts as a keystone predator in intertidal rocky communities. By preying upon competitive dominants (such as mussels), *Pisaster* keeps prey densities below competitive exclusion thresholds. When all *Pisaster* were removed, **more than 10 species of invertebrates became extinct within a single year** due to intense, unchecked interspecific competition. **Answer: (C)**

**Q9.** The Monarch butterfly avoids predation by insectivorous birds through which sophisticated defensive mechanism?

**(A)** Possessing brilliant camouflage coloration that blends seamlessly into tree foliage
**(B)** Secreting a blinding neurotoxic aerosol from specialized abdominal repellent glands
**(C)** Stinging predatory birds with venomous barbs located on the distal margin of its wings
**(D)** Accumulating a bitter, toxic cardiac glycoside during its larval stage by feeding on poisonous milkweed

**Sol. Q9.** The **Monarch butterfly** is exceptionally distasteful to birds due to a special chemical defense. It does not synthesize this poison as an adult; rather, it acquires and accumulates a potent **cardiac glycoside** toxin during its **caterpillar stage by feeding on a poisonous milkweed plant**. Birds learn to avoid Monarchs based on visual recognition and bitter toxicity. **Answer: (D)**

**Q10.** Gause's Competitive Exclusion Principle states that two closely related species competing for the exact same limiting resources cannot coexist indefinitely. Coexistence between competing species is frequently enabled in nature through:

**(A)** Complete mutualistic merging of the two genomes through horizontal gene transfer
**(B)** Resource partitioning, where competing species evolve differing foraging times or distinct spatial feeding niches
**(C)** One species abandoning heterotrophic nutrition to become fully autotrophic photosynthetic symbionts
**(D)** Periodic synchronized reproductive diapause that eliminates predatory pressure entirely

**Sol. Q10.** While Gause's principle predicts that the inferior competitor will be driven to local extinction if resources are strictly limiting, competing species frequently avoid exclusion through **Resource Partitioning**. Robert MacArthur proved that five closely related species of **Warblers** successfully coexisted on the same spruce tree by foraging in different zones of the canopy and hunting at different times of day. **Answer: (B)**

# Practice Questions — Part 3 (Q11 to Q15)

**Q11.** In the rocky intertidal sea coasts of Scotland, field experiments conducted by Joseph Connell demonstrated that the superior barnacle *Balanus* competitively excluded which species from the upper intertidal zone?

**(A)** *Chthamalus* **(B)** *Pisaster*
**(C)** *Abingdon* tortoise **(D)** *Ophrys*

**Sol. Q11.** Connell's classic field experiments on the rocky sea coast of Scotland demonstrated that the larger, competitively dominant barnacle ***Balanus*** occupies the intertidal zone and competitively excludes the smaller barnacle ***Chthamalus*** from that favorable intertidal zone. **Answer: (A)**

**Q12.** Which of the following biological associations represents a classic ecological example of Brood Parasitism?

**(A)** A female mosquito sucking blood from a human host during the night **(B)** A sea anemone attached to the dorsal carapace of a hermit crab
**(C)** An Asian koel (cuckoo) laying its eggs in the nest of a crow **(D)** A dodder plant (*Cuscuta*) growing haustoria into the vascular bundles of a hedge plant

**Sol. Q12.** **Brood parasitism** is an extreme behavioral adaptation in birds where a parasitic bird lays its eggs in the nest of another host species, leaving the host to incubate and feed the hatchlings. The classic NCERT example is the **Asian Koel (cuckoo) laying its eggs in the nest of the crow**. Over evolution, the cuckoo's eggs have evolved to closely mimic the host's eggs in size and coloration to avoid ejection. **Answer: (C)**

**Q13.** Which of the following species pairs exhibits a Commensal $(\frac{+}{0})$ relationship where one organism benefits while the other is completely unaffected?

**(A)** An epiphytic orchid growing on the branch of a mango tree **(B)** A fungus and a photosynthetic cyanobacterium living mutualistically as a lichen
**(C)** A tiger stalking, capturing, and consuming a spotted deer in a forest **(D)** *Penicillium* mold secreting penicillin antibiotic that destroys bacterial colonies

**Sol. Q13.** An **epiphytic orchid growing on a mango tree branch** is a classic **commensalism** $(\frac{+}{0})$ association. The orchid derives physical support, height, and access to sunlight, while the mango tree receives zero benefit and suffers zero harm (neutral, score 0). Option B is mutualism (+/+), C is predation (+/-), and D is amensalism (-/0). **Answer: (A)**

**Q14.** The Mediterranean orchid *Ophrys* secures cross-pollination by male bees through a unique behavioral phenomenon termed "pseudocopulation". This evolutionary strategy relies on:

**(A)** Secreting vast droplets of sweet, energy-rich nectar that nourish the foraging male bee **(B)** Sexual deceit, where one floral petal mimics the female bee in shape, color, and scent
**(C)** Synthesizing potent insecticidal toxins that paralyze the male bee until cross-pollination is complete **(D)** Releasing airborne ultrasound waves that mimic the buzzing frequencies of female bees in flight

**Sol. Q14.** The Mediterranean orchid ***Ophrys*** employs **sexual deceit**. One petal of its flower exhibits an uncanny resemblance to the female bee in anatomical form, color markings, and chemical scent. The male bee mistakes the petal for a female and attempts to copulate with it (**pseudocopulation**), during which pollen is transferred. If the female bee evolves morphological changes, the orchid must **co-evolve** matching modifications to maintain pollination. **Answer: (B)**

**Q15.** The ecological relationship known as Amensalism is designated by the sign interaction $(\frac{-}{0})$. Which of the following is a verified real-world example of amensalism?

**(A)** Cattle egrets foraging near grazing cattle to capture insects flushed from the grass **(B)** *Penicillium* mold secreting penicillin that inhibits surrounding bacterial growth without receiving direct benefit
**(C)** Clownfish seeking shelter and physical protection among stinging sea anemone tentacles **(D)** Barnacles growing attached to the thick dorsal skin of a migratory baleen whale

**Sol. Q15.** **Amensalism** $(\frac{-}{0})$ is an association where one species is inhibited or harmed, while the second organism is unaffected. The classic NCERT case is **antibiosis**, where the mold ***Penicillium*** secretes penicillin, destroying neighboring bacteria (−) while the mold itself is completely unaffected and gains no direct nutritional benefit (0). Cattle egrets, clownfish, and barnacles on whales are all commensal (+/0) associations. **Answer: (B)**

© OurMocks · www.ourmocks.com · CUET UG Biology · Organisms and Populations · 2027 Cycle