OurMocks

2027 CYCLE

MOCK · PRACTICE · CRACK

CUET UG

Biology

Biodiversity and Conservation

Official Syllabus Unit X · Class 12 · Comprehensive NCERT Theory, Diagrams & 15 Solved MCQs

INSIDE
Concept Notes

INSIDE
Solved Examples

INSIDE
Practice MCQs + Solutions

Aligned to the latest official syllabus · Concept-first explanations

www.ourmocks.com

Syllabus Alignment & Exam Blueprint

Sub-Topic	Official Syllabus Core Coverage (NCERT Class 12 Unit X)	Exam Focus & Question Types
Biodiversity Concept & Levels	Definition by Edward Wilson (1992); three hierarchical levels: Genetic diversity (*Rauvolfia vomitoria* reserpine potency/concentration, >50,000 rice strains, >1,000 mango varieties), Species diversity (Western Ghats > Eastern Ghats for amphibians), and Ecological diversity (India's deserts, rainforests, mangroves vs Norway); global species estimates (Robert May: 7 million species, 22% documented).	Identifying the three levels from real-world examples; taxonomic proportions (insects >70% of animals, fungi > combined vertebrates).
Patterns of Biodiversity	Latitudinal diversity gradient (equator to poles; Colombia 1,400 birds vs Greenland 56 birds; Amazonian rainforest: >40,000 plants, 3,000 fishes, 1,300 birds, 427 mammals/amphibians, >125,000 invertebrates); 3 evolutionary hypotheses explaining tropical richness; Alexander von Humboldt's species-area relationship ($S = CA^Z$, $\log S = \log C + Z \log A$; slope Z values: 0.1–0.2 small regions, 0.6–1.2 continents, 1.15 frugivores).	Interpreting Z slope steepness; naming 3 tropical richness hypotheses; species counts in Amazon and latitudinal gradients.
Importance of Biodiversity	David Tilman's long-term outdoor ecosystem plots (diversity = biomass stability + higher productivity); Paul Ehrlich's Rivet Popper hypothesis (airplane = ecosystem, rivets = species, wing rivets = keystone species); reasons to conserve: narrowly utilitarian (food, timber, bioprospecting, >25% modern medicines from plants), broadly utilitarian (Amazon produces 20% O_2, pollination, soil), ethical arguments.	Differentiating utilitarian categories; interpreting Rivet Popper wing analogy; David Tilman's experimental conclusions.
Biodiversity Loss & The Evil Quartet	IUCN Red List 2004 (784 species extinct in 500 yrs: 338 vertebrates, 359 invertebrates, 87 plants); recent extinctions: Dodo (Mauritius), Quagga (Africa), Thylacine (Australia), Steller's sea cow (Russia), 3 tiger subspecies (Bali, Javan, Caspian); current 6th mass extinction (100–1000x faster, anthropogenic); The Evil Quartet: 1. Habitat loss & fragmentation (1 driver; Amazon cleared for soy/cattle), 2. Over-exploitation, 3. Alien species invasions (Nile perch, *Lantana*, *Eichhornia*, *Parthenium*, *Clarias gariepinus*), 4. Co-extinctions (obligate mutualisms).	Matching extinct species to home countries; identifying the 4 drivers of the Evil Quartet; analyzing invasive weed examples.
Conservation Strategies & Summits	In-situ conservation (on-site): 14 Biosphere Reserves, 90 National Parks, 448 Wildlife Sanctuaries, 34 Global Hotspots (Western Ghats-Sri Lanka, Indo-Burma, Himalaya; cover under 2% land, cut extinction by 30%), Sacred Groves (Khasi & Jaintia in Meghalaya, Aravalli in Rajasthan, Western Ghats, Chanda & Bastar in MP); Ex-situ conservation (off-site): Zoological parks, Botanical gardens, Safari parks, Cryopreservation (-196°C in liquid N_2), Seed/Pollen banks, Tissue culture; International summits: Earth Summit (Rio, 1992), World Summit (Johannesburg, 2002; 190 countries).	Classifying in-situ vs ex-situ methods; identifying sacred grove locations; recalling hotspot criteria and international treaty years.

OurMocks

EXAM PATTERN

CUET UG Scoring Blueprint: Biodiversity and Conservation accounts for **3 to 4 questions (15 to 20 marks)** in the CUET UG Biology paper. NTA tests **exact numerical values** ($Z = 1.15$, −196°C, 784 extinctions, 8.1% Indian diversity), **the four components of The Evil Quartet**, **in-situ vs ex-situ classification**, and **locations of sacred groves**.

The Concept of Biodiversity & Its Three Levels

The term **Biodiversity** was popularized in 1992 by the renowned sociobiologist **Edward Wilson** to describe the combined diversity and biological variation occurring at all levels of biological organization, spanning from macromolecules within cells to whole biomes.

The Three Hierarchical Levels of Biodiversity

Biodiversity is structured across three fundamental hierarchical tiers:

1. **Genetic Diversity (Diversity within a Species):**
   - Represents the total variety of genetic information and allelic variations contained within individual organisms of a single species across its geographical distribution range.
   - **High-Yield NCERT Examples:**
     - The medicinal plant *Rauvolfia vomitoria* growing across different Himalayan ranges shows substantial genetic variation in the **potency and concentration of the active chemical reserpine** that it synthesizes.
     - India harbors more than **50,000 genetically distinct strains of rice** (*Oryza sativa*) and more than **1,000 cultivated varieties of mango** (*Mangifera indica*).
   - High genetic diversity provides ecological resilience, enabling species to adapt to environmental stresses, pathogens, and climatic shifts.
2. **Species Diversity (Diversity among Species within a Community):**
   - Represents the richness and variety of different taxonomic species found within a specific biological community or geographical region.
   - **High-Yield NCERT Example:**
     - The **Western Ghats** harbor a significantly greater **amphibian species diversity** than the **Eastern Ghats**.
3. **Ecological / Community Diversity (Diversity of Ecosystems):**
   - Represents the structural and functional variety of habitats, biotic communities, and ecological processes operating within a vast geographical landscape.
   - **High-Yield NCERT Example:**
     - India, with its extensive variety of **deserts, tropical rainforests, coastal mangroves, coral reefs, wetlands, estuaries, and alpine meadows**, exhibits immensely greater ecosystem diversity than a Scandinavian country like **Norway**.

Biodiversity Level	Core NCERT Mechanism / Definition	Key Diagnostic Example
Genetic Diversity	Allelic and genetic variations occurring **within a single species** across geographical ranges.	*Rauvolfia* reserpine potency; 50,000 rice strains; 1,000 mango varieties.
Species Diversity	Richness and variety of **different taxonomic species** inhabiting a community.	Western Ghats amphibians > Eastern Ghats amphibians.
Ecological Diversity	Diversity of **habitats, biomes, and ecological niches** across a geographical territory.	India (deserts to coral reefs) vs Norway.

Global Species Richness & Robert May's Estimate

- **Total Described Species:** According to the International Union for Conservation of Nature (IUCN, 2004), the total number of documented plant and animal species is slightly more than **1.5 million**.
- **Robert May's Global Estimate:**
  - Biologists have made varied estimates ranging from 20 to 50 million species.
  - However, ecologist **Robert May** proposed a sound, scientifically conservative global estimate of approximately **7 million species**.
  - This implies that **only 22% of all living species on Earth have been scientifically recorded and described so far**! More than 78% of planetary biodiversity remains undiscovered, predominantly in tropical rainforests and deep oceans.

Global Taxonomic Patterns (NCERT Figure 15.1 Breakdown)

- **Animals vs Plants:** Animals constitute **more than 70%** of all recorded living species, whereas plants (including angiosperms, gymnosperms, pteridophytes, bryophytes, algae, and fungi) make up **no more than 22%**.

- **Dominance of Insects:** Among animals, **insects** are the most species-rich taxonomic group, accounting for **more than 70% of all animal species**. This means **7 out of every 10 animals on Earth are insects**!
- **Fungi Dominance:** The total number of **fungi species** globally exceeds the combined total species of **fishes + amphibians + reptiles + mammals**!

India's Megadiversity & Latitudinal Diversity Gradients

India as a Megadiverse Nation

Although India occupies only **2.4% of the world's total land area**, its share of the global species diversity is an astonishing **8.1%**. This remarkable biodiversity places India among the **12 recognized megadiversity countries of the world**.

- Approximately **45,000 species of plants** and nearly **twice as many species of animals** (over 90,000–100,000 species) have been scientifically recorded from India.
- If we apply Robert May's global estimate (where only 22% of species are discovered), India is estimated to harbor more than **100,000 undiscovered plant species** and over **300,000 undiscovered animal species**!

Patterns of Biodiversity: Latitudinal Diversity Gradient

Biodiversity is not uniformly distributed across the globe. Instead, it follows highly consistent biogeographical patterns:

- **The General Rule:** Species diversity decreases progressively as we move away from the **equator toward the poles**. The tropical belt (situated between 23.5° N and 23.5° S) harbors significantly greater species richness than temperate or polar regions.
- **Bird Species Gradients Across Latitudes:**
  - **Colombia** (located near the equator, 0°): Harbors nearly **1,400 species of birds**.
  - **India** (situated largely in tropical latitudes): Harbors **more than 1,200 species of birds**.
  - **New York** (temperate latitude, 41° N): Harbors **105 species of birds**.
  - **Greenland** (sub-polar latitude, 71° N): Harbors only **56 species of birds**.
  - A tropical forest tract in **Ecuador** contains up to **10 times** as many species of vascular plants as an equal-sized temperate forest in the Midwest USA.

The Amazonian Rainforest: "The Lungs of the Planet"

The tropical Amazonian rainforest in South America holds the **greatest biodiversity on Earth**:

- Plants: **Over 40,000 species** of vascular flora.
- Fishes: **Over 3,000 species** in Amazon river basins.
- Birds: **1,300 species**.
- Mammals: **427 species**.
- Amphibians: **427 species**.
- Reptiles: **378 species**.
- Invertebrates: **Over 125,000 species** (with scientists estimating that at least 2 million insect species are still waiting to be discovered and named).

Three Hypotheses Explaining Tropical Species Richness

Why do the tropics harbor such exceptionally high biodiversity compared to temperate zones? Ecologists propose three mutually reinforcing hypotheses:

1. **Evolutionary Speciation Time:** Temperate regions were subjected to frequent, catastrophic glaciations (ice ages) in geological history, wiping out existing biota and repeatedly resetting evolutionary clocks. In contrast, **tropical latitudes have remained relatively undisturbed and ecologically stable for millions of years**, allowing a prolonged, uninterrupted period for evolutionary speciation and diversification.
2. **Environmental Constancy & Niche Specialization:** Tropical environments are less seasonal, relatively more constant, and highly predictable compared to harsh temperate climates with extreme freezing winters. Such predictable, stable environmental conditions promote **niche specialization**, allowing a large number of co-occurring species to occupy narrow, specialized ecological niches without competitive exclusion.
3. **Higher Radiant Solar Energy & Productivity:** The tropics receive maximum direct solar insolation throughout the year. Greater solar energy drives **higher primary productivity** (GPP and NPP), which provides an abundant energetic foundation supporting complex, species-rich consumer food webs.




Figure 1: Figure 1: High-yield NCERT analytical framework — Humboldt's Species-Area relationship ($S = CA^Z$, logarithmic slope Z), global latitudinal diversity gradients (Amazonian rainforest richness), and Paul Ehrlich's Rivet Popper ecosystem stability hypothesis.

Key Analytical Takeaways from Figure 1:

1. **Humboldt's Curve:** On a normal arithmetic scale, the species-area relationship is a **rectangular hyperbola**. On a logarithmic scale, it converts into a **straight line** defined by $\log S = \log C + Z \log A$.
2. **Slope (Z) Values:** For small to regional areas, Z remains between **0.1 and 0.2**. For continental areas, Z steepens to **0.6 to 1.2**. For frugivorous birds and mammals in tropical forests across continents, $Z = 1.15$.
3. **Airplane Analogy:** Popping wing rivets (keystone species) directly threatens flight safety (ecosystem collapse), whereas popping seat rivets causes minor functional impairment.

Species-Area Relationship & Ecosystem Stability

Alexander von Humboldt's Species-Area Law

During his extensive pioneering explorations in the wilderness of South American jungles, the great German naturalist and geographer **Alexander von Humboldt** observed that within a given region, **species richness increases with increasing explored area, but only up to a limit**.

Mathematical Formulation

- **Linear Scale Formulation:**

$$S = C \cdot A^Z$$

Where: S = Species richness; A = Explored geographical area; C = Y-intercept (constant reflecting taxon density at unit area); Z = Slope of the regression line (regression coefficient). On a standard Cartesian coordinate graph (S vs A), the curve forms a **rectangular hyperbola**.

- **Logarithmic Scale Formulation:**

$$\log S = \log C + Z \log A$$

On a logarithmic scale, the relationship is represented by a **straight line**, where Z is the exact slope of the line.

Numerical Interpretation of the Regression Coefficient (Z)

- **Small to Medium Regional Areas ($Z = 0.1$ to 0.2):** Regardless of taxonomic group or continent (plants in Britain, birds in California, molluscs in New York), slope falls in the narrow range of **0.1 to 0.2**.
- **Continental Scale Areas ($Z = 0.6$ to 1.2):** When analyzed across enormous continental scales, the regression slope becomes **significantly steeper**, with Z values ranging between **0.6 and 1.2**.
- **Frugivorous Birds and Mammals ($Z = 1.15$):** In tropical forests across different continents, slope Z is measured at **1.15** (high-yield CUET numerical value).

Does Species Diversity Matter to Ecosystem Functioning?

David Tilman's Long-Term Outdoor Plot Experiments

Ecologist **David Tilman** conducted definitive long-term outdoor field experiments on Minnesota grassland plots, providing quantitative empirical proof:

1. **Biomass Stability:** Experimental plots with **higher species diversity showed less year-to-year variation in total biomass** compared to species-poor plots.
2. **Higher Productivity:** Increased species diversity directly contributed to **higher overall primary productivity** due to complementary resource utilization.

Paul Ehrlich's "Rivet Popper" Hypothesis

Stanford ecologist **Paul Ehrlich** proposed a brilliant analogy:

- An ecosystem is compared to an **airplane**, and species to the **rivets** holding parts together.
- Popping rivets from seats/windows (minor species extinction) causes minor impairment.
- But popping rivets from the **wings (keystone species like pollinators or apex predators)** poses an immediate, catastrophic threat to flight safety (ecosystem collapse).

Biodiversity Loss, Extinctions & The Evil Quartet

The Contemporary Extinction Crisis & IUCN Red List Data

Human colonization and industrial exploitation have accelerated global species extinctions:

- **Dodo:** Extinct from **Mauritius**; **Quagga:** Extinct from **Africa**; **Thylacine:** Extinct from **Australia**; **Steller's Sea Cow:** Extinct from **Russia / Bering Sea**; **Three Tiger Subspecies: Bali, Javan, and Caspian** tigers went extinct in the 20th century.
- **IUCN Red List (2004):** Records the verified extinction of **784 species** in the last 500 years (**338 vertebrates**, **359 invertebrates**, and **87 plants**).

The Sixth Mass Extinction

- We are currently witnessing the **Sixth Mass Extinction** ("Anthropocene Extinction").
- Current extinction rates are estimated to be **100 to 1,000 times faster** than pre-human background rates, driven entirely by human activities. Nearly 50% of species could be lost within a century.
- **Most Vulnerable Taxa:** Globally, 12% of bird, 23% of mammal, 31% of gymnosperm, and **32% of amphibian species** face extinction. Amphibians are the most vulnerable taxon worldwide.

The "Evil Quartet" — The Four Major Drivers of Biodiversity Loss

1. **Habitat Loss and Fragmentation (1 Leading Cause):** Tropical rainforests once covered **14% of the Earth's land surface**; today they cover **less than 6%**. The Amazon forest is being cleared for **soybeans** and **beef cattle pasture**. Fragmentation severely hits animals requiring large home ranges and migratory habits.
2. **Over-Exploitation (Human Need vs Greed):** Over-harvesting beyond reproduction replacement capacity drove **Steller's sea cow** and the **passenger pigeon** to extinction.
3. **Alien Species Invasions:** Introduction of **Nile perch** into Lake Victoria caused extinction of **over 200 endemic cichlid fishes**. Invasive weeds: ***Parthenium*** (carrot grass), ***Lantana camara***, and ***Eichhornia crassipes*** (water hyacinth). African catfish (***Clarias gariepinus***) threatens native Indian catfishes.
4. **Co-Extinctions:** Obligate dependency: extinction of host fish causes extinction of its **host-specific parasites**; extinction of fig wasp causes extinction of partner fig tree (***Ficus***).

OurMocks




Figure 2: Figure 2: Systematic comparison matrix — In-situ (on-site) vs Ex-situ (off-site) biodiversity conservation methodologies and the detailed structural breakdown of The Evil Quartet driving global extinctions.

Key Comparative Pointers from Figure 2:

1. **In-situ Strategy:** Conserves the whole ecosystem; ideal whenever habitat is intact. Encompasses National Parks (strictly zero human activity), Wildlife Sanctuaries (controlled collection allowed), Biosphere Reserves (zoning), Hotspots, and Sacred Groves.
2. **Ex-situ Strategy:** Protects endangered organisms away from their natural home in specialized facilities. Encompasses Zoological Parks, Botanical Gardens, Safari Parks, Seed Banks, and Cryopreservation (-196°C in liquid N_2).
3. **The Evil Quartet:** Habitat loss & fragmentation (1 driver), Over-exploitation (need to greed), Alien invasions (Nile perch, *Clarias*), and Co-extinctions (obligate mutualisms).

Why Conserve Biodiversity? Three Utilitarian Arguments

1. Narrowly Utilitarian Arguments (Direct Tangible Products)

Human beings derive enormous, quantifiable direct economic benefits:

- **Food & Sustenance:** Cereals, pulses, fruits, vegetables, livestock, and marine fisheries.
- **Forest & Industrial Products:** Firewood, construction timber, pulp, tannins, resins, gums, latex, perfumes.
- **Modern Pharmaceuticals: More than 25% of modern drugs** are derived from plants; 25,000 species used in traditional medicines.
- **Bioprospecting:** Exploring molecular, genetic, and species-level diversity for commercially valuable products.

2. Broadly Utilitarian Arguments (Ecosystem Services)

Vital life-support services that cannot be assigned a simple market price tag:

- **Oxygen Production:** The Amazonian rainforest produces an estimated **20% of total oxygen** in Earth's atmosphere.
- **Pollination:** Bees, bumblebees, butterflies, moths, birds, and bats pollinate agricultural crops and wild plants.
- **Flood & Drought Mitigation:** Forest tree roots bind soil, absorb monsoon run-off, replenish groundwater tables, and prevent erosion.
- **Aesthetic Pleasures:** Psychological and aesthetic joy from walking through forests and observing nature.

3. Ethical Arguments (Moral Duty of Stewardship)

Every biological species has an **intrinsic value**. We hold a moral duty to care for their well-being and pass our biological legacy safely to future generations.

OurMocks

In-situ & Ex-situ Conservation Strategies

In-situ Conservation (On-Site Protection)

Conserving and protecting the entire natural ecosystem simultaneously protects all species:

- **Biodiversity Hotspots:** Exceptional species richness and high endemism facing accelerated habitat destruction. 34 globally covering under 2% land; protecting them reduces extinction rates by 30%. Three cover India: **Western Ghats-Sri Lanka**, **Indo-Burma**, and **Himalaya**.
- **Protected Area Network in India:**
  - **14 Biosphere Reserves:** Conserving whole ecosystems under UNESCO MAB Program (Core, Buffer, Transition zones).
  - **90 National Parks:** Strictly for wildlife betterment; **strictly zero private ownership, settlement, forestry, or grazing**.
  - **448 Wildlife Sanctuaries:** Focus on wild fauna; regulated timber collection/grazing permitted without disturbing wildlife.
- **Sacred Groves (Tribal Refugia):** Ancient tracts where trees and wildlife are protected by religious veneration:
  1. **Khasi and Jaintia Hills in Meghalaya:** Last refuges for numerous rare and threatened endemic plants.
  2. **Aravalli Hills in Rajasthan:** Protected by Bishnoi and local communities.
  3. **Western Ghats regions of Karnataka and Maharashtra**.
  4. **Sarguja, Chanda, and Bastar areas of Madhya Pradesh and Chhattisgarh**.

Ex-situ Conservation (Off-Site Protection)

Endangered organisms taken out of natural habitats into specialized facilities under human care:

- **Zoological Parks, Botanical Gardens & Safari Parks:** Captive management and live plant collections.
- **Cryopreservation (-196°C in Liquid Nitrogen):** Freezing viable, fertile gametes for indefinitely long periods.
- **In Vitro Tissue Culture / Micropropagation:** Clonal propagation of explants on synthetic nutrient media.
- **Seed Banks & Pollen Banks:** Dehydrated cold storage of genetic strains for decades.

International Conventions & Treaties on Biodiversity

- **The Earth Summit (Rio de Janeiro, 1992):** UN conference calling for biodiversity conservation, sustainable utilization, and fair benefit-sharing.
- **World Summit on Sustainable Development (Johannesburg, 2002): 190 countries** pledged to achieve by 2010 a significant reduction in global biodiversity loss rates.

Key Facts & Terms Cheatsheet

MASTER DIAGNOSTIC & CONCEPTUAL CHEATSHEET — BIODIVERSITY AND CONSERVATION

Key Subject Pointer	Core NCERT Mechanism / Definition	Exam Identifier
Edward Wilson (1992)	Sociobiologist who popularized the term "Biodiversity" for all organizational tiers.	Popularized Term
Genetic Diversity	Allelic variation within one species (*Rauvolfia* reserpine, 50,000 rice strains).	Within One Species
Species Diversity	Variety of species in a region (Western Ghats amphibians > Eastern Ghats).	Among Species
Ecological Diversity	Ecosystem variety across biomes (India deserts to coral reefs vs Norway).	Ecosystem Variety
Total Described Species	Over 1.5 million species documented globally (IUCN 2004).	>1.5 Million
Robert May's Estimate	Scientifically conservative global estimate of approximately 7 million species.	7 Million Species
Percentage Discovered	Only 22% of total planetary species have been scientifically documented.	22% Documented
Taxonomic Proportions	Animals >70%, Plants 22%, Insects >70% of animals, Fungi > all vertebrates.	Insects > 70%
India's Land vs Diversity	India occupies 2.4% world land area, but holds 8.1% of global species diversity.	2.4% Land, 8.1% Species
Latitudinal Gradient	Species richness peaks at equator and declines steadily toward polar latitudes.	Equator to Poles
Amazonian Rainforest	Greatest biodiversity on Earth: 40k plants, 3k fishes, 1.3k birds, 427 mammals.	Lungs of the Planet
Species-Area Law	Rectangular hyperbola: $S = CA^Z$; log straight line: $\log S = \log C + Z \log A$.	Alexander von Humboldt
Z-value (Small Regions)	Slope $Z = 0.1$ to 0.2 for small to regional areas regardless of taxon/continent.	0.1 to 0.2
Z-value (Continents)	Slope $Z = 0.6$ to 1.2 for entire continental land masses.	0.6 to 1.2
Z-value (Frugivores)	Slope $Z = 1.15$ for frugivorous birds and mammals in tropical forests.	Z = 1.15
David Tilman's Finding	Outdoor plots with higher diversity showed less biomass variation and higher yield.	Diversity = Stability
Rivet Popper Hypothesis	Paul Ehrlich's airplane analogy: wing rivets = keystone species vital to safety.	Wing Rivet = Keystone
IUCN Red List (2004)	784 species extinct in 500 yrs: 338 vertebrates, 359 invertebrates, 87 plants.	784 Extinct Species
Recent Extinction Flagships	Dodo (Mauritius), Quagga (Africa), Thylacine (Australia), Steller's sea cow (Russia).	Species-Country Match
The Evil Quartet	1. Habitat loss & fragmentation (1 driver), 2. Over-exploitation, 3. Alien invasion, 4. Co-extinction.	4 Major Drivers
Alien Species Examples	Nile perch in Lake Victoria; *Parthenium*; *Lantana*; *Eichhornia*; *Clarias gariepinus*.	Exotic Invasives
Narrowly Utilitarian	Direct consumable economic benefits: food, timber, fiber, >25% modern medicines.	Direct Economic Goods
Broadly Utilitarian	Indirect ecosystem services: Amazon 20% O_2, insect pollination, flood control.	Ecosystem Services
In-situ Conservation	14 Biosphere Reserves, 90 National Parks, 448 Wildlife Sanctuaries, Sacred Groves.	On-Site Protection
Biodiversity Hotspots	34 globally (under 2% land, cut extinction by 30%); 3 in India: Western Ghats, Indo-Burma, Himalaya.	High Endemism + Loss
Sacred Groves	Khasi & Jaintia (Meghalaya), Aravalli (Rajasthan), Western Ghats, Chanda & Bastar (MP).	Tribal Refugia
Ex-situ Conservation	Zoos, Botanical gardens, Safari parks, Cryopreservation (-196°C), Seed banks.	Off-Site Protection
Global Summits	Earth Summit (Rio 1992); World Summit (Johannesburg 2002; 190 countries).	Rio 1992 & Joburg 2002

Common Errors and High-Yield CUET Traps

COMMON STUDENT MISTAKES & DIAGNOSTIC TRAPS — BIODIVERSITY

Common Student Mistake	Scientifically Verified Reality	The Exact Trap Mechanism
Confusing Endemic with Extinct	**Endemic** means confined exclusively to a given geographic region and found nowhere else. **Extinct** means permanently lost from Earth.	NTA options present "Endemic species are those that have already vanished from Earth." Endemism describes geographic distribution restriction, not extinction status.
Ignoring Both Hotspot Criteria	A region qualifies as a biodiversity hotspot ONLY if it meets **BOTH criteria simultaneously**: (1) Exceptionally high species richness and endemism, AND (2) High degree of habitat threat/destruction.	NTA questions test whether students pick areas simply with "high species count." High richness alone does not create a hotspot; accelerated habitat loss is mandatory.
Classifying Sacred Groves as Ex-situ	Sacred groves are **100% IN-SITU conservation**. The undisturbed forest trees and wildlife are protected right in their native ancestral habitat by tribal faith.	Students mistakenly associate "religious veneration" with human management and select ex-situ. Sacred groves are natural ecosystems protected on-site.
Classifying Nile Perch under Over-Exploitation	The extinction of 200+ cichlid fishes in Lake Victoria is the classic NCERT example of **Alien Species Invasions**, NOT over-exploitation.	Students see fish deaths and incorrectly assume over-fishing. The cichlids were decimated because an introduced exotic predator was added to the lake.
Misidentifying Cryopreservation Temperature	Cryopreservation stores living gametes in **liquid nitrogen at exactly −196°C** (Celsius), NOT −196°F or 0°C.	NTA examiners switch units or signs, offering +196°C, −196 K, or −80°C to catch rushed candidates.
Treating National Parks as Allowing Grazing	In **National Parks, zero human exploitation, private ownership, forestry, or grazing is permitted**. Regulated grazing is permitted only in **Wildlife Sanctuaries**.	NTA match-the-following questions swap the permitted human activities between National Parks and Wildlife Sanctuaries.
Misunderstanding Log Slope Z Values	For normal regional areas, $Z = 0.1$ to 0.2. For entire continents, $Z = 0.6$ to 1.2. For frugivorous birds/mammals, $Z = 1.15$.	Students reverse the numbers, thinking small areas have steep slopes. The larger and more diverse the continental area, the steeper the regression slope.
Assuming Ex-situ Means Releasing into Wild	Ex-situ means keeping endangered organisms **outside their natural home under captive human care** (zoos, gene banks). Release into the wild is a secondary post-breeding goal.	Questions test the fundamental definition: on-site (in-situ) vs off-site (ex-situ).

The Whole Chapter on One Page




Figure 3: Figure 3: Master synthesis mindmap — high-yield integrated architecture of Biodiversity and Conservation covering organizational levels, global patterns, Humboldt's law, The Evil Quartet, and conservation toolkits.

Practice Questions — Part 1 (Q1 to Q5)

Q1. The term "Biodiversity" was popularized in 1992 by which of the following scientists to describe the combined diversity at all levels of biological organization?

(A) Alexander von Humboldt (B) Edward Wilson
(C) Robert May (D) Paul Ehrlich

Sol. Q1. The term **Biodiversity** was popularized in 1992 by the sociobiologist **Edward Wilson** to describe the combined diversity at all levels of biological organization (genetic, species, and ecological). Alexander von Humboldt formulated the species-area relationship, Robert May estimated global species richness (7 million), and Paul Ehrlich proposed the Rivet Popper hypothesis. **Answer: (B)**

Q2. In the medicinal plant *Rauvolfia vomitoria* growing across different Himalayan ranges, the variation observed in the potency and concentration of the active chemical reserpine is an example of which level of biodiversity?

(A) Species diversity (B) Ecological diversity
(C) Genetic diversity (D) Community diversity

Sol. Q2. Variation in the **potency and concentration of reserpine** across different Himalayan populations of *Rauvolfia vomitoria* is a classic NCERT example of **Genetic Diversity** (variation occurring **within a single species** over its geographical distribution range). Other examples include India's >50,000 distinct rice strains and >1,000 mango varieties. **Answer: (C)**

Q3. According to ecologist Robert May's scientifically sound and conservative global estimates, the total number of living species on Earth is approximately:

(A) 1.5 million (B) 20 to 50 million
(C) 7 million (D) 100 million

Sol. Q3. Robert May placed the global species diversity at approximately **7 million species**. Currently, slightly more than 1.5 million species have been formally described (IUCN 2004), meaning that only **22%** of all living species on Earth have been cataloged so far, while >78% remain undiscovered. **Answer: (C)**

Q4. Regarding global taxonomic proportions of documented species, which of the following statements is scientifically accurate according to NCERT?

(A) Plants comprise more than 70% of all recorded living species

(B) Insects constitute more than 70% of all animal species, meaning 7 out of every 10 animals on Earth are insects

(C) The number of mammalian species in the world exceeds the total number of fungi species

(D) Amphibian species diversity in the Eastern Ghats is significantly higher than in the Western Ghats

Sol. Q4. According to official NCERT data: (1) Animals constitute >70% of all recorded species, while plants make up 22%; (2) **Insects constitute more than 70% of all animal species** (7 out of 10 animals are insects); (3) The number of fungi species exceeds the combined total of fishes, amphibians, reptiles, and mammals; and (4) Amphibian diversity is higher in the Western Ghats than in the Eastern Ghats. Thus, option B is correct.
Answer: (B)

Q5. Colombia situated near the equator harbors nearly 1,400 species of birds, whereas Greenland at 71° N harbors only 56 species. This pattern illustrates which ecological principle?

(A) Species-Area Relationship (B) Latitudinal Diversity Gradient
(C) Rivet Popper Hypothesis (D) Biomagnification Gradient

Sol. Q5. The progressive decline in species richness from the equator toward the poles (Colombia with 1,400 bird species vs New York with 105 vs Greenland with 56) exemplifies the **Latitudinal Diversity Gradient**. The tropics harbor maximum diversity due to longer evolutionary speciation time without glaciations, predictable stable climates promoting niche specialization, and higher radiant solar energy. **Answer: (B)**

Practice Questions — Part 2 (Q6 to Q10)

Q6. In Alexander von Humboldt's species-area relationship represented by the equation $\log S = \log C + Z \log A$, what is the measured value of the regression coefficient (Z) when analyzing fruit-eating (frugivorous) birds and mammals in tropical forests across different continents?

(A) 0.1 to 0.2 **(B)** 0.6 to 1.2
(C) 1.15 **(D)** 2.05

Sol. Q6. In the logarithmic species-area equation ($\log S = \log C + Z \log A$), the slope Z is:

- **0.1 to 0.2** for small to regional areas (plants in Britain, birds in California).
- **0.6 to 1.2** for entire continental land masses.
- **1.15** for **frugivorous birds and mammals in tropical forests across different continents**.

Option C is the exact NCERT-stipulated value. **Answer: (C)**

Q7. David Tilman's pioneering long-term outdoor ecological plot experiments in Minnesota grasslands demonstrated that:

(A) Plots with fewer species produced substantially higher annual biomass than diverse plots
(B) Plots with more species showed less year-to-year variation in total biomass and contributed to higher productivity
(C) Species richness is completely independent of ecosystem stability and resilience
(D) Keystone predators invariably cause competitive exclusion and collapse of primary productivity

Sol. Q7. David Tilman's long-term outdoor grassland plot experiments provided conclusive empirical evidence that: (1) Plots with **greater species richness showed less year-to-year variation in total biomass**; and (2) Increased biodiversity directly contributed to **higher overall productivity**. **Answer: (B)**

Q8. In Paul Ehrlich's "Rivet Popper" hypothesis comparing an ecosystem to an airplane, the removal of "rivets from the wings" symbolizes:

(A) Loss of an abundant primary producer with high reproductive turnover
(B) Loss of a keystone species that drives critical, major ecosystem functions
(C) Introduction of an exotic alien herbivore into an aquatic food web
(D) Extinction of a secondary parasite that has multiple alternative hosts

Sol. Q8. In Paul Ehrlich's airplane analogy, popping rivets from seats or windows represents the extinction of minor, redundant species. In contrast, popping rivets from the **airplane wings** represents the loss of **keystone species** (species that perform vital, irreplaceable ecosystem functions, such as top predators or key pollinators), which poses an immediate, catastrophic threat to overall flight safety (ecosystem collapse). **Answer: (B)**

Q9. According to the IUCN Red List (2004), how many total species are documented to have gone extinct across the globe in the last 500 years?

(A) 2,000 species **(B)** 784 species
(C) 125,000 species **(D)** 45,000 species

Sol. Q9. The IUCN Red List (2004) documents the verified extinction of exactly **784 species** in the last 500 years. This total comprises **338 vertebrates**, **359 invertebrates**, and **87 plants**. Notable examples include the Dodo (Mauritius), Quagga (Africa), Thylacine (Australia), Steller's sea cow (Russia), and three tiger subspecies (Bali, Javan, Caspian). **Answer: (B)**

Q10. Which of the following historical extinctions was caused primarily by human "over-exploitation" rather than habitat loss or alien species invasion?

(A) Extinction of over 200 cichlid fish species in Lake Victoria
(B) Extinction of Steller's sea cow and Passenger pigeon
(C) Extinction of native plants in Meghalaya sacred groves
(D) Decline of indigenous catfishes due to *Clarias gariepinus*

Sol. Q10. Both **Steller's sea cow** and the **Passenger pigeon** were driven to extinction in the last two centuries purely due to excessive, uncontrolled **human over-exploitation** (hunting for food and commercial trade). In contrast, the cichlid fish extinction in Lake Victoria was caused by an **alien species invasion** (Nile perch), and the decline of indigenous catfishes was caused by the invasive African catfish *Clarias gariepinus*. **Answer: (B)**

Practice Questions — Part 3 (Q11 to Q15)

Q11. The accidental introduction of which invasive weed with imported wheat led to widespread ecological disruption and allergic health hazards across India?

(A) *Eichhornia crassipes* (Water hyacinth)
(B) *Parthenium hysterophorus* (Carrot grass)
(C) *Lantana camara*
(D) *Rauvolfia vomitoria*

Sol. Q11. ***Parthenium hysterophorus*** (Carrot grass) entered India as an accidental contaminant with imported wheat and spread aggressively as an invasive weed across the entire country, displacing native ground flora and causing severe pollen allergies in humans. *Eichhornia* (water hyacinth) is an aquatic invasive weed, and *Lantana* is an exotic terrestrial shrub. **Answer: (B)**

Q12. When a host fish species becomes extinct in an aquatic ecosystem, its unique assemblage of host-specific monogenean and digenean parasites also invariably perishes. This phenomenon is classified under which component of the Evil Quartet?

(A) Habitat Fragmentation
(B) Alien Species Invasions
(C) Over-Exploitation
(D) Co-Extinctions

Sol. Q12. This is a classic manifestation of **Co-Extinction**. When an organism goes extinct, any species obligately linked with it in a host-parasite or mutualistic partnership (such as co-evolved plant-pollinator mutualisms between *Ficus* and fig wasps) inevitably faces extinction because it cannot survive or reproduce on alternative hosts. **Answer: (D)**

Q13. To qualify officially as a global "Biodiversity Hotspot", a geographical region must satisfy which of the following criteria?

(A) It must possess the largest total geographic land area in a continent
(B) It must exhibit exceptionally high species richness and high endemism, while experiencing severe, accelerated habitat loss
(C) It must be completely devoid of human settlements and have zero commercial agriculture
(D) It must harbor only marine aquatic mammals with low reproductive turnover

Sol. Q13. To qualify as a **Biodiversity Hotspot**, a region must meet two strict criteria simultaneously: (1) It must possess **exceptionally high levels of species richness with a high degree of endemism** (species confined exclusively to that area); and (2) It must be under **accelerated, severe threat of habitat loss**. All 34 global hotspots cover under 2% of Earth's land, but protecting them reduces mass extinction rates by 30%. **Answer: (B)**

Q14. In which of the following Indian states are the famous "Khasi and Jaintia Hills" sacred groves located, serving as the last refuges for numerous rare and threatened plants?

(A) Rajasthan
(B) Meghalaya
(C) Madhya Pradesh
(D) Karnataka

Sol. Q14. The **Khasi and Jaintia Hills** sacred groves are located in **Meghalaya**. They represent ancient, tribal-protected forest refugia where human interference is prohibited by traditional religious faith, preserving numerous rare and endangered endemic plant species that have vanished from surrounding landscapes. Other sacred groves are in the Aravalli Hills (Rajasthan), Western Ghats (Karnataka/Maharashtra), and Chanda/Bastar (MP/Chhattisgarh). **Answer: (B)**

Q15. Preserving viable and fertile gametes of threatened animal species at −196°C in liquid nitrogen for indefinitely long periods is termed:

(A) In-situ Micropropagation
(B) Cryopreservation
(C) Stratified Tissue Culture
(D) Sacred Germplasm Storage

Sol. Q15. Storing living gametes, tissues, or embryos at the ultra-low temperature of **−196°C in liquid nitrogen** is termed **Cryopreservation**. This is an advanced **Ex-situ (off-site)** conservation technique that allows viable germplasm to be preserved indefinitely for assisted reproductive technologies like in vitro fertilization (IVF). **Answer: (B)**