OurMocks

2027 CYCLE

MOCK · PRACTICE · CRACK

CUET UG

Biology

Ecosystem

Official Syllabus Unit X · Class 12 · Comprehensive NCERT Theory, Diagrams & 15 Solved MCQs

INSIDE
Concept Notes

INSIDE
Solved Examples

INSIDE
Practice MCQs + Solutions

Aligned to the latest official syllabus · Concept-first explanations

www.ourmocks.com

Syllabus Alignment & Exam Blueprint

Sub-Topic	Official Syllabus Core Coverage (NCERT Class 12 Unit X)	Exam Focus & Question Types
Ecosystem Structure & Productivity	Ecosystem concept, abiotic and biotic components; stratification (vertical layering); primary productivity: Gross Primary Productivity (GPP) and Net Primary Productivity (NPP), $NPP = GPP - R$; secondary productivity (consumer biomass); global annual biosphere NPP (170 billion tons; oceans 55 billion tons vs land 115 billion tons).	Computing NPP from GPP and R; contrasting oceanic vs terrestrial productivity; defining stratification in forests and ponds.
Decomposition & Detritus Food Chain	5 sequential steps of decomposition: fragmentation (earthworm), leaching, catabolism (bacterial/fungal enzymes), humification (humus: dark, amorphous, colloidal, microbial resistant), and mineralization; rate determinants: chemical composition (lignin/chitin vs nitrogen/sugars) and climatic control (temperature and moisture).	Ordering the 5 steps; identifying factors that accelerate or retard decay; physical and biological properties of humus.
Energy Flow & Food Chains	Photosynthetically Active Radiation (PAR, under 50% of solar radiation); 2–10% PAR captured by autotrophs; Lindeman's 10% energy transfer law; trophic levels; Grazing Food Chain (GFC, major in aquatic) vs Detritus Food Chain (DFC, dominant in terrestrial ecosystems); food webs.	Mathematical calculation of energy across 4 trophic levels; identifying reasons for short food chains; comparing GFC vs DFC.
Ecological Pyramids	Pyramid of numbers (upright in grassland, inverted/spindle in tree); pyramid of biomass (upright in forests, inverted in marine/aquatic oceans); pyramid of energy (always upright, never inverted); standing crop (biomass) vs standing state (nutrients); limitations of ecological pyramids.	Explaining marine biomass inversion (phytoplankton turnover); proving energy pyramid universality; naming pyramid limitations.
Ecological Succession & Nutrient Cycles	Primary vs secondary succession; pioneer species, seres, climax community; hydrarch (water to mesic) vs xerarch (rock to mesic); gaseous (carbon: 71% in oceans, 1% atmosphere) vs sedimentary (phosphorus: rock reservoir, zero respiration); Robert Costanza ecosystem valuation (USD 33 trillion/yr).	Arranging seral stages in hydrarch and xerarch; identifying differences between Carbon and Phosphorus cycles; Costanza service percentages.

EXAM PATTERN

CUET UG Scoring Blueprint: Ecosystem accounts for **3 to 4 questions (15 to 20 marks)** in the CUET UG domain paper. NTA questions focus heavily on **productivity calculations** ($NPP = GPP - R$), the **exact 5-step sequence and chemical controls of decomposition**, the **inversion of the marine biomass pyramid**, the **energy capture percentages** (PAR under 50%, 2–10% capture), and **hydrarch/xerarch seral successions**.

Ecosystem Structure, Stratification & Productivity

An **Ecosystem** is a self-sustaining structural and functional unit of nature where living biological organisms (biotic community) interact continuously with each other and with their non-living physical surroundings (abiotic environment). Ecosystems range in scale from a microscopic water droplet or garden pond to a vast tropical forest, ocean, or the entire global **Biosphere**.

Structural Components of an Ecosystem

Every ecosystem functions through four coordinated integrated processes:

1. **Productivity:** The rate of synthesis of organic biomass by autotrophs through the conversion of radiant solar energy.
2. **Decomposition:** The enzymatic breakdown of dead, complex organic detritus into simple inorganic nutrients.
3. **Energy Flow:** The unidirectional transfer of captured energy from producers through successive consumer trophic levels, dissipating as metabolic heat.
4. **Nutrient Cycling:** The biogeochemical recycling of essential bio-elements (C, N, P, S) between the biotic and abiotic reservoirs.

- **Stratification:** The vertical distribution of different species occupying different vertical physical levels (strata) in an ecosystem:
  - In a mature temperate or tropical forest:
    - **Top Canopy Layer:** Formed by giant emergent trees.
    - **Understory Layer:** Composed of smaller trees and tall shrubs.
    - **Undergrowth Layer:** Composed of shade-tolerant herbs and ferns.
    - **Forest Floor:** Covered by creeping mosses, lichens, and leaf litter.
  - In an aquatic pond ecosystem:
    - **Surface Layer:** Dominated by microscopic floating phytoplankton and floating macrophytes.
    - **Mid-Water Column:** Inhabited by free-swimming nekton (fishes, diving beetles).
    - **Benthic Floor Layer:** Inhabited by rooted submerged plants, benthic invertebrates, fungi, and decomposer bacteria.

Primary & Secondary Productivity

Productivity refers to the rate of biomass production per unit area over a specified time interval. It is expressed in units of mass ($\frac{\text{g}}{\text{m}^2 \cdot \text{yr}}$) or energy ($\frac{\text{kcal}}{\text{m}^2 \cdot \text{yr}}$ or $\frac{\text{kJ}}{\text{m}^2 \cdot \text{yr}}$):

1. **Primary Productivity:** The amount of biomass or organic matter produced per unit area over a time period by photosynthetic autotrophs (green plants, phytoplankton):
   - **Gross Primary Productivity (GPP):** The **total rate of synthesis of organic matter** during photosynthesis by producers per unit area per unit time. A significant fraction of GPP is utilized and oxidized by the plant itself for maintenance and metabolic cellular respiration (R).
   - **Net Primary Productivity (NPP):** The net biomass remaining after subtracting respiratory losses (R) from GPP. NPP represents the **actual stored organic biomass available for consumption by heterotrophs** (herbivores and decomposers):
   $$\text{NPP} = \text{GPP} - R$$
2. **Secondary Productivity:** The rate of formation and assimilation of new organic matter by **consumers** (herbivores and carnivores) per unit area per unit time. (Consumers do not synthesize food de novo; they assimilate ingested primary biomass).
3. **Global Biosphere Productivity Facts:**
   - The annual net primary productivity of the **entire biosphere** is estimated at approximately **170 billion tons (dry weight)** of organic matter.
   - **The Oceanic Paradox:** Although oceans cover approximately **70% of the Earth's surface**, their annual primary productivity is only **55 billion tons**! In contrast, terrestrial ecosystems (occupying just 30% of the surface) generate **115 billion tons**.

- **Why are oceans less productive?** Solar light penetrates only the thin euphotic zone (upper 100–200 m); deep waters are completely dark; and crucial mineral nutrients (such as nitrogen, iron, and phosphorus) are scarce in open ocean waters.

OurMocks

Decomposition & Energy Flow

The Five Steps of Decomposition

Decomposition is an oxygen-requiring biological process in which complex dead organic matter (**detritus**) is broken down by decomposers into simple inorganic substances like carbon dioxide, water, and mineral nutrients:

- **Detritus:** The raw material for decomposition, consisting of dead plant remains (fallen leaves, bark, twigs, flowers) and dead animal remains, including carcasses and fecal matter.
- **The Five Coordinated Stages:**
  1. **Fragmentation:** Detritivores (e.g., the **earthworm**) physically chew, pulverize, and break down coarse detritus into smaller particulate fragments. This drastically increases the exposed surface area available for enzymatic action.
  2. **Leaching:** Water-soluble inorganic nutrients dissolve in percolating soil water, seep downwards through the soil horizon, and precipitate as **unavailable mineral salts**.
  3. **Catabolism:** Bacterial and fungal saprophytes secrete extracellular digestive enzymes directly onto the fragmented detritus, enzymatically degrading complex polymers into simpler inorganic compounds.
  4. **Humification:** Leads to the accumulation of a dark-colored, amorphous, colloidal organic substance called **Humus**.
     - **Key NCERT Properties of Humus:** It is **extremely resistant to microbial action**, decomposes at an exceptionally slow rate, possesses high cation-exchange capacity, and acts as a massive **reservoir of plant nutrients**.
  5. **Mineralization:** Humus is degraded slowly by specialized soil microbes, releasing free inorganic mineral ions ($CO_2, H_2O, Ca^{2+}, Mg^{2+}, K^+, NH_4^+, PO_4^{3-}$) into the soil solution for root absorption.

Factors Regulating Decomposition Rate

Decomposition is largely an aerobic process governed by two main determinants:

- **Chemical Composition of Detritus:**
  - Decomposition is **very slow** if detritus is heavily rich in **lignin** and **chitin** (recalcitrant structural polymers).
  - Decomposition is **very rapid** if detritus is rich in **nitrogen** and **water-soluble substances** (such as sugars and amino acids).
- **Climatic Factors (Temperature & Moisture):**
  - **Warm and moist climatic conditions** strongly accelerate microbial enzyme activity, promoting rapid decomposition (e.g., tropical rainforests, where leaf litter disappears within weeks).
  - **Low temperatures** and **anaerobic conditions (waterlogged soils)** inhibit microbial metabolic activity, drastically slowing decomposition and resulting in the accumulation of massive undecomposed organic matter (e.g., peat bogs and coal deposits).

Energy Flow, PAR & The 10% Transfer Law

Energy flow through all ecosystems on Earth is strictly **unidirectional**—originating from the Sun and dissipating irreversibly as heat:

- **Solar Energy & PAR:**
  - Of the total incident solar radiation reaching the Earth's upper atmosphere, **less than 50%** falls within the **Photosynthetically Active Radiation (PAR)** spectrum (wavelengths between 400 nm and 700 nm).
  - Photosynthetic plants and cyanobacteria capture only **2% to 10% of PAR** (which corresponds to just **1% to 5% of total incident solar radiation**). This minuscule fraction sustains all biotic life on Earth!
- **Lindeman's 10% Energy Transfer Law (1942):**

- When green plants are eaten by primary consumers (herbivores), and herbivores are consumed by carnivores, only about **10% of the energy** stored in biomass at one trophic level is transferred and fixed into the biomass of the next trophic level.
- The remaining **90% of energy** is lost:
  - Dissipated as non-recoverable **metabolic heat** during cellular respiration (obeying the Second Law of Thermodynamics).
  - Expended in locomotion, hunting, feeding, and tissue maintenance.
  - Trapped in unconsumed or unassimilated waste materials (feces, bones).
- **Why Food Chains Are Short:** Because 90% of energy is lost at each step, after 3 to 4 trophic transfers, the remaining energy is too small to support another viable breeding population of apex predators.

Food Chains: Grazing (GFC) vs Detritus (DFC)

- **Grazing Food Chain (GFC):**
  - Starts directly with living photosynthetic primary producers:

    Grass (Producers, T1) → Deer (Herbivore, T2) → Tiger (Carnivore, T3)
  - In **aquatic / marine ecosystems**, GFC is the **dominant pathway** of energy flow (phytoplankton → zooplankton → small fish → large fish).
- **Detritus Food Chain (DFC):**
  - Begins with dead organic matter (detritus), which is consumed by decomposers (saprotrophs: fungi and bacteria):

    Dead Leaf Litter (Detritus) → Earthworm (Detritivore) → Sparrow (Carnivore) → Hawk (Apex)
  - In **terrestrial ecosystems**, a **much larger fraction of total energy flows through the DFC** than through the GFC! Dead leaves and wood far exceed the biomass eaten by herbivores alive.
  - DFC and GFC are interconnected: detritivores from DFC are eaten by predators of GFC, forming an interconnected **Food Web**.




Figure 1: Figure 1: Detailed geometric configurations of ecological pyramids — Number (grassland vs tree), Biomass (terrestrial vs inverted marine), and the universally upright Pyramid of Energy.

Ecological Pyramids

An **Ecological Pyramid** is a graphical representation of the trophic structure and function of an ecosystem, with the producer level forming the broad base and successive consumer levels stacked sequentially toward the apex.

The Three Classic Ecological Pyramids

1. **Pyramid of Numbers:**
   - Plots the absolute number of individual organisms present at each successive trophic level per unit area.
   - **Grassland Ecosystem:** Strictly **Upright**. Millions of grass plants (producers, T1) support thousands of grasshoppers (T2), fewer frogs (T3), and very few eagles (T4).
   - **Tree Ecosystem (The Classic Inversion):**
     - A single giant oak or banyan tree (1 producer, T1) supports thousands of herbivorous insects and fruit-eating birds (T2).
     - These birds in turn host tens of thousands of ectoparasitic lice, mites, and ticks (T3).
     - The pyramid of numbers here is **Inverted or Spindle-Shaped**!
2. **Pyramid of Biomass:**
   - Plots the total standing crop dry weight of living organic tissue ($\frac{g}{m^2}$ or $\frac{kg}{ha}$) present at each trophic level at a specific point in time.
   - **Terrestrial Ecosystems (Forests, Grasslands):** Generally **Upright**. The dry biomass of trees and grasses vastly exceeds that of primary herbivores and secondary carnivores.
   - **Marine / Aquatic Ecosystems (The Famous Inverted Pyramid):**
     - The pyramid of biomass in sea and ocean ecosystems is sharply **INVERTED**!
     - The standing crop biomass of microscopic **phytoplankton** (producers, 4 g/m²) is **much smaller** than the biomass of the zooplankton (21 g/m²) and predatory fishes that feed on them!
     - **Why?** Phytoplankton have microscopic size and exceedingly short lifespans (high turnover rate). Their rapid, explosive cell division sustains a massive standing biomass of long-lived zooplankton and fishes!
3. **Pyramid of Energy:**
   - Plots the total rate of energy flow through each trophic level over time ($\frac{kJ}{m^2 \cdot yr}$).
   - **Universal Law:** The pyramid of energy is **ALWAYS UPRIGHT, NEVER INVERTED** in any ecosystem under any circumstances!
   - When energy flows from autotrophs to top carnivores, substantial energy is irreversibly lost as respiratory heat at each successive trophic step (Lindeman's 10% law). Therefore, energy at lower trophic levels is invariably higher than at higher levels.

Standing Crop vs Standing State

- **Standing Crop:** The total amount of **living biological tissue / biomass** (or total count of living organisms) present in a specific trophic level of an ecosystem at a particular time. Measured as fresh weight or **dry weight** (dry weight is far more scientifically accurate because water content fluctuates).
- **Standing State:** The total amount of **non-living inorganic nutrients** (such as carbon, nitrogen, phosphorus, calcium, potassium) present in the soil, water, or sediment of an ecosystem at any given time.

Fundamental Limitations of Ecological Pyramids

While ecological pyramids provide valuable trophic insights, they possess three significant scientific limitations:

1. They assume a **simple, linear food chain** that rarely exists in real nature; they cannot accommodate interconnected **food webs**.
2. They cannot accommodate a single species that operates across **multiple trophic levels simultaneously** (e.g., a sparrow acts as a primary consumer when eating cereal seeds, but acts as a secondary consumer when eating caterpillars).

3. They **completely ignore saprophytes and decomposers**, giving them zero position in ecological pyramids, despite the fact that decomposers process a massive fraction of an ecosystem's total energy!




Figure 2: Figure 2: Material cycling and decomposition pathways — the 5 sequential stages of detrital mineralization and the global dynamics of Carbon (gaseous) and Phosphorus (sedimentary) cycles.

OurMocks

Ecological Succession & Biogeochemical Cycles

Principles of Ecological Succession

Ecological Succession is the orderly, sequential, and predictable process of directional change in species composition, community structure, and physical environment of a given ecological area over time:

- **Pioneer Species:** The initial pioneering community of organisms that first colonize a completely barren, uninhabited physical substrate.
- **Seres (Seral Stages):** The entire sequence of transitional, intermediate communities that successively replace one another in an area during succession.
- **Climax Community:** The final, self-perpetuating terminal community that achieves stable, dynamic equilibrium with the regional macroclimate (e.g., a mature mesic forest). The climax community persists indefinitely unless disrupted by severe external catastrophic disturbances.
- **Two Types of Succession:**
  - **Primary Succession:** Initiates on a sterile, barren substrate that has **never previously hosted life** (e.g., newly cooled volcanic lava, bare granite rock, newly formed sand dunes, deep newly formed reservoir). It is exceptionally slow, requiring hundreds to thousands of years because fertile soil must first be formed through rock weathering and organic matter accumulation.
  - **Secondary Succession:** Initiates in an area where an existing biological community was completely destroyed or cleared by disturbance (e.g., abandoned farmland, burned or clear-cut forest, flooded river plains). Because fertile soil, seeds, and vegetative roots already persist in the substrate, secondary succession proceeds **much faster** than primary succession.

Hydrarch vs Xerarch Succession

Both aquatic (hydrarch) and terrestrial (xerarch) successional sequences steadily progress from extreme hydrological conditions toward the exact same intermediate, moderately moist condition: the **Mesic** condition!

1. **Hydrarch Succession (In Water):**
   - Begins in wet aquatic environments (ponds, shallow lakes):

ytoplankton (Pioneers) → Submerged Plants → Submerged Free-Floating → Reed-Swamp Stage → Marsh-Meadow → Scrub → Climax

   - Silt and organic detritus accumulate on the pond floor, raising the bottom and transforming the deep open-water body into a shallow marsh, and eventually into a terrestrial mesic forest.
2. **Xerarch Succession (On Bare Rock):**
   - Initiates in extremely dry xeric environments (bare dry rock):

Crustose Lichens (Pioneers) → Foliose Lichens → Mosses → Annual Grasses → Perennial Herbs → Shrubs → Climax Forest (Me

   - Pioneer **crustose lichens** secrete organic carbonic acids that corrode and weather hard granite rock, creating tiny pockets of fine mineral dust. **Mosses** arrive next, forming dense moisture-trapping carpets. Their decay builds rich organic humus, paving the way for rooted grasses, shrubs, and finally a dense mesic forest.

Biogeochemical Nutrient Cycling: Carbon vs Phosphorus

Nutrients are never lost from an ecosystem; they cycle indefinitely between living organisms and physical reservoirs:

Parameter	The Carbon Cycle (Gaseous)	The Phosphorus Cycle (Sedimentary)
Primary Reservoir	Gaseous / Oceanic: The **oceans hold 71% of total global carbon** as dissolved bicarbonate; the atmosphere holds about 1%.	Sedimentary: Lithosphere; **phosphate rock minerals** formed over geological epochs act as the principal reservoir.
Input Mechanism	Atmospheric carbon dioxide (CO_2) fixed via plant and phytoplankton photosynthesis (4×10^{13} kg fixed annually).	Weathering and erosion of phosphate rocks release trace soluble phosphates (PO_4^{3-}) into soil solution for root absorption.

Parameter	The Carbon Cycle (Gaseous)	The Phosphorus Cycle (Sedimentary)
Atmospheric Exchange	Substantial gaseous exchange between living organisms and atmosphere via **respiration** (CO_2 release).	Negligible: **Zero respiratory release of phosphorus** occurs into the atmosphere by any living organism.
Rainfall Inputs	Rainfall dissolves significant atmospheric carbon dioxide.	Atmospheric inputs of phosphorus via rainfall are completely negligible.

Ecosystem Services & Robert Costanza's Valuation

Healthy ecosystems provide indispensable natural purification and life-support functions termed **Ecosystem Services**:

- Purifying atmospheric air and water, mitigating floods and droughts, generating fertile soil, cycling nutrients, pollinating agricultural crops, providing wildlife habitat, and sustaining spiritual and aesthetic values.
- **Robert Costanza's Valuation (1997):**
  - Ecologist Robert Costanza and colleagues attempted to put an economic price tag on nature's life-support services.
  - They calculated that global ecosystem services have an average value of **USD 33 trillion per year**! (By comparison, the total Global Gross National Product (GNP) at that time was only USD 18 trillion per year).
  - **Breakdown of Service Costs:**
    - **Soil Formation:** Accounts for approximately **50%** of the total valuation (USD 16.5 trillion).
    - **Recreation:** Accounts for about **10%**.
    - **Nutrient Cycling & Climate Regulation:** Account for roughly **6% each**.

OurMocks

Key Facts & Terms Cheatsheet

MASTER DIAGNOSTIC & CONCEPTUAL CHEATSHEET — ECOSYSTEM

Key Subject Pointer	Core NCERT Mechanism / Definition	Exam Identifier
Stratification	Vertical layering of diverse species occupying distinct vertical strata.	Forest / Pond Layers
NPP Equation	Net primary productivity: organic matter stored after respiratory losses.	$NPP = GPP - R$
Global Biosphere NPP	170 billion tons total: land = 115 billion tons, oceans = 55 billion tons.	170 Billion Tons
Decomposition Steps	Fragmentation → Leaching → Catabolism → Humification → Mineralization.	5 Fixed Steps
Humus Properties	Dark, amorphous, colloidal substance; highly resistant to microbial decay.	Nutrient Reservoir
Decomposition Inhibitors	High lignin/chitin content, low temperature, and anaerobic conditions.	Lignin / Anaerobiosis
Decomposition Promoters	High nitrogen and water-soluble sugar content, warm and moist climate.	Nitrogen / Moisture
PAR Radiation	Photosynthetically Active Radiation (400–700 nm); under 50% of incident sunlight.	PAR (under 50%)
Solar Energy Captured	Plants capture only 2–10% of PAR (1–5% of total incident sunlight).	2–10% of PAR
Lindeman's 10% Law	Only 10% of energy is transferred and fixed into next trophic level biomass.	10% Energy Transfer
GFC vs DFC	GFC dominates in oceans; DFC channels most energy in terrestrial forests.	DFC > GFC on Land
Tree Pyramid of Numbers	A single tree supports thousands of insects; pyramid is inverted/spindle.	Inverted / Spindle
Ocean Biomass Pyramid	Phytoplankton standing crop is smaller than zooplankton/fishes; inverted.	Inverted in Oceans
Energy Pyramid Law	ALWAYS upright, never inverted in any ecosystem under any conditions.	Always Upright
Standing Crop	Total dry biomass of living organisms in a trophic level at a given time.	Dry Biomass Stock
Standing State	Amount of inorganic non-living nutrients present in soil at any given time.	Soil Nutrients
Hydrarch Succession	Aquatic succession starting from phytoplankton, ending in mesic forest.	Water to Mesic
Xerarch Succession	Terrestrial succession starting from crustose lichens, ending in mesic forest.	Rock to Mesic
Ocean Carbon Reservoir	Oceans hold 71% of global dissolved carbon; atmosphere holds 1%.	71% in Oceans
Phosphorus Cycle	Sedimentary reservoir (rocks); weathering releases PO_4^{3-}; no respiration.	Sedimentary Rock
Robert Costanza Valuation	USD 33 trillion/year for global ecosystem services; soil formation = 50%.	USD 33 Trillion/yr

Common Errors and Special Cases

HIGH-YIELD CUET PITFALLS & DIAGNOSTIC DISTINCTIONS

Common Student Mistake	Scientifically Verified Reality	The Exact Trap Mechanism
Subtracting Decomposition in NPP	$NPP = GPP - R$. Only **respiration (R)** is subtracted; decomposition is an external recycling process.	NTA options present formulas subtracting decomposition or herbivory from GPP to trap students.
Assuming Oceanic Productivity is Highest	Oceans cover 70% of Earth but generate only **55 billion tons** of NPP (vs land generating **115 billion tons**).	Students assume larger surface area guarantees higher productivity; open ocean waters are nutrient-poor.
Omitting Leaching in Decomposition	**Leaching is step 2**, where water-soluble inorganic nutrients seep into soil and precipitate as unavailable salts.	Students remember fragmentation and catabolism but completely forget the physical leaching step.
Believing Energy Pyramids Can Invert	The pyramid of energy is **ALWAYS UPRIGHT, NEVER INVERTED** due to inevitable thermodynamic heat loss.	Questions describe massive apex predators (whales/lions) to entice students into choosing an inverted energy pyramid.
Confusing Standing Crop with Standing State	**Standing crop** is living biological biomass; **Standing state** is non-living inorganic mineral nutrients in soil.	Exams swap the two terms in matching questions; memorize that "crop" is living plants/animals, "state" is soil chemistry.
Claiming DFC is Insignificant on Land	In terrestrial ecosystems, **far more energy flows through DFC than GFC** because most plant biomass dies unconsumed.	Students assume herbivores eat most vegetation; in reality, leaf litter and dead wood dominate terrestrial energy.
Thinking Succession Culminates in Dry or Wet State	Both hydrarch and xerarch successions progress towards the exact same climax: intermediate **mesic** condition!	Students think hydrarch ends in a swamp and xerarch ends in a desert; both end in mesic climax forests.
Asserting Phosphorus Has Gaseous Exchange	Phosphorus has **zero respiratory release** into the atmosphere; it is strictly a **sedimentary** cycle.	Exam questions ask which cycle releases gas via respiration; Carbon does, Phosphorus never does.

The Whole Chapter on One Page




Figure 3: Master synthesis mindmap — high-yield integrated architecture of Ecosystem covering productivity, decomposition, energy flow, pyramids, succession, and biogeochemical cycles.

OurMocks

Practice Questions — Part 1 (Q1 to Q5)

Q1. In an undisturbed tropical grassland ecosystem, the primary producers synthesize organic biomass at a Gross Primary Productivity (GPP) of $25,000 \frac{\text{kJ}}{\text{m}^2 \cdot \text{yr}}$. If autotrophic respiratory losses (R) consume $9,000 \frac{\text{kJ}}{\text{m}^2 \cdot \text{yr}}$, what is the Net Primary Productivity (NPP) available to primary consumers?

(A) $34,000 \frac{\text{kJ}}{\text{m}^2 \cdot \text{yr}}$ **(B)** $16,000 \frac{\text{kJ}}{\text{m}^2 \cdot \text{yr}}$
(C) $2,777 \frac{\text{kJ}}{\text{m}^2 \cdot \text{yr}}$ **(D)** $9,000 \frac{\text{kJ}}{\text{m}^2 \cdot \text{yr}}$

Sol. Q1. Net Primary Productivity (NPP) represents the stored chemical energy remaining after subtracting the plant's metabolic respiratory losses (R) from the Gross Primary Productivity (GPP):

$$\text{NPP} = \text{GPP} - R = 25,000 - 9,000 = 16,000 \frac{\text{kJ}}{\text{m}^2 \cdot \text{yr}}$$

Option B is the exact mathematically verified value. **Answer: (B)**

Q2. Regarding the annual primary productivity of the entire biosphere, which statement accurately reflects global ecological data as documented in NCERT Class 12 Biology?

(A) The total annual net primary productivity of the biosphere is 55 billion tons, with oceans contributing 70%
(B) The total annual net primary productivity of the biosphere is 170 billion tons, with oceans contributing only 55 billion tons
(C) Oceans produce 115 billion tons of dry biomass annually due to high phytoplankton density across open seas
(D) Terrestrial ecosystems contribute only 55 billion tons of dry biomass because deserts and ice cover vast land areas

Sol. Q2. According to NCERT, the annual net primary productivity of the **entire biosphere is approximately 170 billion tons (dry weight)** of organic matter. Despite covering approximately **70% of the Earth's surface**, the **oceans contribute only 55 billion tons**, while terrestrial ecosystems produce the remaining **115 billion tons**. **Answer: (B)**

Q3. During the biological decomposition of dead plant detritus in soil, what is the correct chronological sequence of the five major stages?

(A) Fragmentation → Catabolism → Leaching → Mineralization → Humification
(B) Fragmentation → Leaching → Catabolism → Humification → Mineralization
(C) Leaching → Fragmentation → Humification → Catabolism → Mineralization
(D) Catabolism → Fragmentation → Leaching → Mineralization → Humification

Sol. Q3. The official NCERT sequence of decomposition consists of five coordinated steps: (1) **Fragmentation** (detritivores pulverize detritus); (2) **Leaching** (water-soluble nutrients percolate into soil); (3) **Catabolism** (fungal and bacterial enzymes degrade detritus); (4) **Humification** (accumulation of dark amorphous humus); and (5) **Mineralization** (microbial release of inorganic ions). **Answer: (B)**

Q4. The rate of decomposition of detritus is heavily regulated by its chemical composition and ambient climatic factors. Decomposition proceeds most rapidly when the detritus is:

(A) Rich in lignin and chitin, and the soil environment is cold and waterlogged
(B) Rich in nitrogen and water-soluble substances like sugars, under warm and moist climatic conditions
(C) Rich in cellulose and tannins, under highly anaerobic conditions
(D) Rich in silica and suberin, under arid desert conditions with freezing night temperatures

Sol. Q4. Decomposition rate is accelerated when detritus is chemically **rich in nitrogen and water-soluble substances (like sugars)**, and the climate is **warm and moist** (which optimizes microbial and fungal enzymatic activity). Conversely, detritus rich in **lignin and chitin** decomposes very slowly, and cold or anaerobic conditions drastically inhibit decomposition. **Answer: (B)**

Q5. What proportion of total incident solar radiation falls within the Photosynthetically Active Radiation (PAR) spectrum, and what percentage of PAR is actually captured and fixed by autotrophic plants?

(A) PAR constitutes over 80% of incident radiation; plants capture 50% of PAR
(B) PAR constitutes under 50% of incident solar radiation; plants capture 2% to 10% of PAR
(C) PAR constitutes 50% of incident radiation; plants capture 100% of PAR
(D) PAR constitutes 10% of incident radiation; plants capture 1% to 5% of PAR

Sol. Q5. Of the total incident solar radiation, **less than 50%** falls within the **Photosynthetically Active Radiation (PAR)** spectrum (400–700 nm). Furthermore, photosynthetic plants capture only **2% to 10% of this PAR** (which amounts to just **1% to 5% of total incident solar radiation**), which sustains the entire living world. **Answer: (B)**

Practice Questions — Part 2 (Q6 to Q10)

Q6. If 100,000 Joules of radiant solar energy strikes a forest canopy, and plants capture 1% of this total incident energy into net primary producer biomass, how much energy will reach the tertiary consumer level under Lindeman's 10% Law?

(A) 10 Joules
(B) 1 Joule
(C) 100 Joules
(D) 0.1 Joule

Sol. Q6. Calculating stepwise under Lindeman's 10% Law:

1. Total incident solar energy = $100,000$ J.
2. Captured by Producers (T1) at 1% = $100,000 \times 0.01 = 1,000$ J.
3. Primary Consumers (Herbivores, T2) receive 10% of T1 = $1,000 \times 0.10 = 100$ J.
4. Secondary Consumers (Carnivores, T3) receive 10% of T2 = $100 \times 0.10 = 10$ J.
5. **Tertiary Consumers (Apex, T4) receive 10% of T3 = $10 \times 0.10 = 1$ Jue (1 J).**

Option B is the exact mathematically verified value. **Answer: (B)**

Q7. Regarding the major pathways of energy flow in aquatic versus terrestrial ecosystems, which of the following statements is ecologically accurate?

(A) In oceans, the Detritus Food Chain (DFC) channels 95% of total energy flow
(B) In terrestrial ecosystems, a much larger fraction of energy flows through the Detritus Food Chain (DFC) than through the Grazing Food Chain (GFC)
(C) In all ecosystems, the Grazing Food Chain (GFC) invariably channels over 90% of all community energy
(D) The Detritus Food Chain operates exclusively in aquatic ecosystems and is absent on land

Sol. Q7. In aquatic ecosystems, the Grazing Food Chain (GFC) is the major conduit for energy flow. In **terrestrial ecosystems**, however, a **much larger fraction of energy flows through the Detritus Food Chain (DFC) than through the GFC**, because a vast majority of forest and grassland vegetative biomass (wood, fallen leaves, roots) dies and enters the detrital pool rather than being consumed alive by herbivores. **Answer: (B)**

Q8. Why is the ecological pyramid of biomass in open sea and marine ecosystems inverted?

(A) Marine herbivores are homeothermic and lose 99% of consumed energy as respiratory heat
(B) The standing crop biomass of microscopic phytoplankton is small at any given time, but their high turnover rate supports a larger biomass of zooplankton and fishes
(C) Marine secondary carnivores synthesize their own organic food through chemosynthesis
(D) Phytoplankton possess thick silica frustules that prevent them from being weighed accurately

Sol. Q8. The pyramid of biomass in oceans is **inverted** because the standing crop biomass of **phytoplankton** (producers) at any single moment is very small compared to the biomass of **zooplankton and fishes**. Phytoplankton have microscopic size and very short lifespans (high turnover rate), reproducing rapidly enough to continuously sustain a substantially larger standing biomass of long-lived consumers. **Answer: (B)**

Q9. Under which of the following environmental conditions can an ecological pyramid of energy become inverted?

(A) In a deep marine ecosystem dominated by apex killer whales
(B) In an ancient temperate oak forest hosting millions of hyper-parasites
(C) In an abandoned agricultural field undergoing secondary succession
(D) Under no circumstances whatsoever; a pyramid of energy is always upright in all ecosystems

Sol. Q9. The pyramid of energy is **ALWAYS UPRIGHT, NEVER INVERTED** in any ecosystem under any circumstances. In accordance with the Second Law of Thermodynamics, energy is irreversibly lost as metabolic heat at every successive trophic transfer (Lindeman's 10% law). Consequently, energy available at lower trophic levels is invariably greater than at higher trophic levels. **Answer: (D)**

Q10. During primary hydrarch succession initiating in a deep freshwater lake, which seral community precedes the reed-swamp stage?

(A) Marsh-meadow stage
(B) Submerged free-floating plant stage
(C) Scrub stage
(D) Climax mesic forest stage

Sol. Q10. The official NCERT sequence of seral stages in hydrarch succession is:

on (Pioneers) → Submerged plant stage → Submerged free-floating stage → Reed-swamp stage → Marsh-meadow stage → Scrub stage → For

Thus, the **submerged free-floating plant stage** directly precedes the reed-swamp stage. **Answer: (B)**

Practice Questions — Part 3 (Q11 to Q15)

Q11. Primary xerarch succession begins on bare, dry rock surfaces. The pioneer species that first colonize this extreme habitat and initiate soil formation are:

(A) Foliose and fruticose lichens
(B) Crustose lichens
(C) Perennial bryophyte mosses
(D) Annual xerophytic grasses

Sol. Q11. In xerarch succession on bare rock, the pioneer species are **crustose lichens** (e.g., *Rhizocarpon*). They adhere tightly to bare granite and secrete carbonic and organic acids that corrode and weather the rock surface, creating fine mineral particles that enable foliose lichens and mosses to establish later. **Answer: (B)**

Q12. A fundamental ecological difference between the biogeochemical Carbon cycle and the Phosphorus cycle is that:

(A) Phosphorus possesses a major gaseous atmospheric reservoir, whereas carbon does not
(B) Atmospheric inputs of phosphorus via rainfall are substantial compared to carbon inputs
(C) There is zero respiratory release of phosphorus into the atmosphere, unlike carbon which is respired as CO_2
(D) Phosphorus is cycled exclusively by primary producers without any decomposer involvement

Sol. Q12. The **Phosphorus cycle is a sedimentary cycle** whose reservoir lies in rock minerals. Unlike the Carbon cycle (where respiration releases massive gaseous CO_2 into the atmosphere), there is **zero respiratory release of phosphorus into the atmosphere** by any organism. Furthermore, rainfall inputs of phosphorus are negligible, and oceans hold 71% of global carbon. **Answer: (C)**

Q13. The term "Standing State" in ecology refers specifically to:

(A) The total fresh weight of living autotrophs in a forest at a given time
(B) The amount of inorganic non-living nutrients (such as carbon, nitrogen, phosphorus) present in the soil at any given time
(C) The number of primary consumers supported by a single mature tree
(D) The total annual rate of biomass production by decomposers in a wetland

Sol. Q13. **Standing State** refers to the total amount of **inorganic non-living nutrients** (such as nitrogen, phosphorus, calcium, and potassium) present in the soil or sediment of an ecosystem at any given time. In contrast, **Standing Crop** refers to the living biological biomass present at a trophic level. **Answer: (B)**

Q14. Ecological pyramids suffer from several inherent biological limitations. Which of the following is a recognized limitation of ecological pyramids as highlighted in NCERT?

(A) They can only accommodate aquatic marine food webs and fail on land
(B) They completely exclude saprophytes and decomposers, despite their critical functional role in ecosystems
(C) They assume an inverted pyramid of energy for all terrestrial carnivores
(D) They cannot calculate primary productivity in units of energy

Sol. Q14. Ecological pyramids suffer from three major limitations: (1) They assume a simple food chain and cannot accommodate **food webs**; (2) They cannot handle species belonging to **multiple trophic levels** (like omnivores); and (3) **They completely exclude decomposers and saprophytes**, giving them no position in the pyramid despite their vital functional role in ecosystems. **Answer: (B)**

Q15. In 1997, ecologist Robert Costanza and his colleagues estimated the economic value of fundamental global ecosystem services at an average of USD 33 trillion per year. In their valuation, which single ecosystem service accounted for approximately 50% of this total cost?

(A) Recreation and ecotourism
(B) Climate regulation
(C) Soil formation
(D) Nutrient cycling

Sol. Q15. In Robert Costanza's landmark valuation of nature's life-support services (USD 33 trillion annually), **Soil Formation** accounted for approximately **50%** of the total cost (USD 16.5 trillion). Recreation accounted for 10%, and climate regulation and nutrient cycling accounted for 6% each. **Answer: (C)**