OurMocks

2027 CYCLE

MOCK · PRACTICE · CRACK

CUET UG

Biology

Microbes in Human Welfare

Official Syllabus Unit VIII · Class 12 · CUET UG 2027 Cycle Notes

INSIDE
Concept Notes

INSIDE
Solved Examples

INSIDE
Practice MCQs + Solutions

Aligned to the latest official syllabus · Concept-first explanations

www.ourmocks.com

Syllabus Alignment & Exam Blueprint

Sub-Topic	Official NTA CUET UG Syllabus Scope	High-Yield Exam Focus
Household Food Processing	Microbes in household products: Curd, dough, bread, traditional foods, cheese varieties.	LAB (Vitamin B_{12} boost); *S. cerevisiae* (CO_2 puffing); Swiss cheese (*P. sharmanii*) vs Roquefort (*P. roqueforti*).
Industrial Fermentation	Microbes in industrial products: Fermented beverages, distillation, commercial organic acids, and enzymes.	Brewer's yeast; distilled (whisky/rum) vs non-distilled (wine/beer); *A. niger* (citric), *A. aceti* (acetic); lipases & pectinases.
Bioactive Molecules & Drugs	Antibiotics (penicillin discovery) and bioactive molecules (streptokinase, cyclosporin A, statins).	Streptokinase (clot buster); Cyclosporin A (*T. polysporum*, immunosuppressant); Statins (*M. purpureus*, HMG-CoA reductase).
Sewage Treatment (STP)	Microbes in sewage treatment: Primary (physical) vs Secondary (biological) treatment; BOD dynamics.	Sequential filtration & grit settling; Aeration tank & **Flocs**; **BOD** metric; Activated sludge & river action plans (GAP/YAP).
Biogas Generation	Microbes in production of biogas; methanogenic bacteria, cattle rumen, and biogas plant design.	*Methanobacterium* (anaerobic); Biogas mix ($CH_4 + CO_2 + H_2S$); Rumen cellulose breakdown; IARI & KVIC plant.
Biocontrol Agents	Microbes as biocontrol agents: Natural predators, Bt endotoxin, and baculoviruses in IPM.	Ladybird (aphids); Dragonfly (mosquitoes); *Bacillus thuringiensis* (Bt cry toxin); Baculoviruses (NPV narrow-spectrum).
Biofertilisers	Microbes as biofertilisers: Symbiotic/free-living nitrogen fixers, mycorrhiza, and cyanobacteria.	*Rhizobium* vs *Azotobacter*/*Azospirillum*; Mycorrhiza (*Glomus* phosphorus uptake); Cyanobacteria (*Anabaena*, *Nostoc*) in paddy.

CUET HIGH-YIELD EXAM ARCHITECTURE & WEIGHTAGE GUIDE

Microbes in Human Welfare is one of the most reliable and high-scoring memory-based chapters in CUET UG Biology, yielding **3 to 5 compulsory questions (12–20 marks)** per test session. Questions focus strictly on precise two-part matching pairs (Organism → Product/Function): (1) Household fermentation and cheese makers (*P. sharmanii* vs *P. roqueforti*); (2) Organic acids and their source fungal/bacterial species; (3) The "Holy Trinity" of bioactive molecules: Streptokinase (clot buster), Cyclosporin A (immunosuppressant), and Statins (HMG-CoA reductase inhibitor); (4) Sewage treatment sequence, flocs composition, and BOD relationships; and (5) Specific ecological attributes of Baculoviruses (NPV) and Mycorrhiza (*Glomus*).

Microbes in Household Food Processing

Microorganisms (bacteria, fungi, protozoans, microscopic viruses) are ubiquitous in the biosphere, inhabiting soil, water, air, thermal vents up to 100°C, acidic sulfataras, and deep brine deposits. While some microbes cause infectious disease, the vast majority perform indispensable ecological and industrial metabolic conversions.

Lactic Acid Bacteria (LAB) & Dairy Fermentation

- **The Curdling Process:** Milk is converted into curd through the inoculation of a small starter culture (inoculum) containing millions of active **Lactic Acid Bacteria (LAB)**, primarily *Lactobacillus* and related species.
- **Mechanism:** At ambient temperatures, LAB multiply rapidly in milk, fermenting the milk sugar (lactose) into **lactic acid**. The accumulating lactic acid coagulates and partially digests the primary milk protein (**casein**).
- **Nutritional & Health Benefits:**
  - **Vitamin Enrichment:** LAB fermentation significantly improves the nutritional quality of curd by markedly increasing its **Vitamin B_{12}** content.
  - **Gut Immunity:** Within the human gastrointestinal tract, thriving colonies of LAB play a crucial protective role by inhibiting the proliferation of pathogenic, disease-causing microbes.

Fermentation in Traditional Foods & Baking

- **Dough for Dosa and Idli:** The batter for traditional Indian foods like dosa and idli is fermented by bacteria. The characteristic **puffed-up appearance** of the rising dough is caused by the entrapment and effervescence of **carbon dioxide (CO_2)** gas produced during anaerobic fermentation.
- **Bread Making (Baker's Yeast):** The dough used for leavening bread is fermented by the common unicellular yeast ***Saccharomyces cerevisiae***, designated industrially as **Baker's yeast**. Rapid alcoholic fermentation releases CO_2, creating thousands of tiny air pockets that render the baked loaf soft, spongy, and porous.
- **Traditional Beverages & Foods:**
  - **Toddy:** A traditional beverage of parts of southern India, prepared by tapping and fermenting the sugary sap of palms using wild yeasts.
  - **Fermented Substrates:** Microbes are utilized to ferment soybeans (yielding tempeh, miso, and soy sauce), tender bamboo shoots, and fish into protein-rich dietary staples across Asian cuisines.

Cheese Processing & Ripening

Cheese is one of the oldest dairy products preserved by microbial fermentation. Different varieties of cheese are characterized by their unique texture, flavor, and taste, governed by the specific microbial species used in ripening:

- **Swiss Cheese:** Characterized by large, conspicuous holes throughout the cheese body. These holes are formed by the slow, continuous production of large volumes of **carbon dioxide (CO_2)** gas by the bacterium ***Propionibacterium sharmanii***.
- **Roquefort Cheese:** A semi-hard blue cheese ripened by inoculating specific spores of the filamentous fungus ***Penicillium roqueforti***, which grow through the curd to impart its distinctive marbling, sharp aroma, and piquant flavor.
- **Camembert Cheese:** Ripened on its outer surface by the mold *Penicillium camemberti*.

Industrial Fermentation & Commercial Products

On an industrial scale, microbes are cultivated in massive vessels called **fermenters** (bioreactors) ranging from thousands to hundreds of thousands of liters under strictly controlled temperature, pH, aeration, and nutrient parameters.

Fermented Beverages

- **Brewer's Yeast:** The identical yeast species used in baking, ***Saccharomyces cerevisiae***, is designated as **Brewer's yeast** when utilized for industrial alcohol production. It ferments malted cereals (barley, wheat) and crushed fruit juices into **ethanol** and CO_2.
- **Distilled vs Non-Distilled Beverages:**
  - **Non-Distilled Beverages:** Produced without post-fermentation distillation, resulting in lower alcohol concentration (limited by yeast ethanol toxicity to 10–15%). Examples: **Wine** (fermented grape juice) and **Beer** (fermented malted barley).
  - **Distilled Alcoholic Beverages:** Produced by fractional distillation of the fermented mash to achieve higher ethanol concentrations (40–50% or more). Examples: **Whisky**, **Brandy**, **Rum**, Gin, and Vodka.

Antibiotics: Discovery & Clinical Triumph

Antibiotics (Greek: *anti* = against, *bios* = life) are chemical substances produced by certain microorganisms that inhibit the growth of or kill other disease-causing microorganisms at low concentrations.

- **The Accidental Discovery: Alexander Fleming** (1928) discovered the first antibiotic, **Penicillin**, while working on the bacterium *Staphylococcus*. He observed that a fungal contaminant—the mold ***Penicillium notatum***—growing on an unwashed culture plate created a clear zone of inhibition where *Staphylococcus* colonies failed to grow. He named the active inhibitory substance Penicillin.
- **Clinical Development:** The full therapeutic potential and industrial isolation of penicillin as an effective antibiotic was established over a decade later by **Ernest Chain** and **Howard Florey**. Penicillin was extensively deployed to treat wounded Allied soldiers during World War II.
- **Nobel Prize:** Alexander Fleming, Ernest Chain, and Howard Florey were jointly awarded the **Nobel Prize in Physiology or Medicine in 1945**.
- **Impact:** Antibiotics transformed human medicine, bringing once-fatal bacterial scourges under control: plague, whooping cough (*kali khansi*), diphtheria (*gal ghotu*), and leprosy (*kusht rog*).




Figure 1: Industrial microbes, bioactive molecules & biocontrol architecture — curated classification of household fermentation, organic acids, industrial enzymes, clot-busters, immunosuppressants, and biofertilisers.

Organic Acids & Industrial Enzymes

Microbes serve as bio-factories for large-scale commercial production of organic acids and industrial enzymes:

Organic Chemical / Enzyme	Microbial Source	Taxonomic Class & Commercial Application
Citric Acid	***Aspergillus niger***	Filamentous Ascomycete fungus; food preservative, flavoring in fizzy drinks, and metal pickling.

Organic Chemical / Enzyme	Microbial Source	Taxonomic Class & Commercial Application
Acetic Acid (Vinegar)	***Acetobacter aceti***	Gram-negative bacterium; food seasoning, condiment, and antimicrobial preservative.
Butyric Acid	***Clostridium butylicum***	Obligate anaerobic bacterium; plasticizer synthesis and chemical intermediate (rancid butter odor).
Lactic Acid	***Lactobacillus*** species	Bacterium; food acidification, curing, and synthesis of biodegradable polylactic acid (PLA) polymers.
Lipases	Various fungi and bacteria	Enzymes formulated into commercial laundry detergents to hydrolyze triacylglycerols and remove **oily stains**.
Pectinases & Proteases	Microbial fermentation broths	Enzymes added to clarify commercial bottled fruit juices, hydrolyzing colloidal pectins to yield crystal-clear juice.

The Sacred Bioactive Molecules (NCERT Holy Trinity)

Three bioactive molecules represent the single highest-frequency testing target in this unit:

1. **Streptokinase ("The Clot Buster"):**
   - **Source Microbe:** Produced by the pathogenic bacterium ***Streptococcus***, subsequently modified by recombinant DNA genetic engineering.
   - **Physiological Action:** Acts as a potent fibrinolytic agent, activating plasminogen into plasmin to enzymatically dissolve intravascular blood clots (thrombi).
   - **Clinical Indication:** Administered intravenously as a **"clot buster"** to emergency patients suffering from **myocardial infarction** (heart attack) to restore coronary blood flow.
2. **Cyclosporin A (The Immunosuppressive Drug):**
   - **Source Microbe:** Extracted from the soil-dwelling filamentous fungus ***Trichoderma polysporum***.
   - **Physiological Action:** Selectively inhibits T-lymphocyte activation and interleukin-2 synthesis.
   - **Clinical Indication:** Used worldwide as an essential **immunosuppressive agent** in patients undergoing **organ transplantation** (kidney, liver, heart) to prevent cell-mediated graft rejection.
3. **Statins (Blood Cholesterol-Lowering Agents):**
   - **Source Microbe:** Produced by the red mold yeast ***Monascus purpureus***.
   - **Mechanism of Action:** Statins function as **competitive inhibitors** of the enzyme **HMG-CoA reductase** (β-hydroxy-β-methylglutaryl-CoA reductase), the rate-limiting enzyme in hepatic cholesterol biosynthesis.
   - **Clinical Indication:** Prescribed to lower elevated blood low-density lipoprotein (LDL) cholesterol and prevent atherosclerotic cardiovascular disease.

Microbes in Sewage Treatment (STP)

Municipal wastewater generated daily in cities and towns—termed **sewage**—contains over 99% water, suspended solids, human fecal excreta, organic waste, and dangerous pathogenic organisms. Untreated sewage cannot be discharged directly into natural water bodies like rivers and streams without causing fatal ecological collapse and water-borne epidemics.

Sewage is rendered safe through multi-stage processing in **Sewage Treatment Plants (STPs)** using natural heterotrophic microbes. Treatment occurs in two fundamental stages: **Primary (Physical)** and **Secondary (Biological)**, followed where required by **Tertiary (Physicochemical)** disinfection.

Primary Treatment (Physical Filtration & Sedimentation)

Primary treatment involves the physical removal of large and small particulate matter from raw sewage through mechanical operations:

1. **Sequential Filtration:** Raw sewage flows through sequential wire-mesh bar screens to filter out large floating debris (plastics, rags, leaves, paper).
2. **Grit Settling:** The screened effluent enters a grit chamber where heavier mineral grit (soil particles, pebbles, sand) settles out under gravity.
3. **Primary Sedimentation:** The sewage enters large settling tanks where suspended solid organic matter settles to the bottom, forming the **Primary Sludge**. The supernatant liquid above the sludge forms the **Primary Effluent**.
4. **Effluent Transfer:** The primary sludge is scraped out, while the primary effluent is pumped forward into the secondary biological treatment units.

Secondary Treatment (Aerobic Biological Degradation)

Secondary treatment is a living biological process that harnesses natural aerobic microorganisms to degrade dissolved organic pollutants:

1. **Aeration Tanks:** The primary effluent is channeled into massive **aeration tanks**. Mechanical agitators constantly churn the fluid, while high-pressure blowers continuously pump air into the tank.
2. **Formation of Flocs:** Continuous aeration promotes the vigorous, rapid proliferation of useful aerobic heterotrophs into mesh-like clusters termed **Flocs**—defined strictly as **masses of bacteria associated with fungal filaments to form mesh-like structures**.
3. **Degradation of Organic Matter:** The active flocs consume the vast majority of dissolved organic compounds in the effluent for their growth and cellular respiration.
4. **Drastic Reduction in BOD:** As the microbes digest the organic waste, the **Biochemical Oxygen Demand (BOD)** of the effluent declines sharply.
5. **Settling Tank & Activated Sludge:** Once the BOD is reduced to acceptable thresholds, the effluent flows into a secondary **settling tank**. In this tranquil tank, the bacterial flocs settle down under gravity, forming the **Activated Sludge**.
6. **Inoculum Recycling:** A small fraction of the activated sludge is pumped back into the primary aeration tank to serve as the active microbial **inoculum** (starter culture).
7. **Anaerobic Digestion:** The remaining major portion of the activated sludge is pumped into sealed, unoxygenated vessels called **Anaerobic Sludge Digesters**.




Figure 2: Sewage Treatment Plant (STP) & Biogas generation dynamics — sequential physical filtration/sedimentation, aerobic flocs and BOD reduction, activated sludge recycling, and anaerobic methanogenesis producing combustible biogas.

The Sacred BOD Metric & Polluting Potential

- **Definition of BOD: Biochemical Oxygen Demand (BOD)** represents the total quantity of dissolved oxygen (in milligrams per liter) that would be consumed if all the biodegradable organic matter in one liter of water were completely oxidized by aerobic bacteria.
- **The Infallible BOD Rule:** The magnitude of BOD directly reflects the concentration of organic matter in the water sample. Therefore:

$$\text{Higher BOD} \iff \text{Higher Organic Pollution} \iff \text{Higher Polluting Potential}$$

$$\text{Lower BOD} \iff \text{Lower Organic Pollution} \iff \text{Cleaner Water}$$

- Secondary biological treatment is maintained continuously until the BOD of the effluent has dropped by 80–90%, making it safe for discharge into rivers.

Anaerobic Digestion, Tertiary Polishing & River Action Plans

- **Anaerobic Sludge Digesters:** In these sealed tanks, strictly **anaerobic bacteria** (including methanogenic species) attack and digest both the bacteria and fungi present in the activated sludge.
- **Biogas Production:** During this anaerobic breakdown, the microbes liberate a mixture of gases comprising primarily **methane (CH_4)**, **carbon dioxide (CO_2)**, and **hydrogen sulfide (H_2S)**. This gas mixture is inflammable and constitutes **Biogas**, which is harnessed as clean fuel for electricity generation or heating.
- **Tertiary Treatment (Physicochemical Polishing):** Where treated water is reclaimed for municipal or industrial reuse, a tertiary polishing step is applied. This involves chemical precipitation of residual phosphates/nitrates followed by disinfection using **chlorine gas**, **ozone**, or **ultraviolet (UV) radiation** to destroy lingering pathogenic viruses and bacterial cysts.
- **Government Initiatives:** Due to exploding urban populations, sewage volumes exceed STP capacities across India. To prevent holy rivers from turning into toxic sewers, the Ministry of Environment and Forests initiated comprehensive national cleanup frameworks:
  - **Ganga Action Plan (GAP)** — Initiated in 1985 to construct sewage diversion canals and STPs along major cities on the Ganga.
  - **Yamuna Action Plan (YAP)** — Launched to remediate sewage inflow into the Yamuna river across Delhi, Haryana, and Uttar Pradesh.

Microbes in Production of Biogas

Biogas is a clean, combustible gaseous fuel produced through the anaerobic biological degradation of biomass (animal manure, crop residues, municipal sludge) by methanogenic microorganisms.

Methanogens & The Three Biochemical Stages

- **Methanogens:** Archaebacteria and bacteria that grow strictly anaerobically on cellulosic materials, converting organic acids and hydrogen into methane. The principal archetype is ***Methanobacterium***.
- **Gas Composition:**
  - **Methane (CH_4):** $50\% - 70\%$ (the primary combustible fuel).
  - **Carbon Dioxide (CO_2):** $30\% - 40\%$.
  - **Trace Gases:** Hydrogen (H_2), Nitrogen (N_2), and Hydrogen Sulfide (H_2S).
- **The Three Stages of Methanogenesis:**
  1. **Stage 1 (Hydrolysis):** Facultative anaerobic microbes hydrolyze complex polymers (cellulose, hemicellulose, proteins, lipids) into simple soluble monomers (glucose, amino acids, fatty acids).
  2. **Stage 2 (Acidogenesis & Acetogenesis):** Acid-forming bacteria convert monomers into volatile organic acids (propionic acid, butyric acid), which acetogenic bacteria convert into **acetic acid**, CO_2, and H_2.
  3. **Stage 3 (Methanogenesis):** Strictly anaerobic methanogens (***Methanobacterium***) reduce acetic acid and $CO_2 + H_2$ into **methane (CH_4)** gas ($CH_3COOH \rightarrow CH_4 + CO_2$ and $CO_2 + 4H_2 \rightarrow CH_4 + 2H_2O$).

- **The Rumen of Cattle:** Methanogens are naturally abundant in the **rumen** (a specialized compartment of the stomach) of ruminant cattle. A significant fraction of ruminant diet consists of cellulosic plant cell walls. Rumen methanogens produce cellulase enzymes that digest cellulose, playing an indispensable role in cattle nutrition.
- **Gobar Gas:** Because cattle dung (locally termed **gobar**) contains large amounts of undigested cellulosic fibers and rich resident populations of methanogens, it serves as the premier substrate for rural biogas generation, universally known as **Gobar Gas**.

Design & Operation of a Biogas Plant

Biogas production technology was pioneered and standardized in India through the dedicated research efforts of the **Indian Agricultural Research Institute (IARI)** and the **Khadi and Village Industries Commission (KVIC)**.

- **Structural Components:**
  1. **Digester Pit:** A cylindrical concrete or masonry tank, typically **10 to 15 feet deep**, sunk into the earth.
  2. **Inlet Tank & Slurry:** Cattle dung is mixed with water in a 1:1 ratio in an inlet chamber to form a uniform **slurry**, which is fed via an inlet pipe into the bottom of the digester tank.
  3. **Floating Gas Holder:** A cylindrical steel or fiberglass cover placed directly over the digesting slurry. As methanogens metabolize the slurry anaerobically, liberating gas, the floating holder **rises upwards**, providing a direct visual indicator of gas volume and pressure.
  4. **Gas Outlet Pipe:** An outlet valve and pipe connected to the gas holder conveys biogas directly to village households for smokeless cooking and illumination.
  5. **Spent Slurry Outlet:** An overflow chamber collects the spent, de-methanated slurry. This residue is periodically removed and dried, serving as exceptionally rich, odorless **organic fertilizer/ manure** rich in nitrogen and phosphorus.

Microbes as Biocontrol Agents & Biofertilisers

Modern industrial agriculture relies heavily on chemical insecticides, pesticides, and synthetic inorganic fertilizers. These toxic agrochemicals poison groundwater tables, destroy beneficial soil microbiota, cause bioaccumulation up trophic food chains, and induce pesticide resistance. Sustainable agriculture utilizes biological organisms as **biocontrol agents** and **biofertilisers**.

Biocontrol of Pests & Plant Diseases

Biological control (biocontrol) is an agro-ecological strategy of managing insect pests and plant pathogens by utilizing natural biological predators and parasites, rather than applying chemical poisons:

- **Natural Predators:**
  - **Ladybird Beetle:** Recognized by its red body with black spots; an efficient natural predator utilized to biological control **aphids** on agricultural crops.
  - **Dragonflies:** Swift aerial predators utilized to eradicate **mosquito** populations around standing water bodies.
- **Bacillus thuringiensis (Bt):**
  - A soil bacterium utilized as a microbial biocontrol agent against caterpillars of butterfly and moth pests.
  - Commercial formulations containing dried bacterial spores are mixed with water and sprayed onto vulnerable crops (brassicas, fruit trees).
  - **Molecular Mechanism:** The bacterium synthesizes crystalline **Cry** protein protoxins. When swallowed by insect larvae, the crystal dissolves only in the **alkaline pH (>9.0) of the insect midgut**. Midgut proteases cleave the protoxin into active endotoxin, which binds to epithelial cell receptors, creating lytic pores that cause gut swelling, epithelial perforation, and larval death. Bt toxin genes have been cloned into plants (Bt cotton).
- **Trichoderma Species:**

- Free-living, soil-dwelling filamentous fungi exceptionally common in the **root rhizosphere ecosystems**.
- They act as effective biocontrol agents against numerous fungal soil-borne plant pathogens through mycoparasitism and antibiotic secretion.

- **Baculoviruses (Genus Nucleopolyhedrovirus - NPV):**
  - Viruses that attack insects and other arthropods.
  - **The IPM Gold Standard:** Baculoviruses are the premier biological control agents in Integrated Pest Management (IPM) programs because they are **species-specific, narrow-spectrum insecticidal agents**.
  - **Zero Non-Target Toxicity:** They have been rigorously proven to have **no negative side-effects** on non-target organisms—including plants, mammals, birds, fish, or beneficial pollinating insects. They are vital when beneficial insects must be preserved in ecologically fragile landscapes.

Microbes as Biofertilisers

Biofertilisers are living organisms that enrich the nutrient and mineral quality of agricultural soil through biological nitrogen fixation, phosphorus solubilization, and organic matter synthesis. The three primary categories are **Bacteria**, **Fungi**, and **Cyanobacteria**.

Category	Representative Microorganisms	Agronomic Mechanism & Benefits
Symbiotic Bacteria	***Rhizobium*** species	Inhabits root nodules of leguminous plants; fixes inert atmospheric N_2 into organic ammonium/nitrate ions, directly nourishing host crops.
Free-Living Bacteria	***Azotobacter*** (aerobic) and ***Azospirillum***	Free-living soil heterotrophs; fix atmospheric nitrogen into the soil matrix without plant association, enriching the general soil nitrogen reserve.
Mycorrhizal Fungi	Genus ***Glomus*** and related fungal families	Forms symbiotic **mycorrhiza** with plant root systems. Fungal hyphae explore vast soil volumes to absorb **phosphorus** and translocate it to the plant. Confers resistance to root pathogens, high tolerance to salinity and drought, and overall growth vigor.
Cyanobacteria (BGA)	***Anabaena***, ***Nostoc***, ***Oscillatoria***	Autotrophic, photosynthetic nitrogen-fixing blue-green algae. In **paddy (rice) fields**, cyanobacteria serve as the principal biofertiliser, dramatically boosting fertility and contributing valuable organic humus to the soil.

- **Cyanobacterial Heterocysts:** Genera like *Anabaena* and *Nostoc* possess specialized, thick-walled, pale cells called **heterocysts**. Heterocysts lack Photosystem II (preventing photosynthetic oxygen generation) and maintain an anaerobic micro-niche to protect the oxygen-sensitive enzyme **nitrogenase** from irreversible inactivation.

Key Facts & Terms Cheatsheet

MASTER DIAGNOSTIC & CONCEPTUAL CHEATSHEET — MICROBES IN HUMAN WELFARE

Key Subject Pointer	Core NCERT Mechanism / Definition	Exam Identifier
Curd Fermentation	Coagulates milk protein casein and boosts Vitamin B_{12} content.	*Lactobacillus* (LAB)
Bread & Beer Yeast	Unicellular ascomycete driving alcoholic fermentation and CO_2 release.	*Saccharomyces cerevisiae*
Swiss Cheese Holes	Large circular cavities formed by CO_2 gas release during fermentation.	*Propionibacterium sharmanii*
Roquefort Ripening	Specific fungal ripening conferring unique texture and marbling.	*Penicillium roqueforti*
First Antibiotic	Extracted from mold inhibiting *Staphylococcus*; developed by Chain/Florey.	Penicillin (*P. notatum*)
Industrial Citric Acid	Aerobic fungal fermentation yielding high titers of commercial citric acid.	*Aspergillus niger*
Vinegar Acid	Bacterial oxidation of ethanol into commercial acetic acid.	*Acetobacter aceti*
Laundry Lipases	Enzymes incorporated into detergents to digest lipid triacylglycerols.	Removes Oily Stains
Juice Clarification	Enzymatic digestion of colloidal fruit pectins in commercial bottling.	Pectinases & Proteases
Myocardial Clot Buster	Genetically modified bacterial enzyme dissolving intravascular fibrin clots.	Streptokinase (*Streptococcus*)
Organ Immunosuppressant	Fungal bioactive peptide preventing graft rejection in transplant patients.	Cyclosporin A (*T. polysporum*)
Cholesterol-Lowering Drug	Competitive inhibitor of HMG-CoA reductase for cholesterol synthesis.	Statins (*Monascus purpureus*)
Primary STP Action	Physical removal of large and small particles via filtration and gravity.	Primary Sludge vs Effluent
STP Aeration Flocs	Mesh-like clusters of aerobic bacteria intertwined with fungal filaments.	Consumes Organic Waste
Pollution Index (BOD)	Oxygen consumed by microbes to oxidize all organic matter in 1 L water.	High BOD = High Pollution
Activated Sludge Role	Sedimented flocs; small part recycled as inoculum, rest digested anaerobically.	Inoculum Recirculation
Biogas Gas Mixture	Combustible fuel generated anaerobically: 50–70% CH_4 + CO_2 + trace H_2S.	*Methanobacterium*
Narrow-Spectrum Biocontrol	Species-specific insecticidal virus with zero non-target ecological toxicity.	Baculovirus (NPV)
Phosphorus Biofertiliser	Fungal-root symbiotic mycorrhiza extracting phosphorus from soil.	Genus *Glomus*
Paddy Field Biofertiliser	Photosynthetic nitrogen-fixing cyanobacteria enriching rice soils.	*Anabaena* & *Nostoc*

Common Errors and Special Cases

HIGH-YIELD CUET PITFALLS & DIAGNOSTIC DISTINCTIONS

Common Student Mistake	Scientifically Verified Reality	The Exact Trap Mechanism
Attributing Swiss Cheese Holes to Yeasts	The holes are formed by the bacterium ***Propionibacterium sharmanii***, NOT yeast.	Students associate all CO_2 bubbles with yeast fermentation; Swiss cheese is strictly bacterial.
Confusing Fleming's and Florey's Roles	**Alexander Fleming** discovered penicillin (1928); **Ernest Chain and Howard Florey** developed it into a clinical drug (1942).	Exam questions deliberately mix discovery with pharmaceutical purification and mass deployment.
Classifying Primary STP as Biological	Primary treatment is purely **physical** (screening and sedimentation); secondary treatment is **biological** (flocs and BOD reduction).	Any question mentioning flocs, microbes, or aeration belongs strictly to secondary treatment.
Assuming BOD Increases in Clean Water	**Lower BOD indicates cleaner water** with less organic pollution. As treatment succeeds, BOD **falls dramatically**.	Students invert the BOD scale; remember that BOD is a measure of waste hunger—less waste means less oxygen demand.
Swapping Bioactive Molecule Sources	Cyclosporin A is from the fungus ***Trichoderma polysporum***; Statins are from the yeast ***Monascus purpureus***; Streptokinase is from the bacterium ***Streptococcus***!	NTA frequently pairs Cyclosporin A with yeast or Statins with bacteria; memorize the kingdom and species of each.
Claiming Baculoviruses Kill All Insects	Baculoviruses are **strictly species-specific and narrow-spectrum**, sparing beneficial pollinators and predators.	Broad-spectrum activity applies to synthetic pesticides; baculoviruses are the gold standard for narrow IPM target safety.
Assigning Nitrogen Fixation to Mycorrhiza	Mycorrhiza (*Glomus*) absorbs **phosphorus and water**; it does **NOT** fix atmospheric nitrogen.	Nitrogen fixation belongs exclusively to diazotrophs (*Rhizobium*, *Azotobacter*, *Azospirillum*, Cyanobacteria).

The Whole Chapter on One Page




Figure 3: Master synthesis mindmap — high-yield integrated architecture of Microbes in Human Welfare covering household foods, industrial fermentation, bioactive drugs, STP operations, biogas, biocontrol, and biofertilisers.

OurMocks

Practice Questions — Part 1 (Q1 to Q5)

Q1. During the conversion of milk into curd, Lactic Acid Bacteria (LAB) induce coagulation and partial digestion of milk proteins while simultaneously enhancing its nutritional quality by significantly increasing the concentration of:

(A) Vitamin C (ascorbic acid) **(B)** Vitamin B_{12} (cyanocobalamin)
(C) Vitamin D (calciferol) **(D)** Vitamin A (retinol)

Sol. Q1. When milk is converted into curd by Lactic Acid Bacteria (LAB), predominantly *Lactobacillus*, the bacteria ferment lactose into lactic acid, which coagulates and partially digests casein milk proteins. In addition to creating curd, this fermentation significantly enhances the nutritional value of the milk by increasing its **Vitamin B_{12}** content. LAB also play a protective role in the human stomach by checking pathogenic microbes. **Answer: (B)**

Q2. The puffed-up, spongy appearance observed during the leavening of bread dough and the fermentation of traditional dosa batter is driven by the metabolic liberation of carbon dioxide gas through the action of:

(A) The anaerobic bacterium *Clostridium butylicum* **(B)** The unicellular ascomycete yeast *Saccharomyces cerevisiae*
(C) The filamentous mold *Penicillium notatum* **(D)** The methanogenic archaeon *Methanobacterium*

Sol. Q2. Bread dough rises and develops a soft, spongy texture due to alcoholic fermentation mediated by Baker's yeast, ***Saccharomyces cerevisiae***. The yeast metabolizes sugars to produce ethanol and **carbon dioxide (CO_2)** gas. As the CO_2 bubbles expand and become trapped within the gluten network, they cause the characteristic puffed-up appearance of the dough. **Answer: (B)**

Q3. Which of the following organic acids is correctly matched with its industrial microbial source organism?

(A) Citric acid — *Acetobacter aceti* (bacterium) **(B)** Acetic acid — *Aspergillus niger* (fungus)
(C) Butyric acid — *Clostridium butylicum* (bacterium) **(D)** Lactic acid — *Trichoderma polysporum* (fungus)

Sol. Q3. The precise verified NCERT pairings are: (1) **Citric acid** is produced by the filamentous fungus ***Aspergillus niger***; (2) **Acetic acid** is produced by the bacterium ***Acetobacter aceti***; (3) **Butyric acid** is produced by the obligate anaerobic bacterium ***Clostridium butylicum***; and (4) **Lactic acid** is produced by the bacterium ***Lactobacillus***. Therefore, option C is correctly matched. **Answer: (C)**

Q4. Statins, a premier class of pharmacological agents prescribed globally to lower blood cholesterol levels, are extracted from which microorganism, and through what precise biochemical mechanism do they function?

(A) From *Trichoderma polysporum*; they stimulate macrophage phagocytosis of circulating lipids **(B)** From *Monascus purpureus*; they competitively inhibit the enzyme HMG-CoA reductase
(C) From *Streptococcus*; they enzymatically digest lipid plaques within arterial walls **(D)** From *Aspergillus niger*; they block intestinal absorption of dietary triacylglycerols

Sol. Q4. Statins are bioactive molecules produced by the red mold yeast ***Monascus purpureus***. Statins act as **competitive inhibitors** of the enzyme **HMG-CoA reductase**, which catalyzes the rate-limiting step in the hepatic biosynthesis of cholesterol. By occupying the active site of HMG-CoA reductase, statins prevent the conversion of HMG-CoA to mevalonate, effectively lowering blood cholesterol levels. **Answer: (B)**

Q5. Biochemical Oxygen Demand (BOD) is a standard regulatory parameter evaluated in municipal wastewater analysis. A high BOD measurement for an effluent sample indicates that:

(A) The water has been completely purified and contains zero organic contaminants **(B)** The water carries a high concentration of biodegradable organic waste and possesses high polluting potential
(C) Aerobic microorganisms have completed the oxidation of all waste materials **(D)** The water is safe for direct human consumption and municipal distribution

Sol. Q5. Biochemical Oxygen Demand (BOD) refers to the amount of dissolved oxygen required by aerobic microbes to oxidize all biodegradable organic matter in one liter of water. Because microbes consume oxygen in direct proportion to the quantity of organic substrate present, a **high BOD directly signifies heavy organic pollution and high polluting potential**. Once secondary treatment successfully reduces the organic load, the BOD drops significantly. **Answer: (B)**

Practice Questions — Part 2 (Q6 to Q10)

Q6. During the secondary (biological) stage of sewage treatment in an aeration tank, vigorous mechanical agitation and air pumping stimulate the formation of flocs. Flocs are biologically defined as:

(A) Colonies of anaerobic methanogens forming spherical granules
(B) Masses of aerobic bacteria associated with fungal filaments to form mesh-like structures
(C) Inert mineral precipitates formed by gravitational settling of silt and grit
(D) Dense mats of photosynthetic cyanobacteria floating on the water surface

Sol. Q6. In secondary sewage treatment aeration tanks, continuous air supply and agitation foster the growth of **flocs**, which are strictly defined in NCERT as **masses of bacteria associated with fungal filaments to form mesh-like structures**. As these aerobic flocs metabolize, they consume the dissolved organic matter in the effluent, dramatically lowering its Biochemical Oxygen Demand (BOD). **Answer: (B)**

Q7. Biogas is a combustible fuel produced in anaerobic sludge digesters and rural gobar gas plants. Which of the following conditions is mandatory for methanogens to produce this gas, and what is its primary chemical component?

(A) Continuous aeration with oxygen; Carbon monoxide (CO)
(B) Completely anaerobic environment; Methane (CH_4)
(C) High acidity with sulfuric acid; Hydrogen sulfide (H_2S)
(D) Bright ultraviolet illumination; Carbon dioxide (CO_2)

Sol. Q7. Methanogens, such as ***Methanobacterium***, are obligate anaerobes that grow **strictly in the absence of molecular oxygen**. In anaerobic sludge digesters and cattle dung plants, they digest cellulosic organic materials to produce **Biogas**, which contains $50\% - 70\%$ **methane (CH_4)** as its primary combustible constituent, along with $30\% - 40\% CO_2$ and trace quantities of H_2S and H_2. **Answer: (B)**

Q8. A farmer wishes to protect crop roots against soil-borne fungal pathogens without applying synthetic chemical fungicides. Which free-living soil fungus, exceptionally common in root ecosystems, is widely employed as a biological control agent?

(A) *Rhizobium*
(B) *Trichoderma*
(C) *Glomus*
(D) *Penicillium*

Sol. Q8. ***Trichoderma*** is a genus of free-living, soil-dwelling filamentous fungi that thrive in the root ecosystems (rhizosphere). They are highly effective **biocontrol agents** against several fungal plant root pathogens, suppressing disease through competitive rhizosphere colonization, mycoparasitism, and the secretion of antimicrobial hydrolytic enzymes. **Answer: (B)**

Q9. Root nodules of leguminous plants host symbiotic bacteria capable of converting inert atmospheric nitrogen into organic nitrogen compounds. This symbiotic biofertiliser organism is:

(A) *Azotobacter*
(B) *Rhizobium*
(C) *Azospirillum*
(D) *Nostoc*

Sol. Q9. The Gram-negative bacterium ***Rhizobium*** establishes a specific symbiotic association with the roots of leguminous plants (pulses, beans, peas), developing specialized root nodules. Inside these anaerobic nodules, the bacterial enzyme nitrogenase reduces atmospheric dinitrogen (N_2) into ammonia, supplying vital organic nitrogen to the host plant while receiving carbohydrates. **Answer: (B)**

Q10. A patient is rushed to the hospital emergency department exhibiting acute chest pain diagnosed as myocardial infarction. The physician immediately administers an intravenous infusion of a genetically engineered bacterial enzyme known as a "clot buster". This enzyme is:

(A) Lipase obtained from *Candida*
(B) Streptokinase produced by *Streptococcus*
(C) Protease extracted from *Aspergillus*
(D) Pectinase derived from *Penicillium*

Sol. Q10. **Streptokinase** is an enzyme produced by the bacterium ***Streptococcus*** and subsequently modified by genetic engineering. It functions as a potent **clot buster**, activating plasminogen to dissolve intravascular fibrin clots (thrombi) that occlude coronary blood vessels, thereby restoring myocardial blood perfusion in acute heart attack patients. **Answer: (B)**

Practice Questions — Part 3 (Q11 to Q15)

Q11. The large, conspicuous holes observed in traditional Swiss cheese are produced as a result of the metabolic activity of which specific microorganism, and what gas is responsible?

(A) *Saccharomyces cerevisiae*; Ethanol vapor
(B) *Propionibacterium sharmanii*; Carbon dioxide (CO_2)
(C) *Penicillium roqueforti*; Oxygen (O_2)
(D) *Lactobacillus acidophilus*; Hydrogen gas (H_2)

Sol. Q11. The large holes characteristic of Swiss cheese are produced by the bacterium ***Propionibacterium sharmanii***. During the slow ripening process, this bacterium ferments lactic acid to produce propionic acid and large volumes of **carbon dioxide (CO_2)** gas. As bubbles of CO_2 accumulate and escape through the firm cheese matrix, they carve out the conspicuous holes. **Answer: (B)**

Q12. Commercially bottled fruit juices purchased from retail stores appear distinctly clearer and less cloudy than freshly squeezed homemade fruit juices. This commercial clarity is achieved by treating the juice with:

(A) Streptokinase and Amylases
(B) Pectinases and Proteases
(C) Lipases and Cellulases
(D) DNA ligases and Polymerases

Sol. Q12. Fresh homemade fruit juices appear cloudy and turbid due to suspended insoluble plant cell-wall pectins and proteinaceous colloids. Industrial fruit juice manufacturers treat the raw pressed juice with the hydrolytic enzymes **Pectinases** and **Proteases**. These enzymes hydrolyze the structural pectin and protein polymers into soluble fragments, rendering bottled juices crystal-clear. **Answer: (B)**

Q13. Cyclosporin A is a cyclic non-ribosomal peptide widely administered in modern clinical medicine. What is its source organism and its primary clinical application?

(A) Bacterium *Clostridium butylicum*; Dissolution of gallstones
(B) Yeast *Monascus purpureus*; Treatment of atherosclerosis
(C) Fungus *Trichoderma polysporum*; Immunosuppressive agent in organ transplant recipients
(D) Bacterium *Streptococcus*; Prevention of bacterial sepsis

Sol. Q13. **Cyclosporin A** is an 11-amino-acid cyclic peptide isolated from the soil fungus ***Trichoderma polysporum***. It acts as a powerful **immunosuppressive agent** by inhibiting calcineurin phosphatase and suppressing T-lymphocyte activation. It is administered to patients undergoing organ transplantation (e.g., kidney, liver, heart) to prevent host immune rejection of the grafted organ. **Answer: (C)**

Q14. Baculoviruses belonging to the genus *Nucleopolyhedrovirus* (NPV) are considered exceptional biological control agents in Integrated Pest Management (IPM) programs primarily because they are:

(A) Broad-spectrum poisons lethal to all soil microbes and plant roots
(B) Species-specific, narrow-spectrum insecticides with zero non-target effects on mammals, birds, or beneficial insects
(C) Rapidly mutating genetic elements that kill pests by creating antibiotic resistance
(D) Airborne fungal pathogens that sterilize all flying insects in agricultural zones

Sol. Q14. Baculoviruses of the genus ***Nucleopolyhedrovirus*** (NPV) are celebrated in sustainable agro-ecology because they are strictly **species-specific and narrow-spectrum insecticidal pathogens**. They infect specific arthropods without causing any harm to plants, mammals, birds, fish, or non-target insects such as bees and predatory beetles, making them ideal for Integrated Pest Management (IPM). **Answer: (B)**

Q15. In terrestrial ecosystems, many plant families establish a mutualistic fungal symbiosis termed mycorrhiza, exemplified by the genus *Glomus*. What is the primary nutritional benefit acquired by the plant partner from this fungal association?

(A) Direct synthesis and donation of sucrose carbohydrates
(B) Absorption of phosphorus from the soil and its translocation to the plant
(C) Enzymatic conversion of atmospheric nitrogen gas into nitrate
(D) Production of toxic alkaloids that accelerate leaf abscission

Sol. Q15. In a mycorrhizal mutualism, such as those formed by the fungal genus ***Glomus***, the extensive fungal mycelial network acts as an extension of the plant's root system, absorbing **phosphorus** from the soil and translocating it directly to the host plant. Additionally, mycorrhizal associations confer heightened resistance to root-borne pathogens and enhance tolerance to drought and salinity. **Answer: (B)**

© OurMocks · www.ourmocks.com · CUET UG Biology · Microbes in Human Welfare · 2027 Cycle