OurMocks

2027 CYCLE

MOCK · PRACTICE · CRACK

CUET UG

Biology

Evolution

Official Syllabus Unit VII · Class 12 · CUET UG 2027 Cycle Notes

INSIDE
Concept Notes

INSIDE
Solved Examples

INSIDE
Practice MCQs + Solutions

Aligned to the latest official syllabus · Concept-first explanations

www.ourmocks.com

Syllabus Alignment & Exam Blueprint

Sub-Topic	Official NTA CUET UG Syllabus Scope	High-Yield Exam Focus
Origin of Life	Origin of life; Miller's experiment; abiogenesis; Oparin-Haldane hypothesis; early atmosphere; RNA world.	Miller-Urey gas proportions (CH_4, NH_3, H_2, H_2O at 800°C), absence of free O_2, amino acid synthesis.
Concepts & Theories	Concepts of evolution; Lamarckism; Darwin's contribution; Wallace; Hugo de Vries mutation theory.	Branching descent & natural selection; Saltation (single-step large mutation) vs Darwinian gradualism; Malthus.
Evidences for Evolution	Paleontology, comparative anatomy, embryology, homologous vs analogous organs, divergent and convergent evolution.	Divergent = Homology (forelimbs, Bougainvillea/Cucurbita); Convergent = Analogy (wings, eyes, flippers); Haeckel vs von Baer.
Adaptive Radiation	Adaptive radiation; Darwin's finches; Australian marsupials vs placental mammals.	Geographical radiating lines from common ancestor; parallel evolution of placental and marsupial ecological equivalents.
Biological Evolution & Selection	Natural selection; operational modes (stabilising, directional, disruptive); industrial melanism; anthropogenic selection.	**Biston betularia** soot shift; DDT and antibiotic resistance; phenotypic curve shifts (peak height, shift, two peaks).
Hardy-Weinberg Principle	Mechanism of evolution; Hardy-Weinberg law and equilibrium; 5 disturbing factors affecting genetic equilibrium.	$p^2 + 2pq + q^2 = 1$; calculation of allele and genotype frequencies; genetic drift (founder effect, bottleneck) in small pools.
Geological Evolution & Man	Evolution through geological eras; Coelacanth; dinosaur extinction; origin and chronological evolution of man.	Coelacanth (1938); 65 mya dinosaur demise; Hominid ladder & cranial capacities (habilis 650–800, erectus 900, Neanderthal 1400 cc).

CUET HIGH-YIELD EXAM ARCHITECTURE & WEIGHTAGE GUIDE

Evolution (Unit VII) forms an essential conceptual pillar in CUET UG Biology, contributing **4 to 6 questions (16–24 marks)** per exam session. Questions feature heavy rotation on three high-probability archetypes: (1) sorting organ pairs into **Homologous (Divergent)** vs **Analogous (Convergent)**, (2) Hardy-Weinberg population genetics equations ($p^2 + 2pq + q^2 = 1$) where students confuse allele frequencies (p, q) with genotype frequencies $(p^2, 2pq, q^2)$, and (3) matching chronological hominid rungs with their exact cranial capacities, diets (meat vs fruit), and cultural innovations.

Module 1: Origin of Life: Chemistry Before Biology

Life on Earth represents a continuous chain of descent, yet the emergence of the primordial living entity from inanimate chemical matter represents the foundational question of evolutionary biology.

Cosmic Evolution and Early Earth Atmosphere

- **Age of the Universe:** Almost **20 billion years old** (20 bya). The universe originated via the **Big Bang**, a singular catastrophic thermonuclear explosion of unimaginable physical magnitude that expanded spatial dimensions and cooled matter.
- **Origin of Solar System & Earth:** Gravitational condensation of interstellar hydrogen and helium gas clouds led to the formation of galaxies and stellar solar systems. Earth condensed approximately **4.5 billion years ago** (4.5 bya).
- **Early Reducing Atmosphere:** Primordial Earth possessed **no free molecular oxygen (O_2)**. The atmosphere was intensely reducing, composed primarily of water vapour (H_2O), methane (CH_4), ammonia (NH_3), and carbon dioxide (CO_2) discharged from molten volcanic eruptions.
- **Cooling & Primordial Seas:** Solar ultraviolet (UV) radiation cleaved atmospheric water vapour into hydrogen and oxygen ($2H_2O \rightarrow 2H_2 + O_2$). Light hydrogen gas escaped into space. Free oxygen reacted with methane and ammonia to form water and carbon dioxide, while ozone (O_3) layer subsequently developed. As surface temperatures dropped, torrential condensation produced ancient oceans saturated with dissolved gases.

Oparin-Haldane Hypothesis & The Miller-Urey Experiment (1953)

- **Abiogenesis / Chemical Evolution:** Proposed independently by **Alexander Oparin** (Russia) and **J.B.S. Haldane** (England). They postulated that the first living organisms originated spontaneously through step-by-step **chemical evolution** of complex organic molecules from simple inorganic precursors (Inorganic gases → Simple monomers → Macromolecules → Coacervates / Protobionts).
- **Experimental Validation:** In 1953, **Stanley Miller** (working in Harold Urey's laboratory at the University of Chicago) created an experimental apparatus simulating primordial Earth conditions:
  - **Gaseous Mixture:** Methane (CH_4), ammonia (NH_3), and hydrogen (H_2) in a sealed flask, connected to boiling water providing continuous water vapour.
  - **Energy Source:** High-voltage electric discharge passed through tungsten electrodes at approximately **800°C** to simulate intense atmospheric lightning.
  - **Biochemical Results:** After running the closed circulation system for one week, chemical analysis revealed the spontaneous synthesis of **amino acids** (predominantly glycine, alanine, and aspartic acid).
  - **Corroborating Evidence:** Similar experiments conducted by other researchers yielded sugars, nitrogenous bases, purines, pyrimidines, and pigments. Analysis of **meteorite content** revealed identical organic compounds, confirming that chemical abiogenesis occurs naturally across extraterrestrial space.

Parameter	Primordial Earth (Pre-Life)	Contemporary Earth
Atmospheric Redox State	Intensely **reducing** (no free molecular O_2)	**Oxidising** (21% O_2, 78% N_2)
Primary Atmospheric Gases	H_2O vapour, CH_4, NH_3, CO_2, H_2	N_2, O_2, CO_2, Argon, trace moisture

Parameter	Primordial Earth (Pre-Life)	Contemporary Earth
Energy Influx	Unshielded solar UV radiation, lightning, volcanic heat	Attenuated solar radiation filtered by O_3 shield
Biological Status	Abiotic chemical evolution only	Rich biodiversity; biogenesis strictly reigns

Emergence of Cellular Life & The RNA World

- **Timeline of Cellular Emergence:** First non-cellular macromolecules (RNA, proteins, polysaccharides) assembled around **3 billion years ago** (3 bya). The earliest cellular forms of life arose approximately **2,000 million years ago** (2.0 bya) in aquatic environments as anaerobic, chemoheterotrophic single cells.
- **The RNA World Paradigm:** Ribonucleic acid (RNA) was the primordial genetic and catalytic material. RNA possesses the unique dual capacity to store hereditary nucleotide sequences and execute metabolic enzymatic catalysis (as **ribozymes**, e.g., 23S rRNA peptidyl transferase). DNA later evolved from RNA as a chemically more stable repository with repair mechanisms.

Module 2: Evidences for Evolution & Comparative Anatomy

Evolutionary descent with modification is substantiated across independent biological disciplines: paleontology, comparative embryology, anatomy, and molecular biology.

Paleontological Evidence (Fossil Record)

- **Fossils:** Preserved remains, petrified hard skeletal structures, or impressions of ancient organisms entrapped within sedimentary rock strata.
- **Geological Stratification:** Deeper sedimentary rock layers contain older, structurally simpler organisms, whereas upper strata contain progressively complex, recently evolved taxa. Geological history directly parallels biological history.
- **Transitional Links: Archaeopteryx lithographica** represents the classic fossil transition between reptiles and birds, possessing reptilian teeth and long bony tail alongside avian flight feathers and wishbone.

Embryological Evidence: Haeckel vs. Karl Ernst von Baer

- **Ernst Haeckel's Biogenetic Law:** Proposed that **"Ontogeny recapitulates Phylogeny"** — that an individual organism's embryonic developmental stages retrace the adult evolutionary stages of its ancestral lineage (e.g., human embryos possessing pharyngeal gill slits and a post-anal tail).
- **Disproof by Karl Ernst von Baer:** von Baer carefully scrutinized vertebrate embryogenesis and conclusively demonstrated that **embryos never pass through the adult stages of other animal species**. Pharyngeal arches are transient embryonic features shared across vertebrate phyla, not adult fish gill slits.

Morphological & Anatomical Evidence: Homology vs. Analogy

Comparative morphology provides the most frequently tested evidence in entrance examinations, establishing ancestral relationships through structural plan analysis:

- **Homologous Structures (Divergent Evolution):**
  - **Definition:** Organs exhibiting the **same basic anatomical plan, embryonic origin, and skeletal architecture**, but adapted to perform **different physiological functions** in response to divergent ecological niches.

- **Evolutionary Mechanism: Divergent evolution** from a shared common ancestor.
- **Vertebrate Forelimbs:** The forelimbs of whales (swimming paddle), bats (flight wing), cheetahs (running limb), and humans (grasping hand) all share homologous skeletal elements: humerus, radius, ulna, carpals, metacarpals, and phalanges.
- **Plant Examples:** Thorns of **Bougainvillea** and tendrils of **Cucurbita** are homologous organs; both represent modified axillary buds adapted for defence and climbing respectively.
- **Internal Homology:** Vertebrate hearts and brains exhibit homologous architectural progression across fishes, amphibians, reptiles, birds, and mammals.

- **Analogous Structures (Convergent Evolution):**
  - **Definition:** Organs possessing **completely different anatomical structures and developmental origins**, but adapted to execute the **same biological function** due to identical environmental pressures.
  - **Evolutionary Mechanism: Convergent evolution** in unrelated lineages.
  - **Animal Examples:**
    - Wings of butterfly (folds of chitinous ectodermal membrane without bones) vs. wings of bird (feathered, bony pentadactyl forelimbs).
    - Eyes of octopus (retina develops from surface ectoderm invagination) vs. eyes of mammals (retina develops as an outpocketing of the embryonic diencephalon).
    - Flippers of penguins (flightless birds) and dolphins (aquatic placental mammals).
  - **Plant Examples:** Sweet potato (**Ipomoea batatas**, root modification for food storage) and common potato (**Solanum tuberosum**, underground stem tuber for food storage).

Organ Pair	Anatomical Nature	Shared Feature / Divergence	Evolutionary Mode
Whale flipper & Human arm	Bony pentadactyl forelimbs	Same skeletal bones, modified for swimming vs grasping	Homologous (Divergent)
Bat wing & Cheetah leg	Bony pentadactyl forelimbs	Same skeletal plan, modified for flight vs running	Homologous (Divergent)
Bougainvillea thorn & Cucurbita tendril	Modified axillary buds	Same developmental position, defence vs climbing	Homologous (Divergent)
Butterfly wing & Bird wing	Ectodermal fold vs feathered limb	Different anatomy & origin, both adapted for flight	Analogous (Convergent)
Octopus eye & Mammalian eye	Skin invagination vs brain outpocketing	Different retinal embryology, same visual focus	Analogous (Convergent)
Penguin flipper & Dolphin flipper	Avian wing vs mammalian forelimb	Different vertebrate classes, both swim in sea	Analogous (Convergent)
Sweet potato & Potato tuber	Adventitious root vs stem tuber	Different plant organs, both store starch	Analogous (Convergent)




Figure 1: Figure 1: Comparative Anatomical Evidences (Homology vs Analogy, Adaptive Radiation) and Hominid Chronological Ladder

Adaptive Radiation: Speciation Across Ecological Niches

- **Definition:** The evolutionary process wherein an ancestral species colonizes a new geographical area and rapidly radiates into diverse specialized morphological forms, each occupying a distinct ecological niche.
- **Darwin's Finches:** On the isolated Galápagos archipelago, Charles Darwin observed that an original seed-eating ancestral finch species radiated into vegetarian tree finches, insectivorous finches, cactus-dwelling finches, and woodpecker finches, each evolving distinctive beak shapes and sizes tailored to specialized diets.
- **Australian Marsupials:** In the isolated island continent of Australia, an ancestral marsupial underwent adaptive radiation to generate kangaroos, wombats, koalas, Tasmanian devils, and banded anteaters.
- **Parallel Evolution (Convergent Radiation):** When multiple adaptive radiations occur concurrently in isolated geographical regions, placental mammals evolve phenotypic forms that strikingly mirror marsupial forms occupying equivalent ecological niches:
  - Placental Mole ↔ Marsupial Mole
  - Anteater ↔ Numbat (Banded Anteater)
  - Mouse ↔ Marsupial Mouse
  - Flying Squirrel ↔ Sugar Glider
  - Bobcat ↔ Tasmanian Tiger Cat
  - Wolf ↔ Tasmanian Wolf

Module 3: Theories of Evolution: Lamarck, Darwin, & De Vries

The historical progression of evolutionary thought encompasses three major paradigms concerning the origin and inheritance of biological diversity.

Lamarckism (Theory of Inheritance of Acquired Characters)

- Proposed by French naturalist **Jean-Baptiste de Lamarck** in his treatise **Philosophie Zoologique** (1809).
- **Central Postulates:** Use and disuse of organs dictates morphological change. Continual elongation of the giraffe's neck occurred over generations as ancestors stretched to reach foliage on tall trees.
- **Scientific Refutation:** August Weismann's germplasm theory conclusively disproved Lamarckism by cutting off the tails of mice for 22 consecutive generations; offspring were invariably born with normal tails. Somatic modifications cannot alter germ cell DNA.

Darwinian Theory of Evolution (Natural Selection)

- Proposed by **Charles Darwin** based on extensive observations during his voyage around the world on the survey ship **H.M.S. Beagle** (1831–1836), particularly on the Galápagos Islands.
- **Independent Co-Discovery: Alfred Russel Wallace**, a naturalist working in the Malay Archipelago, independently arrived at identical conclusions regarding natural selection. Their joint paper was presented in 1858, followed by Darwin's seminal book **The Origin of Species** (1859).
- **Malthusian Influence:** Darwin was profoundly influenced by **Thomas Malthus's Essay on Population**, which demonstrated that populations produce offspring geometrically, whereas natural food resources increase arithmetically, enforcing an inevitable struggle for survival.
- **Two Key Concepts of Darwinism:**

1. **Branching Descent:** All living organisms are linked by descent from common ancestors through branched lineages.
2. **Natural Selection:** Individuals possessing minor heritable variations best suited to their physical and biotic environment survive and reproduce at higher rates (**differential reproduction**), leaving more progeny.

Mutation Theory of Hugo de Vries

- In 1901, Dutch botanist **Hugo de Vries** proposed the **Mutation Theory of Evolution** based on experimental breeding of the evening primrose (**Oenothera lamarckiana**).
- **Core Postulates:**
  - Evolutionary speciation is caused by **mutations** — sudden, large, heritable phenotypic alterations.
  - Mutations are **random and directionless**, in stark contrast to Darwinian minor, continuous variations which are directional.
  - Speciation occurs discontinuously in a single step termed **Saltation** (single-step large mutation), whereas Darwin viewed evolution as an extraordinarily gradual, continuous process.

Attribute	Darwinian Variations	De Vriesian Mutations (Saltation)
Nature & Magnitude	Small, continuous, minor variations	Large, sudden, discontinuous alterations
Directionality	Directional (favours adaptive survival)	Random and directionless
Tempo of Speciation	Gradual, slow divergence over ages	Sudden, jerky, single-step speciation
Primary Driver	Differential reproductive survival	Novel germline genetic mutation

Module 4: Natural Selection in Action & Anthropogenic Shifts

Natural selection is not merely a theoretical historical deduction; its real-time operation is directly verified through ecological field studies and anthropogenic selection.

Industrial Melanism: The Peppered Moth (Biston betularia)

- **Pre-Industrialization Era (Before 1850 in England):**
  - Tree trunks in countryside forests were heavily carpeted with thick white lichens and free from industrial soot.
  - The **white-winged (melanic recessive)** peppered moths were exquisitely camouflaged against the pale lichen-covered bark.
  - Dark-winged (melanic dominant, **carbonaria**) moths were conspicuous against the white background and were heavily predated by insectivorous birds. Consequently, white moths constituted over 95% of the population.
- **Post-Industrialization Era (By 1920 in Industrial Areas):**
  - Coal smoke, particulate emissions, and industrial soot blackened tree trunks; sensitive lichens perished completely due to sulfur dioxide pollution.
  - Against the blackened, soot-covered bark, the **dark-winged melanic moths** enjoyed superior camouflage protection.
  - White-winged moths became highly conspicuous and suffered intense bird predation.
  - The population frequency of dark moths surged dramatically in industrial cities like Birmingham and Manchester.

- **Key Evolutionary Deductions:**
  - Natural selection operates on **pre-existing genetic variations**; soot did not generate the melanic allele, it simply altered the selective survival value of existing morphs.
  - **No variant was completely eradicated.** In rural unpolluted regions, white moths remained dominant.
  - In recent decades, clean-air legislation reduced coal combustion, causing lichen recovery and an observed decline in dark moth frequency. This proves that natural selection is dynamic and reversible.

Anthropogenic Selection: Resistance to Antibiotics & Pesticides

- **Antibiotic Resistance in Bacteria:** Indiscriminate clinical administration of antibiotics destroys sensitive bacterial populations. Rare pre-existing mutant bacteria possessing antibiotic-resistance plasmids survive, proliferate uninhibitedly, and establish resistant strains within months.
- **Pesticide Resistance:** Excessive spraying of DDT killed susceptible mosquitoes, but selected for rare mutant mosquitoes possessing enzymes capable of detoxifying DDT. Within a few seasons, DDT-resistant mosquito populations dominated.
- **Evolution is Stochastic:** These examples prove that evolution is not a directed teleological process; it is a **stochastic process** governed by chance mutations and environmental selection pressures.




Figure 2: Figure 2: Three Modes of Natural Selection (Stabilising, Directional, Disruptive) and Hardy-Weinberg Dynamics

Module 5: Hardy-Weinberg Principle & Population Genetics

In 1908, British mathematician **G.H. Hardy** and German physician **Wilhelm Weinberg** independently formulated the foundational mathematical law governing population genetics.

The Hardy-Weinberg Principle & Algebraic Formulation

- **The Principle:** In a large, randomly mating diploid population devoid of evolutionary disturbances, allele frequencies and genotype frequencies remain **constant and invariant from generation to generation** — a condition termed **genetic equilibrium** or **Hardy-Weinberg equilibrium**.
- **The Algebraic Binomial Expansion:** Consider a single gene locus with two alternative alleles: dominant allele A and recessive allele a. Let the gene frequency of allele $A = p$ and the frequency of allele $a = q$. Since these are the only two alleles at this locus:

$$p + q = 1$$

The diploid genotype frequencies generated by random union of gametes follow the binomial expansion:

$$(p + q)^2 = p^2 + 2pq + q^2 = 1$$

- p^2 = Frequency of homozygous dominant individuals (AA)
- $2pq$ = Frequency of heterozygous individuals (Aa)
- q^2 = Frequency of homozygous recessive individuals (aa)
- $p^2 + 2pq$ = Total frequency of individuals displaying the dominant phenotype.

- **Evolution Defined:** Any measured deviation of actual genotype frequencies from Hardy-Weinberg expectations indicates that evolutionary forces are actively driving population divergence.

Five Evolutionary Forces Disrupting Hardy-Weinberg Equilibrium

1. **Gene Migration / Gene Flow:** Movement of a section of a population into another geographic region. When emigrants breed with the recipient population, new alleles are introduced into the new gene pool while being lost from the donor pool. Continuous, repeated migration is termed **gene flow**.
2. **Genetic Drift:**
   - **Definition:** Non-directional, random fluctuations in allele frequencies occurring purely **by chance** in **small, isolated populations**.
   - Unlike natural selection, genetic drift operates regardless of whether an allele is beneficial, neutral, or harmful.
   - **Founder Effect:** When a small group of individuals becomes geographically isolated from the parental population, their random allele frequencies establish the genetic foundation of the new colony. The isolated colonists are the "founders", and the resultant divergence is termed the **Founder Effect**.
   - **Bottleneck Effect:** Severe ecological catastrophes (epidemics, floods, volcanic eruptions) drastically reduce population size to a small surviving fraction, randomly eliminating alleles.
3. **Mutation:** Spontaneous alteration in DNA sequences introducing novel alleles into the gene pool. Advantageous mutations are rare, but provide fundamental raw material for long-term evolutionary divergence.
4. **Genetic Recombination:** Crossing over between non-sister chromatids during pachytene of meiosis I, coupled with random assortment of homologous chromosomes, generates novel allelic combinations in gametes.
5. **Natural Selection:** Differential reproductive success of phenotypes resulting in directional shifts in allele frequencies. Natural selection operates through three distinct phenotypic modes:
   - **Stabilising Selection:** Favours individuals with intermediate (mean) phenotypes. The phenotypic distribution curve becomes narrower and taller. **Example:** Human infant birth weight — babies weighing between 2.5 and 3.5 kg exhibit the lowest perinatal mortality.
   - **Directional Selection:** Favours individuals at one extreme of the phenotypic distribution. The entire distribution peak shifts toward that extreme. **Example:** Industrial melanism in **Biston betularia**; evolution of DDT-resistant mosquitoes.
   - **Disruptive Selection:** Favours individuals at both peripheral extremes while penalizing intermediate phenotypes. The distribution curve splits into **two distinct phenotypic peaks**, fostering polymorphism and potential sympatric speciation.

WORKED EXAMPLE — STEP-BY-STEP HARDY-WEINBERG ALLELE & GENOTYPE COMPUTATION

Problem Statement: In a human population of 10,000 individuals adhering to Hardy-Weinberg equilibrium, 900 individuals exhibit an autosomal recessive trait (cystic fibrosis, genotype aa).

- **Step 1: Calculate recessive genotype frequency (q^2):**

$$q^2 = \frac{900}{10000} = 0.09$$

- **Step 2: Calculate recessive allele frequency (q):**

$$q = \sqrt{0.09} = 0.3$$

- **Step 3: Calculate dominant allele frequency (p):** Since $p + q = 1$, we have:

$$p = 1 - 0.3 = 0.7$$

- **Step 4: Calculate homozygous dominant genotype frequency (p^2):**

$$p^2 = (0.7)^2 = 0.49 \quad \rightarrow \quad 49\% \quad (4,900 \text{ individuals})$$

- **Step 5: Calculate heterozygous carrier frequency ($2pq$):**

$$2pq = 2 \times 0.7 \times 0.3 = 0.42 \quad \rightarrow \quad 42\% \quad (4,200 \text{ individuals})$$

- **Step 6: Total dominant phenotype individuals ($p^2 + 2pq$):**

$$0.49 + 0.42 = 0.91 \quad \rightarrow \quad 91\% \quad (9,100 \text{ individuals})$$

Crucial Exam Rule: Never calculate p first by taking the square root of the dominant population fraction, because dominant individuals consist of both p^2 (homozygotes) and $2pq$ (heterozygotes)! Always solve for q from q^2 first!

Module 6: Evolution Through Geological Eras & Major Milestones

Earth's 4.5-billion-year chronological history is subdivided into **Eras** (Palaeozoic, Mesozoic, Cenozoic), which are subdivided into **Periods** and **Epochs**.

Chronological Evolutionary Milestones Across Geological Time

- **2000 mya (Precambrian):** First cellular life forms emerged on Earth.
- **500 mya (Palaeozoic - Cambrian/Ordovician):** Invertebrates formed and dominated marine ecosystems.
- **350 mya (Silurian/Devonian):** Jawless fishes (**Agnatha**) evolved; early fishes dominated the waters. Early plants colonized terrestrial habitats.
- **320 mya (Carboniferous):** Giant seaweeds and primitive terrestrial flora flourished. Vast coal deposits formed from giant ferns and lycopods.
- **The Coelacanth Discovery (1938):** In 1938, a live fish caught in South Africa was identified as a **Coelacanth** (a **Lobefin**), thought to have been extinct for 65 million years. Lobefins possessed sturdy, fleshy, muscular fins that enabled them to crawl onto land and return to water — representing the transitional ancestors of the first terrestrial **Amphibians**.
- **200 mya (Mesozoic - Triassic/Jurassic):** Some land reptiles retreated back into the aquatic environment to evolve into fish-like marine reptiles, exemplified by **Ichthyosaurs**. Giant terrestrial reptiles (**Dinosaurs**) dominated the planet. The most formidable carnivorous dinosaur was **Tyrannosaurus rex**, standing about **20 feet in height** and possessing ferocious dagger-like teeth.

- **65 mya (Cretaceous-Tertiary Boundary):** Dinosaurs abruptly disappeared from the geological record, possibly due to massive bolide asteroid impact and subsequent global climatic collapse. Mammals, which were previously small, shrew-like nocturnal creatures, radiated rapidly into vacated ecological niches.

Geological Era	Period & Chronology	Major Evolutionary Radiations & Extinctions
Palaeozoic (Ancient Life)	Cambrian to Permian (540–250 mya)	Invertebrates thrive (500 mya); jawless fish (350 mya); lobefin Coelacanth crawls ashore; first amphibians; coal forests of giant lycopods.
Mesozoic (Age of Reptiles)	Triassic, Jurassic, Cretaceous (250–65 mya)	Ichthyosaurs return to water (200 mya); dinosaurs and cycads dominate; Archaeopteryx evolves; abrupt dinosaur extinction at 65 mya.
Cenozoic (Age of Mammals)	Tertiary & Quaternary (65 mya to Present)	Adaptive radiation of placental mammals and birds; dominance of angiosperms; hominid evolution from Dryopithecus to modern Homo sapiens.

Module 7: Origin and Evolution of Man (Hominid Phylogeny)

Human evolution represents one of the most meticulously documented and heavily tested evolutionary sequences in the NCERT curriculum.

Chronological Hominid Ladder & Diagnostic Features

1. **Dryopithecus & Ramapithecus (15 mya - Miocene):**
   - Hairy primates existing in Africa and Asia.
   - **Dryopithecus:** More ape-like; possessed long arms; walked like chimpanzees and gorillas (knuckle-walkers).
   - **Ramapithecus:** More man-like; possessed shorter canines and a more erect posture. Discovered in the Shivalik hills of India.
2. **Australopithecines (3–4 to 2 mya - Pliocene):**
   - Fossils discovered in East African grasslands (Tanzania, Ethiopia; famous skeleton **"Lucy"**).
   - Walked **fully erect and bipedal**.
   - **Brain Capacity:** Approximately **400 to 500 cc**.
   - **Habits:** Hunted with primitive stone weapons, but **essentially ate fruit** (herbivorous/ frugivorous).
3. **Homo habilis (2.0 mya - Early Pleistocene):**
   - The **first human-like being (the hominid)**. Known colloquially as the "handy man" or tool-maker.
   - **Brain Capacity:** Diagnostic range of **650 to 800 cc**.
   - **Dietary Trait:** Crucial NCERT distinction — **Homo habilis did NOT eat meat**.
4. **Homo erectus (1.5 mya - Middle Pleistocene):**
   - Fossils discovered in Java, Indonesia in 1891 ("Java Man").
   - **Brain Capacity:** Exactly **900 cc**.
   - **Dietary Trait: Homo erectus probably ate meat** (carnivorous hunter). Mastered fire and crafted advanced Acheulean stone hand-axes.
5. **Homo neanderthalensis (Neanderthal Man; 100,000–40,000 years ago):**
   - Lived across the Near East and Central Asia.
   - **Brain Capacity:** Exactly **1,400 cc** (comparable to or slightly larger than modern humans).

- **Cultural Traits:** Used **animal hides** to insulate and protect their bodies; performed ritualistic burial of their dead. Not a direct ancestor of modern humans, but an evolutionary cousin that coexisted and went extinct.

6. **Homo sapiens (Modern Man; 75,000–10,000 years ago):**
   - Arose in Africa during the last Ice Age (75,000–10,000 years ago) and migrated across continents, diverging into distinct geographical races.
   - **Brain Capacity: 1,350 to 1,450 cc**.
   - **Cultural Milestones:** Prehistoric cave art developed approximately **18,000 years ago** (superb examples preserved at the Bhimbetka rock shelters in Raisen district, Madhya Pradesh); agriculture and human settlements commenced approximately **10,000 years ago**.

Hominid Species	Chronological Age	Cranial Capacity	Salient Traits, Diet, & Culture
Dryopithecus	15 mya	Ape-like (400 cc)	Arboreal, knuckle-walking, hairy; ancestral to both apes and humans.
Ramapithecus	15 mya	More man-like	Walked more erect; discovered in Shivalik hills; smaller canines.
Australopithecines	3–4 mya	400–500 cc	Bipedal upright walking in East Africa (Lucy); stone tools; fruit eaters.
Homo habilis	2.0 mya	650–800 cc	First hominid (toolmaker / "handy man"); strictly non-meat eater.
Homo erectus	1.5 mya	900 cc	Java man (1891); fully upright; ate meat; discovered and used fire.
Neanderthal Man	100k–40k ya	1,400 cc	Near East & Central Asia; used animal hides for clothing; buried dead.
Homo sapiens	75k–10k ya	1,350–1,450 cc	Modern humans; cave paintings (Bhimbetka, 18k ya); agriculture (10k ya).

OurMocks

Key Facts & Concept Cheatsheet

HIGH-YIELD EVOLUTION EXAMINATION CHEATSHEET

Concept / Milestone	Biological Definition / Mechanism	NCERT Anchor Metric / Exam Rule
Miller-Urey Setup (1953)	Spark discharge simulating primitive reducing atmosphere	CH_4:NH_3:H_2 + H_2O at 800°C → Amino acids
Homologous Organs	Same embryonic origin, different functions; divergent evolution	Whale/bat/cheetah/man forelimbs; Bougainvillea/Cucurbita
Analogous Organs	Different origins, same function; convergent evolution	Bird/butterfly wings; sweet potato (root) vs potato (stem)
Adaptive Radiation	Single ancestral stock radiating into diverse ecological niches	Darwin's finches; Australian marsupials vs placentals
Hardy-Weinberg Equation	Genetic equilibrium: allele & genotype frequencies invariant	$p^2 + 2pq + q^2 = 1$; $p + q = 1$; 5 disturbing factors
Genetic Drift	Random allele frequency fluctuations by chance in small pools	Founder effect & bottleneck effect; non-adaptive
Natural Selection Modes	Stabilising (mean), Directional (one extreme), Disruptive (both)	Birth weight (stabilising); industrial melanism (directional)
De Vries Mutation Theory	Evolution proceeds via sudden single-step large mutations	**Saltation**; mutations are random & directionless
Living Fossil Lobefin	Coelacanth caught in South Africa (1938); ancestral to amphibians	Fleshy fins enabled crawling onto land 350 mya
Dinosaur Demise	Dominant Mesozoic reptiles extinct at Cretaceous boundary	Disappeared suddenly 65 mya; T. rex was 20 ft tall
Hominid Brain Capacities	H. habilis → H. erectus → Neanderthal → H. sapiens	650–800 cc → 900 cc → 1400 cc → 1350–1450 cc
Hominid Diets & Culture	Dietary and cultural markers distinguishing Homo species	H. habilis: no meat; H. erectus: ate meat; Neanderthal: buried dead




Figure 3: Figure 3: Whole-Chapter Concept Synthesis Mindmap: Origin of Life, Theories, Evidences, Mechanisms, Geological Milestones, and Hominid Lineage

Practice Questions — Part 1 (Q1 to Q5)

Q1. In his famous 1953 closed-flask simulation of primitive Earth conditions, Stanley Miller passed continuous electric spark discharges through a gaseous mixture maintained at approximately 800°C. Which of the following correctly identifies the gaseous components present inside the reaction vessel?

(A) Carbon dioxide (CO_2), molecular oxygen (O_2), nitrogen (N_2), and water vapour
(B) Methane (CH_4), ammonia (NH_3), hydrogen (H_2), and water vapour
(C) Methane (CH_4), molecular oxygen (O_2), ammonia (NH_3), and carbon monoxide
(D) Carbon dioxide (CO_2), ammonia (NH_3), hydrogen (H_2), and sulfur dioxide

Sol. Q1. Stanley Miller simulated the primordial reducing atmosphere of prebiotic Earth using methane (CH_4), ammonia (NH_3), molecular hydrogen (H_2), and circulating water vapour at 800°C. Crucially, molecular oxygen (O_2) was strictly absent. The reaction yielded amino acids (glycine, alanine, aspartic acid), confirming the Oparin-Haldane chemical evolution hypothesis. **Answer: (B)**

Q2. In England, field collections of the peppered moth (Biston betularia) demonstrated that before industrialization (1850s), white-winged moths were abundant, whereas after industrialization (1920s), dark-winged melanic moths dominated industrial areas. This evolutionary shift constitutes a classical demonstration of:

(A) Stabilising natural selection operating against both phenotypic extremes
(B) Directional natural selection favouring an adaptive phenotypic extreme
(C) Disruptive natural selection splitting the population into two discrete peaks
(D) Lamarckian inheritance of acquired dark pigmentation induced by coal soot

Sol. Q2. Industrial melanism is a textbook example of directional natural selection. Soot from coal combustion covered tree trunks and killed pale lichens in industrial regions. Dark melanic moths (**carbonaria**) were camouflaged against birds on soot-darkened trunks, while white moths were easily predated. The phenotypic distribution shifted directionally toward melanism. No variant was created de novo or completely eliminated. **Answer: (B)**

Q3. Which of the following paired biological structures represents an authentic example of homologous organs resulting from divergent evolutionary descent?

(A) The wing of a butterfly and the wing of a bird
(B) The eye of an octopus and the eye of a mammal
(C) The thorn of Bougainvillea and the tendril of Cucurbita
(D) The flipper of a penguin and the flipper of a dolphin

Sol. Q3. Thorns of **Bougainvillea** and tendrils of **Cucurbita** are homologous organs: both are modified axillary buds sharing the same embryonic origin and anatomical position, but modified to perform different functions (defence vs climbing). In contrast, butterfly/bird wings, octopus/mammal eyes, and penguin/dolphin flippers are analogous organs resulting from convergent evolution. **Answer: (C)**

Q4. Sweet potato (Ipomoea batatas) and common potato (Solanum tuberosum) both function as specialized storage organs for edible carbohydrates. From an evolutionary perspective, these organs are classified as:

(A) Homologous structures that have undergone divergent evolutionary radiation
(B) Analogous structures that have evolved independently via convergent evolution
(C) Vestigial structures inherited from a common ancestral angiosperm
(D) Atavistic organs reflecting ancestral underground root morphology

Sol. Q4. Sweet potato is an anatomically modified adventitious root, whereas common potato is an anatomically modified underground stem tuber (possessing nodes, internodes, and scale leaves). Because they share completely different morphological origins but perform the identical biological function of food storage, they are analogous structures developed through convergent evolution. **Answer: (B)**

Q5. The evolution of diverse forms of Australian marsupials (such as the marsupial mole, wombat, banded anteater, and kangaroo) from a single ancestral marsupial stock within the isolated Australian continent represents an outstanding example of:

(A) Adaptive radiation
(B) Saltation and single-step macromutation
(C) Stabilising natural selection
(D) Anthropogenic selective breeding

Sol. Q5. Adaptive radiation is the evolutionary diversification of a single ancestral species into multiple distinctly specialized forms that radiate into different ecological niches within a geographical territory. Darwin's finches on the Galápagos Islands and Australian marsupials on the island continent of Australia are the two premier examples described in NCERT Class 12. **Answer: (A)**

Practice Questions — Part 2 (Q6 to Q10)

Q6. In a large human population conforming strictly to Hardy-Weinberg equilibrium, the proportion of individuals afflicted with an autosomal recessive condition (albinism) is observed to be 16% (0.16). What is the expected frequency of heterozygous carrier individuals in this population?

(A) 0.16 (16%) **(B)** 0.24 (24%)
(C) 0.48 (48%) **(D)** 0.84 (84%)

Sol. Q6. In Hardy-Weinberg mathematics, homozygous recessive individuals (aa) have a frequency of q^2. Here, $q^2 = 0.16$, which yields $q = \sqrt{0.16} = 0.4$. Since $p + q = 1$, the frequency of the dominant allele A is $p = 1 - 0.4 = 0.6$. The frequency of heterozygous carriers (Aa) is given by $2pq = 2 \times 0.6 \times 0.4 = 0.48$ (or 48%). The common trap is mistaking carrier frequency for $2q$ or forgetting the factor of 2. **Answer: (C)**

Q7. A catastrophic volcanic eruption isolates a tiny cohort of 15 lizards on an offshore oceanic reef. Over subsequent generations, the allele frequencies of the newly established reef population diverge dramatically from the large mainland parental population due to purely stochastic sampling events. This phenomenon exemplifies:

(A) Gene flow resulting from continuous directional migration **(B)** Genetic drift exhibiting the Founder Effect in a small population
(C) Disruptive selection favouring intermediate heterozygotes **(D)** Sympatric speciation driven by polyploidization

Sol. Q7. Genetic drift refers to random, non-directional fluctuations in allele frequencies occurring purely by chance in small, isolated populations. When a minuscule group emigrates and colonizes an isolated territory, their random allelic sample forms the genetic basis of the new lineage — a specific manifestation of genetic drift termed the **Founder Effect**. The new population may become distinct enough to constitute a new species. **Answer: (B)**

Q8. Which of the following operational modes of natural selection acts to penalize both extreme phenotypes while simultaneously increasing the proportion of individuals exhibiting the intermediate mean character state?

(A) Directional selection **(B)** Stabilising selection
(C) Disruptive selection **(D)** Artificial selection

Sol. Q8. Stabilising selection favours the intermediate (mean) character value and eliminates peripheral phenotypes at both extremes. As a consequence, the phenotypic distribution curve becomes narrower and taller over generations. A classic example is human birth weight: infants with birth weights between 2.5 and 3.5 kg have significantly higher survival rates than low-birth-weight (under 2 kg) or heavy (over 4.5 kg) infants. **Answer: (B)**

Q9. Hugo de Vries formulated his Mutation Theory of evolution based on experimental hybridization of evening primrose (Oenothera lamarckiana). According to de Vries, evolutionary speciation differs fundamentally from Darwin's concept because mutations are:

(A) Minor, continuous, and directional variations **(B)** Single-step, large, random, and directionless alterations termed Saltation
(C) Inherited epigenetic modifications induced by somatic usage of organs **(D)** Gradual microscopic changes accumulated exclusively over millions of generations

Sol. Q9. Hugo de Vries postulated that speciation is caused by sudden, discrete, large-scale genetic alterations termed **Saltation** (single-step large mutation). He emphasized that mutations are **random and directionless**, sharply contrasting with Darwinian variations which are small, continuous, gradual, and directional. **Answer: (B)**

Q10. In 1938, a live specimen of Coelacanth (a Lobefin fish) was netted off the coast of South Africa. The biological significance of this discovery resides in the fact that lobefins:

(A) Are the direct ancestors of modern cartilaginous sharks **(B)** Represented transitional living fossils whose muscular fins evolved into the limbs of the first terrestrial amphibians
(C) Proved that marine reptiles such as Ichthyosaurs never existed **(D)** Represented the earliest known jawless agnathan vertebrates

Sol. Q10. The Coelacanth is a lobefin fish that was long presumed to be extinct for 65 million years. Lobefins possessed stout, fleshy, muscular fins containing internal bones, enabling them to crawl onto muddy terrestrial banks and return to water. They represent the direct evolutionary transition between fishes and the earliest terrestrial amphibians that emerged approximately 350 million years ago. **Answer: (B)**

Practice Questions — Part 3 (Q11 to Q15)

Q11. Giant carnivorous reptiles, exemplified by Tyrannosaurus rex, dominated terrestrial ecosystems during the Mesozoic Era before undergoing sudden global extinction. According to the geological record, dinosaurs appeared and abruptly vanished approximately:

(A) 500 million years ago and 200 million years ago respectively
(B) 350 million years ago and 150 million years ago respectively
(C) 230 million years ago and 65 million years ago respectively
(D) 65 million years ago and 10,000 years ago respectively

Sol. Q11. Dinosaurs emerged during the Triassic Period of the Mesozoic Era approximately 230 million years ago and dominated Earth for over 160 million years before disappearing abruptly at the Cretaceous-Tertiary (K-T) boundary approximately **65 million years ago** (65 mya). Following their extinction, small mammalian lineages radiated into terrestrial niches. **Answer: (C)**

Q12. Ernst Haeckel proposed the Biogenetic Law, stating that embryonic development recapitulates ancestral evolutionary history. Which eminent scientist decisively refuted this generalization by demonstrating that embryos never pass through the adult stages of other animals?

(A) Alexander Oparin
(B) Karl Ernst von Baer
(C) Thomas Malthus
(D) Alfred Russel Wallace

Sol. Q12. Karl Ernst von Baer carefully analyzed vertebrate embryogenesis and explicitly refuted Haeckel's biogenetic law. von Baer demonstrated that during development, generalized embryonic features (such as pharyngeal arches and notochord) appear before specialized traits, and vertebrate embryos **never pass through the adult morphological stages of other animal species**. **Answer: (B)**

Q13. Which of the following options arranges the evolutionary ancestors of modern humans in the strictly correct chronological sequence of their historical emergence on Earth?

(A) Homo habilis → Australopithecus → Homo erectus → Homo neanderthalensis → Homo sapiens
(B) Australopithecus → Homo habilis → Homo erectus → Homo neanderthalensis → Homo sapiens
(C) Australopithecus → Homo erectus → Homo habilis → Homo neanderthalensis → Homo sapiens
(D) Ramapithecus → Homo erectus → Australopithecus → Homo habilis → Homo sapiens

Sol. Q13. The verified NCERT chronological sequence of hominid evolution is: Dryopithecus/Ramapithecus (15 mya) → Australopithecines (3–4 mya, Lucy) → Homo habilis (2.0 mya, toolmaker) → Homo erectus (1.5 mya, Java man) → Neanderthal man (100,000–40,000 ya) → Homo sapiens (75,000–10,000 ya, modern humans). Homo habilis preceded Homo erectus. **Answer: (B)**

Q14. Match the hominid species listed in Column I with their corresponding cranial capacities detailed in Column II:

Column I:
(a) Homo habilis
(b) Homo erectus
(c) Neanderthal man
(d) Homo sapiens

Column II:
(i) 900 cc
(ii) 1,400 cc
(iii) 650–800 cc
(iv) 1,350–1,450 cc

(A) (a)-(iii), (b)-(i), (c)-(ii), (d)-(iv)
(B) (a)-(i), (b)-(iii), (c)-(ii), (d)-(iv)
(C) (a)-(iii), (b)-(ii), (c)-(i), (d)-(iv)
(D) (a)-(iv), (b)-(i), (c)-(ii), (d)-(iii)

Sol. Q14. The precise cranial capacities established in NCERT Class 12 are: **Homo habilis** = 650–800 cc; **Homo erectus** = 900 cc; **Neanderthal man** = 1,400 cc; **Homo sapiens** = 1,350–1,450 cc. Matching gives: (a)-(iii), (b)-(i), (c)-(ii), (d)-(iv). **Answer: (A)**

Q15. Which of the following hominid species is correctly paired with its documented dietary habit and cultural behavioral characteristics?

(A) Homo habilis — Exclusively carnivorous hunter that mastered controlled fire
(B) Homo erectus — Herbivorous fruit-eater that buried dead with religious rituals
(C) Homo neanderthalensis — Used animal hides to protect bodies and performed burial of dead
(D) Australopithecus — Created prehistoric cave paintings at Bhimbetka 18,000 years ago

Sol. Q15. Neanderthal man lived in Europe and Central Asia 100,000–40,000 years ago, used animal hides for clothing, and buried their dead with ceremonial rites. In contrast, Homo habilis did **NOT** eat meat; Homo erectus **probably ate meat**; and cave paintings at Bhimbetka were created by **Homo sapiens** 18,000 years ago during the late ice age. **Answer: (C)**

© OurMocks · www.ourmocks.com · CUET UG Biology · Evolution · 2027 Cycle