OurMocks

2027 CYCLE

MOCK · PRACTICE · CRACK

CUET UG

Biology

Human Reproduction

Official Syllabus Unit VI · Class 12 · CUET UG 2027 Cycle Notes

INSIDE
Concept Notes

INSIDE
Solved Examples

INSIDE
Practice MCQs + Solutions

Aligned to the latest official syllabus · Concept-first explanations

www.ourmocks.com

Syllabus Alignment

Sub-topic	Official syllabus pointer (NTA CUET UG Biology 2026 / 2027 Baseline)	Exam Skills & Question Types
Male Reproductive System	Male reproductive system; microscopic anatomy of testis; accessory ducts and glands.	Scrotal thermoregulation (2° C drop); Sertoli vs Leydig cells; duct order; seminal fluid biochemistry.
Female Reproductive System	Female reproductive system; microscopic anatomy of ovary; fallopian tube and uterine wall layers.	Ampullary fertilization site; endometrium cyclical shedding; myometrium oxytocin response; ovarian histology.
Gametogenesis & Sperm/Ovum	Gametogenesis — spermatogenesis and oogenesis; structural organization of gametes.	Spermatogonia to sperm pathway; spermiogenesis vs spermiation; dual female meiotic arrests (Prophase I, Metaphase II).
Menstrual Cycle Endocrinology	Menstrual cycle: phases, hormonal regulation (LH, FSH, estrogen, progesterone).	28-day timeline; mid-cycle LH surge; follicular vs luteal phases; corpus luteum progesterone secretion.
Fertilisation & Implantation	Fertilisation, embryo development up to blastocyst formation, implantation.	Capacitation; acrosome reaction; cortical block to polyspermy; cleavage; morula; blastocyst ICM vs trophoblast.
Pregnancy, Placenta & Birth	Pregnancy and embryonic development; placenta; parturition and lactation.	Placental hormones (hCG, hPL, relaxin); fetal ejection reflex; oxytocin positive feedback; colostrum IgA.

EXAM PATTERN

Official CUET UG Biology Pattern (NTA 2026/2027 Baseline): The Biology domain paper carries **50 compulsory questions** to be answered in **60 minutes** (+5 correct, −1 incorrect). Human Reproduction is among the top three highest-frequency chapters in domain Biology, with heavy emphasis on hormone flux graphs, cellular ploidy, developmental arrest checkpoints, and sequence-based reproductive physiology.

Module 1: The Male Reproductive System & Testicular Histology

Humans are sexually reproducing, viviparous mammals exhibiting marked sexual dimorphism. The male reproductive system is situated predominantly in the pelvic region and comprises primary sex organs (**testes**), secondary accessory ducts, accessory reproductive glands, and external genitalia.

Scrotal Thermoregulation & Testis Micro-Anatomy

- **Scrotal Thermoregulation:** The testes are extra-abdominal organs suspended in a cutaneous pouch called the **scrotum**.
  - **Temperature Differential:** The scrotum functions as a biological thermostat, maintaining testicular temperature **2 degree "C" to 2.5 degree "C" lower than internal body temperature** (34.5° C to 35° C vs 37° C).
  - **Spermatogenic Imperative:** Human germinal epithelium cannot undergo meiosis at core abdominal temperatures. Undescended testes (**cryptorchidism**) result in complete germ cell atrophy and permanent sterility, though androgen production by Leydig cells remains partially functional.
  - **Thermoregulatory Muscles:** The **dartos muscle** in the scrotal subcutaneous tissue contracts in cold weather to reduce surface area, while the **cremaster muscle** elevates the testes closer to the pelvic floor for warmth.
- **Testicular Compartmentalization:**
  - Each adult testis is an oval body (length 4–5 cm, width 2–3 cm, weight 10–15 g) encased in a tough fibrous capsule, the **tunica albuginea**.
  - Internally, fibrous septa partition each testis into approximately **250 testicular lobules**.
  - Each lobule contains **1 to 3 highly convoluted seminiferous tubules**, totaling approximately 750 tubules per testis, with an aggregate length exceeding 250 meters.

Cytology of the Seminiferous Tubule Niche

A transverse section of a seminiferous tubule reveals a complex germinal epithelium surrounded by a specialized interstitial microenvironment:

Cell Type	Ploidy	Histological Architecture & Exact Physiological Role
Spermatogonia (Male Germ Cells)	$2n$ (46 chr)	Line the inner basement membrane; stem cell reserve (**Type A**) renews population, while committed cells (**Type B**) differentiate into primary spermatocytes.
Primary Spermatocytes	$2n$ (46 chr, 92 chromatids)	Enormous cells undergoing Meiosis I; synthesize RNA and proteins during prolonged prophase I before reductional division.
Secondary Spermatocytes	n (23 chr, 46 chromatids)	Short-lived haploid intermediates that immediately undergo Meiosis II (equational division) without an intervening interphase.
Spermatids	n (23 chr, 23 chromatids)	Small, circular, non-motile haploid cells embedded in Sertoli cell apical recesses; undergo spermiogenesis.
Sertoli Cells (Nurse / Sustentacular)	$2n$ (Somatic)	Tall pyramidal epithelial cells spanning basement membrane to lumen. Key roles: (1) physical support and nutritional supply to germ cells; (2) phagocytose discarded spermatid residual bodies; (3) construct the **blood-testis barrier** via basal tight junctions, shielding foreign haploid antigens from the immune system; (4) secrete **Androgen-Binding Protein (ABP)** under FSH stimulation

Cell Type	Ploidy	Histological Architecture & Exact Physiological Role
		to concentrate testosterone; (5) synthesize **Inhibin** to exert selective negative feedback on anterior pituitary FSH.
Leydig Cells (Interstitial Cells)	$2n$ (Somatic)	Clustered in interstitial connective tissue between seminiferous tubules alongside blood vessels and immunocompetent cells. Stimulated by pituitary **LH (ICSH)** to synthesize and secrete **androgens** (principally **Testosterone**).

Male Accessory Ducts & Transport Architecture

Spermatozoa leave the seminiferous tubules and traverse a strict anatomical hierarchy of paired conduits:

Seminiferous Tubules → Rete Testis → Vasa Efferentia → Epididymis → Vas Deferens → Ejaculatory Duct → Urethra

- **Rete Testis:** A labyrinth of anastomosing cuboidal-lined tubules in the mediastinum testis that aggregates sperm from all seminiferous tubules.
- **Vasa Efferentia (10 to 20 fine tubules):** Ciliated ducts that pierce the tunica albuginea and convey non-motile sperms from rete testis into the epididymal head.
- **Epididymis (Head/Caput, Body/Corpus, Tail/Cauda):**
  - A tightly coiled, 6-meter-long tube lined with pseudostratified stereociliated columnar epithelium.
  - **Function:** Serves as the site for **physiological maturation, acquisition of progressive forward motility, membrane remodeling, and temporary storage** of spermatozoa. Sperm can survive in the cauda epididymis for up to a month.
- **Vas Deferens (Ductus Deferens — 45 cm):**
  - A muscular tube that ascends from the cauda epididymis through the **inguinal canal** into the pelvic cavity.
  - Loops over the ureter, expands into the **ampulla**, joins the duct of the seminal vesicle, and forms the **ejaculatory duct** (2 cm), which penetrates the prostate to open into the prostatic urethra.
- **Urethra (20 cm):** Common urogenital conduit divided into prostatic, membranous, and penile/spongy segments, terminating at the external **urethral meatus**.

Accessory Sex Glands & Semen Biochemistry

Seminal plasma is an alkaline, nutrient-dense fluid synthesized by three sets of accessory sex glands:

Accessory Gland	Volume %	Biochemical Composition & Physiological Significance
Seminal Vesicles (Paired, convoluted glands)	60–70%	Secretion is viscous, yellowish, and alkaline. Contains: (1) **Fructose** (exclusive metabolic substrate for sperm mitochondrial ATP generation); (2) **Prostaglandins** (stimulate smooth muscle reverse peristalsis in female uterus/oviducts); (3) **Seminogelin / Clotting Proteins** (coagulates semen post-ejaculation to prevent vaginal loss).
Prostate Gland (Single, walnut-sized)	20–30%	Encircles the urethra below the bladder. Secretes a thin, milky, slightly alkaline fluid (pH 7.3) rich in **citrate** (nutrient), **calcium**, **phosphate**, and proteolytic enzymes

Accessory Gland	Volume %	Biochemical Composition & Physiological Significance
		(**prostate-specific antigen, PSA**, and **fibrinolysin**) that liquify the coagulated semen clot within 15–20 minutes.
Bulbourethral Glands (Cowper's — Paired)	5%	Pea-sized glands located beneath the prostate. Prior to ejaculation, secrete an alkaline, clear, viscous mucus that coats the urethra to **neutralize residual acidic urine** and lubricates the glans penis.

- **Semen Composition:** Average ejaculate is **2.5 to 4 mL**, containing **200 to 300 million spermatozoa** suspended in alkaline seminal plasma (pH $7.2 - 7.7$).
- **WHO Fertility Benchmark:** At least **60% of sperms must exhibit normal morphology**, and at least **40% must demonstrate vigorous progressive motility**.
- **Clinical Anomalies: Oligospermia** (< 15 million sperms/mL), **Azoospermia** (complete absence of sperms), **Asthenozoospermia** (defective motility), and **Teratozoospermia** (abnormal morphology).

External Genitalia: The Penis

- **Erectile Columns:** Composed of three cylindrical vascular bodies bound by fibrous sheaths: two dorsolateral **corpora cavernosa** and one ventromedial **corpus spongiosum** enclosing the penile urethra and expanding distally into the sensitive **glans penis** (covered by retractable **prepuce** / foreskin).
- **Erection & Ejaculation:** Sexual arousal triggers parasympathetic **Nitric Oxide (NO)** release, elevating cGMP to relax arteriolar smooth muscle; blood engorges cavernous spaces producing rigid erection. Ejaculation is a sympathetic spinal reflex expelling semen through the urethral meatus.

Module 2: The Female Reproductive System & Ovarian Histology

The female reproductive system generates ova, receives spermatozoa, provides the fertilization site, sustains 9 months of gestation, and delivers and nourishes the infant.

Ovarian Architecture & Follicular Development

- **Ovaries:** Paired primary sex organs (length 2–4 cm) anchored by the **suspensory ligament** (to pelvic wall) and **ovarian ligament** (to uterus). Covered by cuboidal **germinal epithelium** and fibrous **tunica albuginea**.
- **Ovarian Stroma:** Outer **cortex** (houses ovarian follicles) and inner **medulla** (blood vessels, lymphatics, nerves).
- **Follicular Stratigraphy:**
  - **Primordial Follicle:** Primary oocyte ($2n$, arrested at diplotene) surrounded by a single layer of squamous follicular cells.
  - **Primary Follicle:** Granulosa cells become stratified cuboidal; **zona pellucida** (glycoprotein coat) begins forming.
  - **Secondary Follicle:** Multiple granulosa layers enclosed by an organizing stromal **theca**.
  - **Tertiary Follicle:** Characterized by a fluid-filled cavity (**Antrum**). Theca differentiates into an angiogenic, steroidogenic **theca interna** (synthesizes androgens aromatized to estrogen by granulosa cells) and a fibrous **theca externa**. Primary oocyte completes Meiosis I to form the **Secondary Oocyte (n)** and First Polar Body.

- **Mature Graafian Follicle (18–25 mm):** Large antrum; secondary oocyte suspended on a granulosa mound (**cumulus oophorus**).

Female Accessory Ducts: Fallopian Tubes & Uterus

- **Fallopian Tubes (Oviducts — 10 to 12 cm):** Ciliated simple columnar epithelium with secretory peg cells:
  - **1. Fimbriae:** Finger-like margins that sweep the ovarian surface to catch the ovulated secondary oocyte.
  - **2. Infundibulum:** Funnel-shaped conduit bearing the abdominal ostium.
  - **3. Ampulla:** Widest, longest segment (approx. 7 cm). **The ampullary-isthmic junction is the definitive biological site of fertilization.**
  - **4. Isthmus:** Narrow, thick-walled muscular conduit opening into the uterine cornu.
- **The Uterus (Womb):** Inverted pear-shaped organ (fundus, body, cervix) with a three-layered wall:
  - **Perimetrium:** Outer serous peritoneum.
  - **Myometrium:** Thick interlaced smooth muscle; undergoes hypertrophy in pregnancy and contracts vigorously during labour under **Oxytocin**.
  - **Endometrium:** Inner vascular mucosal coat. Permanent **stratum basale** regenerates the superficial **stratum functionale**, which proliferates cyclically and sloughs off during menstruation.
- **Cervix & Vagina:** Cervical canal + vagina constitutes the **birth canal**. Vagina (8–10 cm) is lined by non-keratinized stratified squamous epithelium; commensal **Lactobacillus acidophilus** ferment shed glycogen into lactic acid (pH $3.8 - 4.5$), creating an acidic antimicrobial barrier.

External Genitalia (Vulva) & Mammary Glands

- **Vulval Structures & Male Homologies:**
  - **Mons Pubis:** Adipose cushion over pubic symphysis covered with pubic hair.
  - **Labia Majora:** Paired fleshy cutaneous folds (**homologous to male scrotum**).
  - **Labia Minora:** Paired delicate mucous folds medial to labia majora (**homologous to penile urethra skin**).
  - **Clitoris:** Sensitive erectile cylinder at anterior junction of labia minora (**homologous to glans penis**).
  - **Hymen:** Vascular mucous membrane partially occluding the vaginal introitus; torn by coitus, sports, or physical activity; **unreliable index of virginity**.
  - **Bartholin's Glands:** Greater vestibular glands (**homologous to bulbourethral glands**); secrete lubricating mucus.
- **Mammary Glands:** Modified sweat glands with 15–20 lobes composed of milk-secreting **alveoli**.
  - **Milk Ejection Route:** Alveoli → Mammary Tubules → Mammary Ducts → Ampulla → Lactiferous Ducts → Nipple.

Module 3: Gametogenesis: Spermatogenesis vs Oogenesis

Gametogenesis is the production of haploid gametes (spermatozoa and ova) from diploid primordial germ cells through meiotic reduction:




Figure 1: Comparative cytogenetic architecture, meiotic progression, hormonal feedback axes, and morphological organization of spermatogenesis versus oogenesis.

Spermatogenesis Cytogenetics & Hormonal Control

- **Initiation:** Initiated strictly at puberty (age 13–14) in response to pulsatile hypothalamic **Gonadotropin-Releasing Hormone (GnRH)**.
- **Cytogenetic Sequence:**
  - **Multiplication Phase:** Spermatogonia ($2n, 46$ chromosomes) situated on the basement membrane divide mitotically. **Type A spermatogonia** remain as stem cells, while **Type B spermatogonia** differentiate and enlarge into **Primary Spermatocytes ($2n, 46$)**.
  - **Growth & Meiosis I:** Primary spermatocytes duplicate their DNA (46 chromosomes, 92 chromatids, $4C$ DNA content) and undergo a prolonged prophase I (pachytene crossing over). Completion of Meiosis I yields two equal haploid **Secondary Spermatocytes ($n, 23$ chromosomes, 46 chromatids, $2C$ DNA content)**.
  - **Meiosis II:** Secondary spermatocytes immediately undergo equational Meiosis II without interkinesis, dividing centromeres to produce four uniflagellate haploid **Spermatids ($n, 23$ chromosomes, 23 chromatids, $1C$ DNA content)**.
- **Spermiogenesis & Spermiation:**
  - **Spermiogenesis:** The cellular metamorphosis of spherical, non-motile spermatids into functional spermatozoa: (1) Golgi apparatus condenses into the acrosomal cap; (2) nucleus elongates and dehydrates; (3) centrioles align; (4) mitochondria spiral around axial filament forming the **Nebenkern**; (5) excess cytoplasm is shed as a residual body.
  - **Spermiation:** The release of finished spermatozoa from Sertoli cell crypts into the seminiferous tubule lumen.
- **Endocrine Circuit:**

$$\text{Hypothalamus (GnRH)} \rightarrow \text{Anterior Pituitary} \begin{cases} \rightarrow \text{LH} \rightarrow \text{Leydig Cells} \rightarrow \text{Testosterone} \rightarrow \text{Spermatogenesis} \\ \rightarrow \text{FSH} \rightarrow \text{Sertoli Cells} \rightarrow \text{ABP \& Factors} \rightarrow \text{Spermiogenesis} \end{cases}$$

  - **Feedback:** High testosterone suppresses hypothalamic GnRH and pituitary LH; Sertoli cells secrete **Inhibin**, which selectively suppresses pituitary FSH synthesis.

Oogenesis Cytogenetics & The Dual Meiotic Arrests

Oogenesis is initiated during embryonic development, ceases postnatally, resumes cyclically at puberty, and terminates permanently at menopause (50 years):

- **Prenatal Phase:** Primordial germ cells migrate from yolk sac endoderm into fetal genital ridges, multiplying mitotically to form approximately **2 million oogonia $(2n, 46)$** per ovary by the 20th week of gestation. **Zero oogonia are formed or added after birth.**
- **Arrest I (Prophase I — Diplotene):** All oogonia enter Meiosis I and arrest at the **diplotene stage of Prophase I**, becoming **primary oocytes $(2n, 46)$** surrounded by pre-granulosa cells (**primordial follicles**).
- **Follicular Atresia:** Most follicles degenerate (**atresia**). By birth, count falls to 1 million; by puberty (menarche), only **60,000 to 80,000 primary follicles** survive in each ovary.
- **Pubertal Resumption:** Each month, FSH stimulates 15–20 primary follicles. A single dominant tertiary follicle matures into a Graafian follicle.
- **Meiosis I Completion:** Hours prior to ovulation, the primary oocyte completes Meiosis I.
  - **Extreme Asymmetric Cytokinesis:** Cleavage furrow forms at the extreme cell periphery, allocating almost all cytoplasm, yolk, glycogen, and maternal organelles to the **Secondary Oocyte $(n, 23)$**, while extruding a tiny nucleus with negligible cytoplasm as the **First Polar Body $(n, 23)$**.
- **Arrest II (Metaphase II) & Ovulation:** The secondary oocyte begins Meiosis II but arrests at **Metaphase II** due to activity of **Cytostatic Factor (CSF)**. **Ovulation discharges the secondary oocyte arrested at Metaphase II.**
- **Meiosis II Completion:** When a fertilizing sperm penetrates the plasma membrane, an intracellular calcium surge degrades CSF and MPF. The secondary oocyte rapidly completes Meiosis II, extruding the **Second Polar Body (n)** and leaving the mature haploid **Ovum (Ootid, n)**.

Morphological Architecture of Mature Gametes

Parameter	Spermatozoon (Male Gamete)	Secondary Oocyte / Ovum (Female Gamete)
Dimensions	Slender, microscopic (length 60 μm, width 3 μm)	Enormous, spherical, non-motile (diameter 100–120 μm)
Motility	Highly motile; flagellar propulsion speed 1.5–3 mm/min	Completely non-motile; propelled passively by oviduct ciliary beat
Nutritional Reserves	Dehydrated, no food reserves; relies on seminal fructose	Enriched with maternal cytoplasm, enzymes, lipids, and glycogen
Envelopes	Covered by continuous plasma membrane	Enclosed by plasma membrane, **Zona Pellucida**, and **Corona Radiata**
Centrioles	Possesses proximal and distal centrioles	Centrosome lost during meiotic divisions; lacks centrioles
Output per MMC	1 primary spermatocyte → 4 viable spermatozoa	1 primary oocyte → 1 functional ovum + 2–3 polar bodies

Module 4: Menstrual Cycle Endocrinology & Uterine Dynamics

The reproductive cycle in non-pregnant female primates (humans, apes, Old World monkeys) is termed the **menstrual cycle**. Its average duration is **28 days** (ranging from 25 to 35 days).




Figure 2: Synchronized 28-day endocrine orchestration of the human menstrual cycle: pituitary gonadotropins (LH, FSH), ovarian steroids (estrogens, progesterone), and endometrial structural remodeling.

The Four Phases of the 28-Day Cycle

- **1. Menstrual Phase (Days 1 to 5):**
  - Initiated when fertilization does not occur. Progesterone and estrogen levels crash due to regression of the corpus luteum.
  - **Vascular Ischemia:** Sudden withdrawal of progesterone triggers sharp spasmodic contractions of endometrial spiral arterioles, causing ischemic tissue necrosis.
  - **Menstrual Flow:** The necrotic stratum functionale sloughs off, discharging approximately 40–80 mL of blood, glandular mucus, and unfertilized secondary oocyte debris through the vagina.
- **2. Follicular / Proliferative Phase (Days 6 to 13):**
  - Hypothalamic GnRH stimulates anterior pituitary **FSH** and **LH** secretion.
  - FSH recruits and matures a cohort of ovarian follicles (Primary → Secondary → Tertiary → Graafian).
  - Growing granulosa and theca interna cells secrete escalating quantities of **Estrogens**.
  - **Endometrial Repair:** High estrogen stimulates rapid mitotic proliferation of uterine stroma, blood vessels, and tubular glands, restoring endometrial thickness to 3–5 mm. Cervical mucus becomes clear, watery, and alkaline to facilitate sperm migration.
- **3. Ovulatory Phase (Day 14 — Mid-Cycle Peak):**
  - Estrogen levels surpass a critical physiological threshold, triggering a neuroendocrine switch from negative feedback to **positive feedback** on the pituitary.
  - **LH Surge:** A massive, sharp spike in LH secretion (accompanied by a smaller FSH surge) reaches peak levels at mid-cycle.
  - **Ovulation:** The LH surge induces proteolytic enzymes that dissolve the follicular wall. The mature Graafian follicle ruptures, discharging the **Secondary Oocyte (*n*, arrested at Metaphase II)** into the peritoneal cavity. Basal body temperature rises by 0.3° C to 0.5° C due to early progesterone synthesis.
- **4. Luteal / Secretory Phase (Days 15 to 28 — Invariable 14 Days):**
  - Under LH influence, the collapsed remnant of the ruptured Graafian follicle transforms into a bright yellow endocrine body called the **Corpus Luteum**.
  - **Progesterone Dominance:** Corpus luteum secretes immense amounts of **Progesterone** (the hormone of pregnancy) and moderate estrogen.

- **Secretory Transformation:** Endometrium reaches maximum thickness (6–8 mm); uterine glands become highly tortuous, coiled, and secrete a glycogen-rich fluid ("uterine milk") to nourish the impending blastocyst.
- **Dual Fate Rule:**
  - **If Unfertilized:** High progesterone exerts strong negative feedback inhibiting pituitary LH; lack of LH causes corpus luteum to degenerate into a fibrous scar called the **Corpus Albicans**. Progesterone plunges, triggering the next menstrual cycle.
  - **If Fertilized:** The implanting blastocyst secretes **hCG**, rescuing the corpus luteum and maintaining high progesterone to prevent menstruation.

WORKED EXAMPLE — MENSTRUAL CYCLE CYCLE CALCULATIONS

Clinical Scenario: A female patient exhibits an irregular menstrual cycle averaging **34 days**. On which day of her cycle is ovulation expected to occur, and what is her fertile window?

Step-by-Step Physiological Calculation:

- **Rule of Fixed Luteal Phase:** In human females, the post-ovulatory luteal (secretory) phase has a strictly fixed biological duration of **14 days** across all cycle lengths. Variations in cycle length are due entirely to changes in the follicular/proliferative phase.
- **Ovulation Day Calculation:**

$$\text{Ovulation Day} = \text{Total Cycle Length} - 14 \text{ days} = 34 - 14 = \text{Day } 20$$

- **Fertile Window:** Secondary oocyte survives 24 hours post-ovulation; viable sperms persist up to 72 hours in the female tract. Therefore, the fertile window spans **Day 17 to Day 21**.

Module 5: Fertilisation, Cleavage & Implantation

- **Insemination & Transport:** During coitus, 200 to 300 million sperms are deposited in the vagina. Sperms swim through the cervix and uterus into the fallopian tube within minutes, assisted by uterine muscular contractions and their own flagellar propulsion.
- **Capacitation:** Freshly ejaculated sperms cannot fertilize an ovum. Over 5 to 7 hours in the female reproductive tract, sperms undergo **capacitation**—removal of inhibitory cholesterol and glycoproteins from the acrosomal membrane and influx of Ca^{2+}, hyperactivating flagellar motility.
- **Fertilisation Locus:** Fertilization occurs strictly at the **Ampullary-isthmic junction (Ampulla)** of the fallopian tube.

The Acrosome Reaction & Prevention of Polyspermy

1. **Acrosome Reaction:** When the capacitated sperm contacts the ovum, it binds to specific **ZP3 glycoprotein receptors** on the zona pellucida. This triggers exocytosis of the acrosomal vesicle, releasing **hyaluronidase** (digests corona radiata cells) and **acrosin** (digests a passage through the zona pellucida).
2. **Fast Block to Polyspermy:** The binding of the first sperm membrane triggers immediate sodium influx, causing **rapid electrical depolarization of the oocyte plasma membrane** within 2–3 seconds, transiently preventing further sperm fusions.
3. **Slow Block (Cortical Reaction — Permanent):** Fusion triggers intracellular calcium wave release, causing thousands of **cortical granules** beneath the oocyte membrane to exocytose enzymes into the perivitelline space. These enzymes hydrolyze ZP3 receptors and cross-link glycoproteins (**zona hardening**), permanently barring any additional sperm from entering.

4. **Syngamy:** The sperm nucleus, proximal centriole, and mitochondria enter the oocyte cytoplasm. Sperm entry induces the secondary oocyte to complete Meiosis II, forming the haploid **female pronucleus** and the **Second Polar Body**. The male and female pronuclei fuse (**karyogamy / syngamy**), restoring the diploid condition: the **Zygote ($2n, 46$)**.

Cleavage & Blastocyst Morphogenesis

Within 24 hours of fertilization, the zygote undergoes rapid mitotic divisions called **cleavage** as it is propelled along the fallopian tube towards the uterus by ciliary currents:

- **Holoblastic Cleavage:** Human cleavage is holoblastic (divides entire zygote) and slightly unequal.
  - 1 cell ($2n$) $\rightarrow$ 2 cells $\rightarrow$ 4 cells $\rightarrow$ 8 cells $\rightarrow$ 16 cells (**Morula**).
  - **Morula:** A solid, mulberry-like ball of 16 blastomeres (Days 3–4 post-fertilization), encased firmly within the intact zona pellucida to prevent ectopic tubal implantation.
- **Blastocyst Formation:**
  - On Day 4–5, fluid enters the morula forming a central cavity (**blastocoel**). The embryo is now a **Blastocyst**.
  - **Trophoblast:** An outer single layer of flattened epithelial cells; later develops into chorionic villi and contributes to the fetal placenta.
  - **Inner Cell Mass (ICM / Embryoblast):** A cluster of cells attached to one pole of the trophoblast; represents the true progenitor of the embryo proper. ICM cells are **pluripotent stem cells** capable of differentiating into all body tissues.
- **Hatching & Implantation:**
  - By Day 6–7, the blastocyst reaches the uterine cavity and sheds its zona pellucida (**hatching**).
  - **Implantation:** The trophoblast cells secrete lytic enzymes that digest uterine endometrial stroma, allowing the blastocyst to embed deeply within the vascular endometrium. Implantation is completed by Day 11–12.

Module 6: Gestation, Placenta, Parturition & Lactation

- **Placenta Formation:** Following implantation, finger-like projections called **chorionic villi** sprout from the trophoblast and interdigitate intimately with maternal endometrial crypts, forming the **Placenta**—a joint structural and functional vascular bridge between fetus and mother. Connected to embryo by the **umbilical cord** (houses 2 umbilical arteries and 1 umbilical vein).
- **Endocrine Placenta:** The placenta acts as a temporary, vital endocrine gland secreting:
  - **hCG (Human Chorionic Gonadotropin):** Rescues corpus luteum to sustain progesterone; excreted in maternal urine (diagnostic marker in pregnancy test kits).
  - **hPL (Human Placental Lactogen):** Modulates maternal metabolism and stimulates breast development.
  - **Estrogens & Progestogens:** Maintain uterine endometrium and suppress maternal gonadotropins.
  - **Relaxin:** Peptide hormone secreted by ovary and placenta in late pregnancy to soften pubic symphysis.
  - **CUET High-Yield Trap: hCG, hPL, and Relaxin are synthesized in females ONLY during pregnancy.**

- **Embryonic Germ Layers & Stem Cells:** Inner cell mass differentiates into three primary germ layers: outer **Ectoderm**, middle **Mesoderm**, and inner **Endoderm**, which collectively give rise to all adult tissues and organs. ICM contains **pluripotent stem cells**.
- **Timeline of Embryonic Milestones (High-Yield NCERT Sequence):**
  - **1st Month:** Embryonic heart forms (first clinical sign: detection of fetal heartbeat via stethoscope).
  - **2nd Month:** Limbs and digits differentiate; primary organogenesis initiates.
  - **12 Weeks (End of 1st Trimester):** Major organ systems established; external genitalia clearly defined.
  - **5th Month: First fetal movements (Quickening)** observed by mother; fine hair appears on head.
  - **24 Weeks (End of 2nd Trimester):** Body covered with fine hair (**lanugo**); eyelids separate; eyelashes form.
  - **9 Months:** Full-term maturity finished; fetus turns cephalic (head-down), prepared for birth.
- **Parturition (Childbirth) & Ferguson Reflex:** Average human gestation period is 9 months (approx. 280 days).
  - **Fetal Ejection Reflex:** Fully mature fetus and placenta initiate mild neuroendocrine signals that induce early contractions.
  - **Oxytocin Positive Feedback Loop:** Impulses travel to maternal posterior pituitary, releasing **Oxytocin**. Oxytocin stimulates powerful myometrial contractions; contractions press fetal head against cervix, sending sensory impulses for increased oxytocin release, generating an escalating positive feedback loop culminating in expulsion of the infant.
- **Lactation & Colostrum:** Mammary alveoli synthesize milk under anterior pituitary **Prolactin**. Milk letdown (ejection) during suckling is mediated by posterior pituitary **Oxytocin**.
  - **Colostrum:** The yellowish, watery milk secreted during initial days post-birth. Highly rich in proteins, minerals, and maternal **secretory IgA antibodies**, providing critical passive immunity to protect the newborn infant against gastrointestinal infections.

High-Yield Formula & Concept Cheatsheet

Concept / Structure	Formula / Ratio	Essential Exam Memory Trigger
Spermatogenesis Output	1 Primary → 4 Sperms	1 primary spermatocyte ($2n$) yields 4 functional motile spermatozoa (n).
Oogenesis Output	1 Primary → 1 Ovum	1 primary oocyte ($2n$) yields only 1 functional ovum (n) + 2–3 polar bodies.
Meiotic Arrest 1	Prophase I (Diplotene)	Initiated prenatally; primary oocytes remain arrested here until puberty.
Meiotic Arrest 2	Metaphase II	Secondary oocyte ovulated in Metaphase II; completed **only upon sperm entry**.
LH Surge Target	Day 14 (Mid-Cycle)	Triggers rupture of Graafian follicle and release of secondary oocyte.
Fertilization Site	Ampulla (Oviduct)	Specific locus at ampullary-isthmic junction; cleavage begins inside oviduct.
Pregnancy Hormones	hCG, hPL, Relaxin	Produced in female body **exclusively** during pregnancy; absent in non-pregnant state.
Milk Reflex Dual Axis	Prolactin vs Oxytocin	Prolactin synthesizes milk; Oxytocin ejects milk ("milk letdown reflex").




Figure 3: Master synthesis mindmap of Human Reproduction: coordinating male/female anatomy, gametogenesis, menstrual hormonal dynamics, fertilization, embryogeny, and parturition.

Practice Questions — Part 1 (Q1 to Q5)

Q1. In human males, the testes are situated outside the abdominal cavity within the scrotal sac primarily because:

(A) The scrotal position protects testes from abdominal hydrostatic pressure
(B) Spermatogenesis requires an optimal temperature 2°C to 2.5°C lower than core internal body temperature
(C) The scrotum provides an increased volume of interstitial tissue for androgen storage
(D) Testicular descent facilitates direct drainage of seminal fluid into the urinary bladder

Sol. Q1. Normal human spermatogenesis is heat-sensitive and cannot proceed at internal body core temperature (37° C). The scrotum functions as a physical thermoregulator, maintaining the testes at 34.5° C (2° C to 2.5° C below core temperature). Undescended testes (**cryptorchidism**) result in sterility due to germ cell degeneration.
Answer: (B)

Q2. Which pituitary gonadotropin directly stimulates Leydig cells to synthesize androgens, and which hormone acts on Sertoli cells to support spermiogenesis?

(A) FSH acts on Leydig cells; LH acts on Sertoli cells
(B) LH acts on Leydig cells; FSH acts on Sertoli cells
(C) Prolactin acts on Leydig cells; Oxytocin acts on Sertoli cells
(D) GnRH acts on Leydig cells; Inhibin acts on Sertoli cells

Sol. Q2. Pituitary Luteinizing Hormone (LH / ICSH) stimulates interstitial Leydig cells to synthesize and secrete androgens (principally testosterone). Follicle Stimulating Hormone (FSH) acts directly on Sertoli cells, stimulating them to secrete androgen-binding protein (ABP) and factors essential for spermiogenesis. **Answer: (B)**

Q3. What is the specific contribution of the proximal centriole brought into the egg by the fertilizing sperm?

(A) It provides the energy-generating mitochondrial sheath for early embryonic motility
(B) It organizes the first mitotic cleavage spindle apparatus in the centriole-deficient ovum
(C) It triggers the cortical reaction by enzymatically hydrolyzing the glycoprotein matrix
(D) It directly induces transcription of embryonic paternal genome alleles

Sol. Q3. The unfertilized human ovum loses its centrosome during the second meiotic division and lacks a functional centriole. The penetrating spermatozoon contributes its proximal centriole, which organizes the pericentriolar material into the spindle apparatus necessary for the first mitotic cleavage of the diploid zygote.
Answer: (B)

Q4. During human oogenesis, at which precise cytological stage does the primary oocyte arrest from fetal life until puberty, and at which stage is the secondary oocyte arrested at ovulation?

(A) Arrest 1: Metaphase I; Arrest 2: Telophase II
(B) Arrest 1: Diplotene of Prophase I; Arrest 2: Metaphase II
(C) Arrest 1: Pachytene of Prophase I; Arrest 2: Anaphase I
(D) Arrest 1: Leptotene of Prophase I; Arrest 2: Prophase II

Sol. Q4. Oogenesis features two prominent developmental arrests: (1) Primary oocytes arrest prenatally at the **diplotene stage of Prophase I** within primordial follicles until puberty; (2) Upon completing Meiosis I prior to ovulation, the secondary oocyte enters Meiosis II and arrests at **Metaphase II**, which is released at ovulation.
Answer: (B)

Q5. The formation of a fluid-filled cavity called the antrum is the distinctive histological hallmark of which ovarian follicular stage?

(A) Primordial follicle
(B) Primary follicle
(C) Secondary follicle
(D) Tertiary follicle

Sol. Q5. The secondary follicle transforms into a tertiary follicle upon the appearance of a fluid-filled follicular cavity called the **Antrum**. The tertiary follicle is also characterized by the differentiation of the surrounding theca into an inner glandular theca interna and an outer fibrous theca externa. **Answer: (D)**

Practice Questions — Part 2 (Q6 to Q10)

Q6. The LH surge that triggers ovulation in a normal 28-day human menstrual cycle occurs precisely at which time point and under what hormonal stimulus?

(A) Day 1 to 5, stimulated by high progesterone from the corpus luteum
(B) Day 14 (mid-cycle), induced by positive feedback from peak estrogen levels
(C) Day 21, stimulated by low levels of both gonadotropins
(D) Day 28, triggered by ischemic necrosis of the stratum basale

Sol. Q6. During the late follicular phase, peak estrogen secreted by the mature Graafian follicle surpasses a critical threshold, triggering positive feedback on the anterior pituitary. This causes a massive, rapid release of LH (**LH surge**) at mid-cycle (Day 14), which induces rupture of the Graafian follicle and ovulation. **Answer: (B)**

Q7. What is the endocrine function of the corpus luteum during the post-ovulatory luteal phase, and what causes its degeneration in the absence of pregnancy?

(A) It secretes massive quantities of FSH; degenerates due to elevated oxytocin levels
(B) It secretes large amounts of progesterone to maintain the endometrium; degenerates due to falling LH support
(C) It secretes relaxin to trigger menstruation; degenerates due to estrogen feedback
(D) It secretes prolactin to stimulate milk ducts; degenerates due to falling body temperature

Sol. Q7. The corpus luteum secretes large amounts of **Progesterone** (and some estrogen), which transforms the endometrium into a glandular, secretory bed. In the absence of fertilization, high progesterone suppresses pituitary LH. Lacking LH trophic support, the corpus luteum regresses into a fibrous **corpus albicans**. **Answer: (B)**

Q8. In the human female reproductive tract, fertilization of the secondary oocyte by a viable sperm occurs definitively at which anatomical location?

(A) Cervical canal
(B) Uterine fundus
(C) Ampullary region of the fallopian tube
(D) Infundibular fimbriae

Sol. Q8. Fertilization in humans takes place specifically within the **ampulla (ampullary-isthmic junction)** of the fallopian tube (oviduct). The ovulated secondary oocyte is swept by fimbriae into the infundibulum and conveyed into the ampulla, where it meets capacitated sperms. **Answer: (C)**

Q9. The permanent, irreversible block to polyspermy in humans is achieved primarily through which physiological mechanism?

(A) Rapid hyperpolarization of the corona radiata cells by sodium ions
(B) Depolarization of the sperm middle piece by mitochondrial calcium release
(C) Exocytosis of cortical granules that enzymatically alter and harden the zona pellucida
(D) Mechanical contraction of the myometrium expelling excess sperms

Sol. Q9. While membrane depolarization provides a transient fast block, the permanent slow block to polyspermy is the **cortical reaction**. Sperm-egg fusion triggers intracellular Ca^{2+} release, causing oocyte cortical granules to exocytose enzymes into the perivitelline space, destroying sperm receptors and hardening the zona pellucida. **Answer: (C)**

Q10. What biological event directly triggers the arrested secondary oocyte to resume and complete its second meiotic division?

(A) Contact of the sperm flagellum with cervical mucus
(B) Binding of LH to luteal receptors on the corona radiata
(C) Penetration and entry of the sperm into the oocyte cytoplasm
(D) Implantation of the morula into the endometrial stroma

Sol. Q10. The secondary oocyte is arrested at Metaphase II. Entry of the fertilizing sperm provides the calcium stimulus that degrades Maturation Promoting Factor (MPF), prompting the oocyte to complete Meiosis II. This unequal division extrudes the **Second Polar Body** and yields the haploid female pronucleus (**Ovum**). **Answer: (C)**

Practice Questions — Part 3 (Q11 to Q15)

Q11. In a developing human blastocyst prior to implantation, what are the respective developmental fates of the trophoblast and the inner cell mass?

(A) Trophoblast forms the nervous system; inner cell mass forms the placenta
(B) Trophoblast attaches to endometrium and forms chorionic villi; inner cell mass forms the embryo proper
(C) Trophoblast forms the amnion; inner cell mass undergoes complete apoptosis
(D) Trophoblast develops into the umbilical cord; inner cell mass forms maternal decidua

Sol. Q11. A blastocyst consists of an outer epithelial layer, the **Trophoblast**, and an internal cluster of cells, the **Inner Cell Mass (ICM)**. The trophoblast attaches to the endometrium, digests uterine stroma, and forms fetal chorionic villi of the placenta. The pluripotent cells of the ICM differentiate into all tissues of the embryo proper. **Answer: (B)**

Q12. Which trio of hormones is synthesized in the human female body exclusively during the state of pregnancy?

(A) Estrogen, Progesterone, and Oxytocin
(B) hCG, hPL, and Relaxin
(C) FSH, LH, and Prolactin
(D) Cortisol, Thyroxine, and Aldosterone

Sol. Q12. Human Chorionic Gonadotropin (**hCG**), Human Placental Lactogen (**hPL**), and **Relaxin** (produced by ovary/placenta in late pregnancy) are synthesized **exclusively during pregnancy**. Other hormones like estrogen and progesterone circulate in non-pregnant cycling females as well. **Answer: (B)**

Q13. The neuroendocrine reflex that triggers the onset of labour (parturition) in humans is termed the:

(A) Milk ejection reflex, mediated by anterior pituitary prolactin
(B) Fetal ejection reflex, initiated by the fully developed fetus and placenta
(C) Baroreceptor reflex, triggered by placental vasoconstriction
(D) Luteal regression reflex, stimulated by sudden estrogen surge

Sol. Q13. Parturition is induced by a complex neuroendocrine mechanism called the **fetal ejection reflex**. Signals originating from the fully developed fetus and placenta induce mild uterine contractions. This stimulates the maternal posterior pituitary to secrete **Oxytocin**, initiating a positive feedback cascade of vigorous contractions. **Answer: (B)**

Q14. During lactation, which hormone is primarily responsible for milk synthesis within mammary alveoli, and which hormone mediates milk ejection (let-down)?

(A) Oxytocin stimulates synthesis; Prolactin stimulates ejection
(B) Prolactin stimulates synthesis; Oxytocin stimulates ejection
(C) Estrogen stimulates synthesis; Progesterone stimulates ejection
(D) hPL stimulates synthesis; Relaxin stimulates ejection

Sol. Q14. Anterior pituitary **Prolactin** stimulates the glandular alveoli of the mammary glands to synthesize and secrete milk. Posterior pituitary **Oxytocin** contracts the myoepithelial cells surrounding the alveoli, causing the physical ejection or "let-down" of milk in response to infant suckling. **Answer: (B)**

Q15. Why is the initial yellowish milk (**colostrum**) secreted by the mother during the first few days postpartum deemed indispensable for the newborn infant?

(A) It contains exceptionally high concentrations of lipids for adipose insulation
(B) It is packed with maternal secretory IgA antibodies that confer essential passive immunity
(C) It provides synthetic growth hormones that induce rapid epiphyseal closure
(D) It contains clotting factors that permanently occlude the ductus arteriosus

Sol. Q15. **Colostrum** is rich in proteins, vitamins, and maternal **secretory IgA (sIgA) antibodies**. Because the newborn's immune system is naive, these ingested IgA antibodies coat the infant's mucosal surfaces, providing critical **passive immunity** against gastrointestinal and respiratory pathogens. **Answer: (B)**

© OurMocks · www.ourmocks.com · CUET UG Biology · Human Reproduction · 2027 Cycle