

WIDE-FIELD EFFECTS IN REDSHIFTED 21 CM POWER SPECTRA

NITHYANANDAN THYAGARAJAN^{1*}, DANIEL C. JACOBS¹, JUDD D. BOWMAN¹, N. BARRY², A. P. BEARDSLEY²,
 G. BERNARDI^{3,4,5}, F. BRIGGS^{6,7}, R. J. CAPPALLO⁸, P. CARROLL², B. E. COREY⁸, A. DE OLIVEIRA-COSTA⁹,
 JOSHUA S. DILLON⁹, D. EMRICH¹⁰, A. EWALL-WICE⁹, L. FENG⁹, R. GOEKE⁹, L. J. GREENHILL⁵, B. J. HAZELTON²,
 J. N. HEWITT⁹, N. HURLEY-WALKER¹⁰, M. JOHNSTON-HOLLITT¹¹, D. L. KAPLAN¹², J. C. KASPER^{13,5}, HAN-SEEK KIM^{14,7},
 P. KITTIWISIT¹, E. KRATZENBERG⁸, E. LENC^{15,7}, J. LINE^{14,7}, A. LOEB⁵, C. J. LONSDALE⁸, M. J. LYNCH¹⁰,
 B. MCKINLEY^{14,7}, S. R. MCWHIRTER⁸, D. A. MITCHELL^{16,7}, M. F. MORALES², E. MORGAN⁹, A. R. NEBEN⁹, D. OBEROI¹⁷,
 A. R. OFFRINGA^{6,7}, S. M. ORD^{10,7}, SOURABH PAUL¹⁸, B. PINDOR^{14,7}, J. C. POBER², T. PRABU¹⁸, P. PROCOPIO^{14,7},
 J. RIDING^{14,7}, A. E. E. ROGERS⁸, A. ROSHI¹⁹, N. UDAYA SHANKAR¹⁸, SHIV K. SETHI¹⁸, K. S. SRIVANI¹⁸,
 R. SUBRAHMANYAN^{18,7}, I. S. SULLIVAN², M. TEGMARK⁹, S. J. TINGAY^{10,7}, C. M. TROTT^{10,7}, M. WATERSON^{10,6},
 R. B. WAYTH^{10,7}, R. L. WEBSTER^{14,7}, A. R. WHITNEY⁸, A. WILLIAMS¹⁰, C. L. WILLIAMS⁹, C. WU²⁰, J. S. B. WYTHE^{14,7}

Draft version March 12, 2015

ABSTRACT

Foreground emission is currently the primary limitation to detection of redshifted H I emission from the epoch of reionization. Modern radio telescopes that target this cosmological signal are typically wide-field instruments. Through modeling of delay spectra measured between antenna pairs, it has recently emerged that wide-field measurements imprint a characteristic *pitchfork*-shaped signature in this Fourier domain. It is characterized by enhanced power from foreground emission mapped to regions near the horizon and plays a significant role in determining the contamination of the cosmological H I signal. With MWA data sensitivity improved by coherently averaging snapshots aligned in local sidereal time across different observing nights, we confirm the prediction from modeling at $> 10\sigma$ level.

Subject headings: cosmology: observations — dark ages, reionization, first stars — large-scale structure of universe — methods: statistical — radio continuum: galaxies — techniques: interferometric

¹ Arizona State University, School of Earth and Space Exploration, Tempe, AZ 85287, USA

² University of Washington, Department of Physics, Seattle, WA 98195, USA

³ Square Kilometre Array South Africa (SKA SA), Park Road, Pinelands 7405, South Africa

⁴ Department of Physics and Electronics, Rhodes University, Grahamstown 6140, South Africa

⁵ Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138, USA

⁶ Australian National University, Research School of Astronomy and Astrophysics, Canberra, ACT 2611, Australia

⁷ ARC Centre of Excellence for All-sky Astrophysics (CAASTRO)

⁸ MIT Haystack Observatory, Westford, MA 01886, USA

⁹ MIT Kavli Institute for Astrophysics and Space Research, Cambridge, MA 02139, USA

¹⁰ International Centre for Radio Astronomy Research, Curtin University, Perth, WA 6845, Australia

¹¹ Victoria University of Wellington, School of Chemical & Physical Sciences, Wellington 6140, New Zealand

¹² University of Wisconsin–Milwaukee, Department of Physics, Milwaukee, WI 53201, USA

¹³ University of Michigan, Department of Atmospheric, Oceanic and Space Sciences, Ann Arbor, MI 48109, USA

¹⁴ The University of Melbourne, School of Physics, Parkville, VIC 3010, Australia

¹⁵ The University of Sydney, Sydney Institute for Astronomy, School of Physics, NSW 2006, Australia

¹⁶ CSIRO Astronomy and Space Science (CASS), PO Box 76, Epping, NSW 1710, Australia

¹⁷ National Centre for Radio Astrophysics, Tata Institute for Fundamental Research, Pune 411007, India

¹⁸ Raman Research Institute, Bangalore 560080, India

¹⁹ National Radio Astronomy Observatory, Charlottesville and Greenbank, USA

²⁰ International Centre for Radio Astronomy Research, University of Western Australia, Crawley, WA 6009, Australia

* e-mail: t_nithyanandan@asu.edu

1. INTRODUCTION

2. WIDE-FIELD EFFECTS IN DELAY SPECTRUM

Thyagarajan et al. (2015) have described in detail the effects of wide-field measurements as seen in the delay spectra of interferometer *visibilities*. Here, we give a brief overview of the wide-field signature predicted therein.

The delay spectrum for a baseline vector, \mathbf{b} , is given by (Parsons et al. 2012a,b; Thyagarajan et al. 2013, 2015):

$$\tilde{V}_b(\tau) \equiv \int V_b(f) W(f) e^{i2\pi f \tau} df, \quad (1)$$

with interferometer visibilities, $V_b(f)$, given by (van Cittert 1934; Zernike 1938; Thompson et al. 2001):

$$V_b(f) = \iint_{\text{sky}} A(\hat{\mathbf{s}}, f) I(\hat{\mathbf{s}}, f) W_i(f) e^{-i2\pi f \frac{\mathbf{b} \cdot \hat{\mathbf{s}}}{c}} d\Omega \quad (2)$$

$$= \iint_{\text{sky}} \frac{A(\hat{\mathbf{s}}, f) I(\hat{\mathbf{s}}, f)}{\sqrt{1-l^2-m^2}} W_i(f) e^{-i2\pi f \frac{\mathbf{b} \cdot \hat{\mathbf{s}}}{c}} dl dm, \quad (3)$$

where, $I(\hat{\mathbf{s}}, f)$ and $A(\hat{\mathbf{s}}, f)$ are the sky brightness and antenna's directional power pattern, respectively, as a function of frequency (f) and direction on the sky denoted by the unit vector $\hat{\mathbf{s}} \equiv (l, m, n)$, $W_i(f)$ denotes instrumental bandpass weights, $W(f)$ is a spectral weighting function that controls the transfer function in the delay transform, $d\Omega = (1-l^2-m^2)^{-1/2} dl dm$ is the solid angle element to which $\hat{\mathbf{s}}$ is the unit normal vector, and c is the speed of light. $\tau = \mathbf{b} \cdot \hat{\mathbf{s}}/c$ is the geometric delay between antenna pairs measured relative to the zenith and provides a mapping to position on the sky.

In wide-field measurements, the steep rise in subtended solid angle near the horizon for a fixed delay bin size significantly enhances the integrated emission near the horizon delay limits. This is found to be true for diffuse emission even on wide antenna spacings because their foreshortening towards the horizon makes them sensitive to large angular scales that match the inverse of their foreshortened lengths.

Typically, an antenna is most sensitive towards its primary field of view relative to the rest of the sky, which in conjunction with the aforementioned wide-field effects results in a characteristic “pitchfork” signature in the delay spectrum.

Although there is marginal evidence for presence of this feature in the *zenith* snapshot presented in Thyagarajan et al. (2015), the high level of thermal noise prevented a robust confirmation. In this paper, we confirm the *pitchfork* signature using deeper data from the MWA.

3. THE MURCHISON WIDEFIELD ARRAY OBSERVATIONS

The MWA instrument configuration, EoR observations, and data analysis used in this study are already described in Thyagarajan et al. (2015). In order to reduce thermal fluctuations while maintaining coherence, it is essential to average independent data sets obtained over the same region of sky with identical beamformer settings. Hence, we select a subset of MWA snapshots each of duration 112 seconds obtained over different nights which are aligned to within 72 seconds of each other in *local sidereal time* (LST) around a mean LST of 0.04 hours with the MWA tile beam pointed at zenith. The database

contains 14 snapshots satisfying these criteria. Two of these snapshots were found to contain amplitude and phase artifacts for a significant duration across different baselines. Hence, they have been excluded from our analysis.

The delay spectra of the rest of the snapshots were verified to be coherent in their amplitudes and phases. These complex valued delay spectra from independent snapshots are averaged together to lower thermal fluctuations without losing coherence from foreground contributions. The results are discussed below.

4. RESULTS

Figure 1 shows the delay spectra obtained by averaging LST aligned delay spectra from individual snapshots from different nights. The dynamic range (in power spectrum units) in the averaged data is $\gtrsim 10$ higher compared to that in the *zenith* snapshot used in Thyagarajan et al. (2015), and is consistent with the improvement expected in averaging 12 independent snapshots. With this improvement in sensitivity, the foreground power near the horizon limits (white dotted lines) has become $\gtrsim 10$ times more prominent. We also note that faint horizontal features appear at $\tau = \pm 0.78 \mu\text{s}$ also as a result of lowering thermal fluctuations which were not seen earlier in the individual snapshots presented in Thyagarajan et al. (2015), thus confirming effective lowering of thermal fluctuations. We identify these faint features as the response in delay space of the MWA coarse band edges flagged periodically every 1.28 MHz.




FIG. 1.— Amplitudes of delay power spectra obtained by averaging 12 snapshots of LST aligned MWA data. The x -axis, denoted by $|\mathbf{b}|$ (and k_\perp), represents angular (and spatial) scales in the plane of the sky while the y -axis, shown in τ and k_\parallel , denotes the spatial scales along the line of sight. White dotted lines are the horizon delay limits. Power near the horizon limits caused by wide-field effects are prominent. Faint horizontal features at $\tau = \pm 0.78 \mu\text{s}$ are visible due to effective lowering of thermal fluctuations and are the response to periodic coarse band edge flagging of MWA data every 1.28 MHz.

Figure 2 shows the amplitudes of averaged delay spectra on three selected baseline vectors oriented northward. Data and noiseless simulations (using foreground and instrument models described in Thyagarajan et al. (2015)) are shown in black and red respectively. The horizontal dotted black line denotes *rms* of thermal fluctuations estimated from data. The vertical dashed line denotes horizon delay limits, and the vertical dot-dashed lines

denote delays at which the responses to coarse band edge flagging are expected.




FIG. 2.— Delay power spectra amplitudes on three antenna spacings oriented northward obtained by coherent averaging of 12 snapshots aligned in LST. The data and models are shown in black and red respectively. The antenna spacings are 83.5 m (*top*), 104.2 m (*middle*), and 171 m (*bottom*). The horizontal dotted line is the *rms* of thermal fluctuations. The vertical dashed lines denote the horizon delay limits. The vertical dot-dashed lines at $\tau = \pm 0.78 \mu\text{s}$ correspond to grating responses of periodic flagging of bandpass at intervals of 1.28 MHz. The dynamic range has increased by a factor ~ 10 as a result of coherent averaging relative to that in Thyagarajan et al. (2015). The peaks close to the horizon delay limits are distinctly visible at $\sim 10\text{--}1000 \sigma$ levels. Differences between model and data are primarily attributed to uncertainties in the foreground model and the MWA tile power pattern.

The delay power spectra morphologies from data and modeling are remarkably similar even while ignoring any differences in the amplitude scales. We attribute these differences to uncertainties in the foreground model, the MWA tile power pattern, thermal fluctuations, and other uncertainties noted in Thyagarajan et al. (2015).

We focus on the level of foreground power near the horizon limits in the data. Typically, the power near the negative horizon limit is seen with a signal-noise ratio (SNR) $\sim 10\text{--}100$, while that around the positive horizon limit is $\sim 100\text{--}1000$. The models of Thyagarajan et al. (2015) have noted that the foreground power near the horizon limits is due to the nature of wide-field measurements, and is predominantly composed of diffuse emission especially on baseline lengths $\lesssim 100$ m. Based on the morphological agreement between the data and the models, we conclude the features noted in our current analysis are a robust detection of the *pitchfork* signature predicted in Thyagarajan et al. (2015).

5. SUMMARY

This work was supported by the U. S. National Science Foundation (NSF) through award AST-1109257. DCJ is supported by an NSF Astronomy and Astrophysics Postdoctoral Fellowship under award AST-1401708. JCP is supported by an NSF Astronomy and Astrophysics Fellowship under award AST-1302774. This work makes use of the Murchison Radio-astronomy Observatory, operated by CSIRO. We acknowledge the Wajarri Yamatji people as the traditional owners of the Observatory site. Support for the MWA comes from the NSF (awards: AST-0457585, PHY-0835713, CAREER-0847753, and AST-0908884), the Australian Research Council (LIEF grants LE0775621 and LE0882938), the U.S. Air Force Office of Scientific Research (grant FA9550-0510247), and the Centre for All-sky Astrophysics (an Australian Research Council Centre of Excellence funded by grant CE110001020). Support is also provided by the Smithsonian Astrophysical Observatory, the MIT School of Science, the Raman Research Institute, the Australian National University, and the Victoria University of Wellington (via grant MED-E1799 from the New Zealand Ministry of Economic Development and an IBM Shared University Research Grant). The Australian Federal government provides additional support via the Commonwealth Scientific and Industrial Research Organisation (CSIRO), National Collaborative Research Infrastructure Strategy, Education Investment Fund, and the Australia India Strategic Research Fund, and Astronomy Australia Limited, under contract to Curtin University. We acknowledge the iVEC Petabyte Data Store, the Initiative in Innovative Computing and the CUDA Center for Excellence sponsored by NVIDIA at Harvard University, and the International Centre for Radio Astronomy Research (ICRAR), a Joint Venture of Curtin University and The University of Western Australia, funded by the Western Australian State government.

REFERENCES

- Parsons, A., Poher, J., McQuinn, M., Jacobs, D., & Aguirre, J. 2012a, *ApJ*, 753, 81
- Parsons, A. R., Poher, J. C., Aguirre, J. E., et al. 2012b, *ApJ*, 756, 165
- Thompson, A. R., Moran, J. M., & Swenson, Jr., G. W. 2001, *Interferometry and Synthesis in Radio Astronomy*, 2nd Edition (Wiley)
- Thyagarajan, N., Udaya Shankar, N., Subrahmanyam, R., et al. 2013, *ApJ*, 776, 6
- Thyagarajan, N., Jacobs, D. C., Bowman, J. D., et al. 2015, *ArXiv e-prints*, arXiv:1502.07596
- van Cittert, P. H. 1934, *Physica*, 1, 201
- Zernike, F. 1938, *Physica*, 5, 785