



IMMUNO-DIAGNOSTIC & PATHOLOGY LABORATORY

Halar Road Cross Lane, Besides L.I.C. Bldg.
Valsad-396 001. Ph.: (02632) 243280. Mo.: 99250 49280**KARIM SATTAR SHAIKH**
Male/60 YearsReg. Date : **21/09/2020**Lab. No **21556-18**Sample No
7620

Ref. Dr.

Dr. KRUNAL RAMTEKE MD
VIBRANT HOSPITAL VAP1

C - REACTIVE PROTEIN TITRE

Test	Result	Unit	Ref. Range
CRP :	49.00	mg/L	Adult < 6.0 mg/L Newborn upto 3 weeks < 4.1 mg/L Infants & Children < 2.8 mg/L
Method	BY IMMUNOTURBIDOMETRIC METHOD.		

LAB USE ONLY (Enter Date/Time:21/09/2020 10:00:00 AM) (Verify Date/Time:21/09/2020 11:26:00 AM)

Dr. Mehul SOLANKY
M.D.(Path & Bact)

KARIM SATTAR SHAIKH
Male/60 Years

Reg. Date : **21/09/2020**
Lab. No **21556-18**
Sample No
7620

Ref. Dr.

Dr. KRUNAL RAMTEKE MD
VIBRANT HOSPITAL VAP1

D - DIMER

Test	Result	Unit	Ref. Range
D - Dimer Level:	<u>1727.45</u>	ng/mL FEU	Up to 500 Fibrinogen Equivalent units

INTERPRETATION

Method - Fluorescent Immunoassay on Minividas/Ichroma 2

Concentrations of approximately ng/mL or more are considered pathological. Increased levels of D-Dimer in plasma indicate that excessive amounts of fibrin have formed in the vascular tree. Through Activated Fibrinolytic system, fibrin is degraded by plasmin to various soluble fragments in which the D-Dimer configuration occurs.

Elevated levels are seen in.

- Deep vein thrombosis
- pulmonary embolism
- Disseminated intravascular coagulation (DIC) disorder.
- Evaluation of acute myocardial infarction and unstable angina.
- Presence of D - Dimer in pre eclampsia suggest a severe disease.

LAB USE ONLY (Enter Date/Time:21/09/2020 12:32:00 PM) (Verify Date/Time:21/09/2020 12:32:00 PM)


Dr. Mehul SOLANKY
M.D.(Path & Bact)

KARIM SATTAR SHAIKH
Male/60 Years

Reg. Date : **21/09/2020**
Lab. No **21556-18**
Sample No
7620

Ref. Dr.
Dr. KRUNAL RAMTEKE MD
VIBRANT HOSPITAL VAP1

I N T E R L E U K I N - 6

<u>Test</u>	<u>Result</u>	<u>Unit</u>	<u>Normal Range</u>
Inter Leukin-6(IL-6):	<u>24.21</u>	pg/ml	0 to 7.0

Note & Interpretation

*InterLeukin-6 is a Pleiotropic helical glycosylated cytokine with wide range of functions.

*Many different cells are capable of IL-6 synthesis, including Monocytes, macrophage, fibroblasts, endothelial cells, keratinocytes, mast cells, T cells and many Tumor cells.

*IL-6 Concentrations in trauma patients may predict later complications from additional stress or indicate missed injuries or complications.

*Sequential measurements in serum or plasma of ICU Patients is useful in evaluating the severity of Systemic Inflammatory Response Syndrome (**SIRS**), **Sepsis** and **Septic shock**.

*IL-6 is also useful as an early alarm marker for detection of neonatal sepsis, acute phase response and regulation of metabolic processes.

Elevated IL-6 levels may occur in a variety of acute and chronic diseases associated with inflammation, such as sepsis, neoplastic disorder, autoimmune diseases, AIDS, Alcoholic Liver disease, and infections, or transplant rejection.

Serum or EDTA/HEPARIN Plasma samples can be used to test IL-6.

Labile parameter, stable for 5 hours at 2- - 25 degree C, if serum is used, use fresh sample, separate immediately after clotting and if not processed immediately store at 2-8 degree C.



IMMUNO-DIAGNOSTIC & PATHOLOGY LABORATORY

Halar Road Cross Lane, Besides L.I.C. Bldg.
Valsad-396 001. Ph.: (02632) 243280. Mo.: 99250 49280

KARIM SATTAR SHAIKH
Male/60 Years

Reg. Date : **21/09/2020**
Lab. No **21556-18**
Sample No
7620

Ref. Dr.
Dr. KRUNAL RAMTEKE MD
VIBRANT HOSPITAL VAP1

Dr. Mehul SOLANKY
M.D.(Path & Bact)