

Valsad-396 001. Ph.: (02632) 243280. Mo.: 99250 49280

KARIM SATTAR SHAIKH Male/60 Years

Reg. Date : Lab. No 18/09/2020 21290-18

Sample No

7251

Ref. Dr.

Dr. KRUNAL RAMTEKE MD VIBRANT HOSPITAL VAP1

C - REACTIVE PROTEIN TITRE

Test	Result	Unit	Ref. Range
CRP:	39.83	mg/L	Adult < 6.0 mg/L Newborn upto 3 weeks < 4.1 mg/L Infants & Children < 2.8 mg/L
Method	BY IMMUNOTURBIDOMETRIC METHOD.		

LAB USE ONLY (Enter Date/Time:18/09/2020 10:34:00 AM) (Verify Date/Time:18/09/2020 11:35:00 AM)

Dr. Mehul SOLANKY M.D.(Path & Bact)



Valsad-396 001. Ph.: (02632) 243280. Mo.: 99250 49280

KARIM SATTAR SHAIKH Male/60 Years

Reg. Date: 18/09/2020 Lab. No

21290-18

Sample No

7251

Ref. Dr.

Dr. KRUNAL RAMTEKE MD **VIBRANT HOSPITAL VAP1**

ANAEMIA PROFILE

Test Result Unit Ref. Range 324.3 S. Ferritin: ng/mL 30 - 300 ng/mL

< 10 ng/mL : Iron Deficiency Anaemia

10 - 20 ng/mL : Borderline >400 ng/mL :Iron Excess

METHOD: ECLIA ON ELECSYS 2010 - ROCHE

LAB USE ONLY (Enter Date/Time:18/09/2020 3:01:00 PM) (Verify Date/Time:18/09/2020 3:01:00 PM)

relus Dr. Mehul SOLANKY M.D.(Path & Bact)



Valsad-396 001. Ph.: (02632) 243280. Mo.: 99250 49280

KARIM SATTAR SHAIKH Male/60 Years

Reg. Date : Lab. No 18/09/2020 21290-18

Sample No

7251

Ref. Dr.

Dr. KRUNAL RAMTEKE MD VIBRANT HOSPITAL VAP1

SERUM LDH LEVEL

Test	Result	Unit	Ref. Range
S.L.D.H:	394.03	U/L	313 - 618 U/L

(Lactate Dehydrogenase)

LAB USE ONLY (Enter Date/Time:18/09/2020 3:02:00 PM) (Verify Date/Time:18/09/2020 3:02:00 PM)

Dr. Mehul SOLANKY M.D.(Path & Bact)



KARIM SATTAR SHAIKH Male/60 Years

Reg. Date : 18/09/2020 Lab. No 21290-18

Sample No

7251

Ref. Dr.

Dr. KRUNAL RAMTEKE MD VIBRANT HOSPITAL VAP1

INTERLEUKIN-6

TestResultUnitNormal RangeInter Leukin-6(IL-6):151.3pg/ml0 to 7.0

Note & Interpretation

- *InterLeukin-6 is a Pleiotropic helical glycosylated cytokine with wide range of functions.
- *Many different cells are capable of IL-6 synthesis, including Monocytes, macrophage, fibroblasts, endothelial cells, keratinocytes, mast cells, T cells and many Tumor cells.
- *IL-6 Concentrations is trauma patients may predict later complications from addistional stress or indicate missed injuries or complications.
- *Sequential measurements in serum or plasma of ICU Patients is useful in evaluating the severity of Systemic Inflammatory Response Syndrome (SIRS), Sepis and Septic shock.
- *ILI-6 is also useful as an early alarm marker for detection of neonatal sepsis. acute phase response and regulation of metabolic processes.

Elevated IL-6 levels may occur in a variety of acute and chronic diseases associated with inflammation, such as

sepsis, neoplastic disorder, autoimmune diseases, AIDS, Alcoholic Liver disease, and infections, or transplant rejection.

Serum or EDTA/HEPARIN Plasma samples can be used to test IL-6.

Labile parameter, stable for 5 hours at 2- - 25 degree C, if serum is used, use fresh sample, seperate immediately after clotting and if not processed immediately store at 2-8 degree C.

LAB USE ONLY (Enter Date/Time:18/09/2020 3:02:00 PM) (Verify Date/Time:18/09/2020 3:02:00 PM)



KARIM SATTAR SHAIKH Male/60 Years

Reg. Date : 18/09/2020 Lab. No 21290-18

Sample No

7251

Ref. Dr.

Dr. KRUNAL RAMTEKE MD VIBRANT HOSPITAL VAP1

> Dr. Mehul SOLANKY M.D.(Path & Bact)