

Individual Case Safety Report (ICSR)

Complete Pharmacovigilance Assessment – Simulated Case Study

1. Case Identification

Field	Case information
Case type	Initial ICSR – simulated educational case
Patient	42-year-old male
Reporter	Healthcare professional / hospital physician
Suspected medicine	Amoxicillin + Clavulanic Acid
Indication	Acute bacterial respiratory infection
Suspected ADR	Generalized urticaria, facial/lip swelling and difficulty in breathing
Case status	Serious ADR

2. Clinical History and ADR Identification

A 42-year-old male with no known drug allergy was prescribed amoxicillin/clavulanic acid for a suspected bacterial respiratory infection. Approximately two hours after the first dose, he developed generalized itching and urticaria followed by facial/lip swelling and difficulty in breathing. No other regular medicines or strong alternative causes were identified in this simulated history.

ADR parameter	Assessment
Onset	Approximately 2 hours after first dose
Clinical presentation	Urticaria, angioedema and dyspnea consistent with acute hypersensitivity
Clinical significance	Potentially life-threatening reaction
Drug action	Suspected medicine discontinued

3. Treatment and Outcome

The suspected medicine was immediately discontinued. The patient received appropriate emergency management, including intramuscular adrenaline/epinephrine, antihistamine and corticosteroid therapy with supportive monitoring. Symptoms improved. The patient was admitted for observation and later discharged in stable condition. Re-exposure was not performed.

4. Seriousness Assessment

Seriousness criterion	Finding
Death	No
Life-threatening	Yes – respiratory difficulty with acute hypersensitivity
Hospitalization	Yes – emergency treatment and observation
Persistent/significant disability	No
Congenital anomaly	Not applicable
Other medically important event	Yes – urgent emergency treatment required
Overall classification	SERIOUS

5. Expectedness Assessment

For this simulated case, acute hypersensitivity/anaphylactic reactions are treated as a known risk. In an actual ICSR, expectedness must be verified against the current approved product information or applicable reference safety information for the exact product, formulation and region.

6. Causality Assessment

Factor	Assessment
Temporal relationship	Strong – reaction occurred about 2 hours after exposure
Dechallenge	Positive – symptoms improved after withdrawal and treatment
Rechallenge	Not performed; ethically inappropriate after a serious hypersensitivity reaction
Alternative causes	No strong alternative cause identified in the simulated case
Clinical compatibility	Presentation is compatible with drug hypersensitivity
Overall causality	PROBABLE / HIGHLY PROBABLE

7. Severity Classification

Severity: Severe. The reaction required emergency treatment and hospitalization and included respiratory difficulty and facial swelling. Severity describes the intensity of the reaction; seriousness is determined by defined clinical/regulatory outcomes.

8. ICSR Data Summary

ICSR element	Recorded information
Patient	42-year-old male
Suspect drug	Amoxicillin + Clavulanic Acid
Indication	Acute bacterial respiratory infection
Reaction	Urticaria, angioedema and dyspnea / acute hypersensitivity
Time to onset	~2 hours after first dose
Seriousness	Serious – life-threatening potential and hospitalization
Severity	Severe
Expectedness	Known/expected type of reaction; verify current reference safety information
Causality	Probable / highly probable
Dechallenge	Positive
Rechallenge	Not performed
Outcome	Recovered/recovering

9. Drug Safety Associate Workflow

1. Confirm the minimum information needed for a valid ICSR: identifiable patient, identifiable reporter, suspected medicinal product and suspected adverse reaction.
2. Perform duplicate search/checking.
3. Review chronology, concomitant medicines, medical history, investigations and treatment.
4. Code the case using the current MedDRA version and applicable product terminology.
5. Assess seriousness, expectedness, causality and outcome.
6. Request follow-up information where important data are missing.
7. Perform medical review and quality control according to company procedures.
8. Submit/escalate within applicable regulatory and company timelines.

10. Pharmacovigilance Conclusion

This simulated ICSR represents a serious and severe suspected hypersensitivity reaction following amoxicillin/clavulanic acid exposure. The close temporal relationship, positive dechallenge, compatible clinical presentation and lack of a strong alternative explanation support a probable/highly probable causal association. Because the event was potentially life-threatening and required hospitalization, it warrants appropriate safety-reporting attention. Rechallenge should not be attempted after a serious hypersensitivity reaction.

Educational note: This is a simulated training case, not a real patient safety report. Actual MedDRA coding, expectedness and regulatory reporting decisions must follow the current applicable reference safety information, regulatory requirements and company SOPs.