

Applications of Derivatives; Maxima and Minima

Maximum and Minimum Values

In this section we see how to use partial derivatives to locate maxima and minima of functions of two variables.

Look at the hills and valleys in the graph of f shown in Figure 1.




Figure 1

Maximum and Minimum Values

There are two points (a, b) where f has a *local maximum*, that is, where $f(a, b)$ is larger than nearby values of $f(x, y)$.

The larger of these two values is the *absolute maximum*.

Likewise, f has two *local minima*, where $f(a, b)$ is smaller than nearby values.

The smaller of these two values is the *absolute minimum*.

Maximum and Minimum Values

1 Definition A function of two variables has a **local maximum** at (a, b) if $f(x, y) \leq f(a, b)$ when (x, y) is near (a, b) . [This means that $f(x, y) \leq f(a, b)$ for all points (x, y) in some disk with center (a, b) .] The number $f(a, b)$ is called a **local maximum value**. If $f(x, y) \geq f(a, b)$ when (x, y) is near (a, b) , then f has a **local minimum** at (a, b) and $f(a, b)$ is a **local minimum value**.

If the inequalities in Definition 1 hold for *all* points (x, y) in the domain of f , then f has an **absolute maximum** (or **absolute minimum**) at (a, b) .

2 Fermat's Theorem for Functions of Two Variables If f has a local maximum or minimum at (a, b) and the first-order partial derivatives of f exist there, then $f_x(a, b) = 0$ and $f_y(a, b) = 0$.

Maximum and Minimum Values

A point (a, b) is called a **critical point** (or *stationary point*) of f if $f_x(a, b) = 0$ and $f_y(a, b) = 0$, or if one of these partial derivatives does not exist.

Theorem 2 says that if f has a local maximum or minimum at (a, b) , then (a, b) is a critical point of f .

However, as in single-variable calculus, not all critical points give rise to maxima or minima.

At a critical point, a function could have a local maximum or a local minimum or neither.

Example 1

Let $f(x, y) = x^2 + y^2 - 2x - 6y + 14$.

Then

$$f_x(x, y) = 2x - 2 \qquad f_y(x, y) = 2y - 6$$

These partial derivatives are equal to 0 when $x = 1$ and $y = 3$, so the only critical point is $(1, 3)$.

By completing the square, we find that

$$f(x, y) = 4 + (x - 1)^2 + (y - 3)^2$$


Example 1

cont'd

Since $(x - 1)^2 \geq 0$ and $(y - 3)^2 \geq 0$, we have $f(x, y) \geq 4$ for all values of x and y .

Therefore $f(1, 3) = 4$ is a local minimum, and in fact it is the absolute minimum of f .

This can be confirmed geometrically from the graph of f , which is the elliptic paraboloid with vertex $(1, 3, 4)$ shown in Figure 2.



Maximum and Minimum Values

The following test, is analogous to the Second Derivative Test for functions of one variable.

3 Second Derivatives Test Suppose the second partial derivatives of f are continuous on a disk with center (a, b) , and suppose that $f_x(a, b) = 0$ and $f_y(a, b) = 0$ [that is, (a, b) is a critical point of f]. Let

$$D = D(a, b) = f_{xx}(a, b)f_{yy}(a, b) - [f_{xy}(a, b)]^2$$

- (a) If $D > 0$ and $f_{xx}(a, b) > 0$, then $f(a, b)$ is a local minimum.
- (b) If $D > 0$ and $f_{xx}(a, b) < 0$, then $f(a, b)$ is a local maximum.
- (c) If $D < 0$, then $f(a, b)$ is not a local maximum or minimum.

In case (c) the point (a, b) is called a **saddle point** of f and the graph of f crosses its tangent plane at (a, b) .

Absolute Maximum and Minimum Values

Absolute Maximum and Minimum Values

For a function f of one variable, the Extreme Value Theorem says that if f is continuous on a closed interval $[a, b]$, then f has an absolute minimum value and an absolute maximum value.

According to the Closed Interval Method, we found these by evaluating f not only at the critical numbers but also at the endpoints a and b .

There is a similar situation for functions of two variables.

Absolute Maximum and Minimum Values

Just as a closed interval contains its endpoints, a **closed set** in $\mathbb{R}^2$ is one that contains all its boundary points.

[A boundary point of D is a point (a, b) such that every disk with center (a, b) contains points in D and also points not in D .]


For instance, the disk

$$D = \{(x, y) \mid x^2 + y^2 \leq 1\}$$


which consists of all points on and inside the circle $x^2 + y^2 = 1$, is a closed set because it contains all of its boundary points (which are the points on the circle $x^2 + y^2 = 1$).

Absolute Maximum and Minimum Values

But if even one point on the boundary curve were omitted, the set would not be closed. (See Figure 11.)



(a) Closed sets



(b) Sets that are not closed

Figure 11

Absolute Maximum and Minimum Values

A **bounded set** in $\mathbb{R}^2$ is one that is contained within some disk.

In other words, it is finite in extent.

Then, in terms of closed and bounded sets, we can state the following counterpart of the Extreme Value Theorem in two dimensions.

8 Extreme Value Theorem for Functions of Two Variables If f is continuous on a closed, bounded set D in $\mathbb{R}^2$, then f attains an absolute maximum value $f(x_1, y_1)$ and an absolute minimum value $f(x_2, y_2)$ at some points (x_1, y_1) and (x_2, y_2) in D .

Absolute Maximum and Minimum Values

To find the extreme values guaranteed by Theorem 8, we note that, by Theorem 2, if f has an extreme value at (x_1, y_1) , then (x_1, y_1) is either a critical point of f or a boundary point of D .

Thus we have the following extension of the Closed Interval Method.

9 To find the absolute maximum and minimum values of a continuous function f on a closed, bounded set D :

1. Find the values of f at the critical points of f in D .
2. Find the extreme values of f on the boundary of D .
3. The largest of the values from steps 1 and 2 is the absolute maximum value; the smallest of these values is the absolute minimum value.

Example 7

Find the absolute maximum and minimum values of the function $f(x, y) = x^2 - 2xy + 2y$ on the rectangle

$$D = \{(x, y) | 0 \leq x \leq 3, 0 \leq y \leq 2\}.$$

Solution:

Since f is a polynomial, it is continuous on the closed, bounded rectangle D , so Theorem 8 tells us there is both an absolute maximum and an absolute minimum.

According to step 1 in [\[9\]](#), we first find the critical points.

Example 7 – *Solution*

These occur when

$$f_x = 2x - 2y = 0$$

$$f_y = -2x + 2 = 0$$

so the only critical point is $(1, 1)$, and the value of f there is $f(1, 1) = 1$.

In step 2 we look at the values of f on the boundary of D , which consists of the four line segments L_1, L_2, L_3, L_4 shown in Figure 12.




Figure 12

Example 7 – *Solution*

cont'd

On L_1 we have $y = 0$ and

$$f(x, 0) = x^2 \qquad 0 \leq x \leq 3$$

This is an increasing function of x , so its minimum value is $f(0, 0) = 0$ and its maximum value is $f(3, 0) = 9$.

On L_2 we have $x = 3$ and

$$f(3, y) = 9 - 4y \qquad 0 \leq y \leq 2$$

This is a decreasing function of y , so its maximum value is $f(3, 0) = 9$ and its minimum value is $f(3, 2) = 1$.

Example 7 – *Solution*

On L_3 we have $y = 2$ and

$$f(x, 2) = x^2 - 4x + 4 \quad 0 \leq x \leq 3$$

Simply by observing that $f(x, 2) = (x - 2)^2$, we see that the minimum value of this function is $f(2, 2) = 0$ and the maximum value is $f(0, 2) = 4$.

Example 7 – *Solution*

cont'd

Finally, on L_4 we have $x = 0$ and

$$f(0, y) = 2y \quad 0 \leq y \leq 2$$

with maximum value $f(0, 2) = 4$ and minimum value $f(0, 0) = 0$.

Thus, on the boundary, the minimum value of f is 0 and the maximum is 9.

Example 7 – *Solution*

In step 3 we compare these values with the value $f(1, 1) = 1$ at the critical point and conclude that the absolute maximum value of f on D is $f(3, 0) = 9$ and the absolute minimum value is $f(0, 0) = f(2, 2) = 0$.

Figure 13 shows the graph of f .




Figure 13

$$f(x, y) = x^2 - 2xy + 2y$$