

CORRECTION NOTICE

Propranolol for Anxiety · 10mg and 40mg tablets

The signed PGD behind this page is unaltered. These corrections take precedence over it.

Two corrections apply, to BOTH the 10mg and the 40mg arms.

1. Quantity is strength-specific

Both arms print the identical line ·Up to 56 tablets (28-day supply at twice daily dosing)·. That is only true at 10mg twice daily. Fifty-six 40mg tablets is 2,240mg, four times the quantity the 10mg arm authorises, and nothing in either arm says when 40mg is indicated rather than 10mg.

Until the corrected version is signed, supply the 10mg strength and no more than one course for a single situational event. Do not supply 56 tablets of the 40mg strength.

2. Screening before supply

The document's own guidance names suicidal ideation, severe depression, PTSD and substance misuse as red flags requiring referral, but none of them appears in the exclusion criteria. Propranolol is cardiotoxic in overdose. Before supplying, ask about and REFER rather than supply where there is:

- **Any suicidal ideation, self-harm or severe depression**
- **PTSD, severe panic disorder or agoraphobia**
- **Comorbid substance or alcohol misuse**

Also check current medicines and do not supply to a patient already taking another beta-blocker, or taking oral verapamil or diltiazem.

Issued 7 September 2026 · Get Real Health. Corrected version to follow after clinical review.

This Patient Group Direction (PGD) is intended only for registered healthcare professionals who have been named and authorised by their organisation to practice under it.

Patient Group Direction

for the administration of Propranolol for the treatment of Anxiety (Situational and Somatic Symptoms)

Summary of NICE / NICE CKS Guidelines for Anxiety (Situational and Somatic Symptoms)

Overview

- Anxiety disorders are common, affecting physical and psychological wellbeing
- Physical symptoms include palpitations, tremor, sweating, and gastrointestinal disturbance

Assessment

- Use GAD-7 score to assess severity and impact
- Differentiate generalised anxiety from panic disorder, social anxiety, and PTSD
- Exclude thyroid disease and cardiac causes for physical symptoms

Management

- Cognitive Behavioural Therapy (CBT) is first-line psychological treatment
- Propranolol for symptomatic relief of physical symptoms in situational/performance anxiety
- Propranolol is not first-line for generalised anxiety disorder

Red flags requiring specialist referral

- Suicidal ideation or severe depression
- Severe panic disorder or agoraphobia
- PTSD or trauma-related anxiety
- Comorbid substance misuse

Patient Group Direction

for the supply/administration of

Propranolol 10mg tablets

for the treatment of

Anxiety (Situational and Somatic Symptoms)

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	• Pharmacist registered and practicing with the GPhC • Pharmacy technician registered and practicing with the GPhC

Training and assessment	Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. • All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines • All users must previously have used PGDs to supply medication • Must work in compliance with the SOPs of their own employer • All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	• All users must be up to date with CPD requirements and appraisal • All users must be up to date with MHRA and safety alerts with regards to medical products • All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	Symptomatic relief of situational/performance anxiety (physical symptoms) in adults
Inclusion criteria	Adults aged 18 years and over Physical symptoms of situational anxiety (palpitations, tremor, sweating) Informed consent
Exclusion criteria	Asthma or history of bronchospasm Uncontrolled heart failure Cardiogenic shock Severe bradycardia (heart rate <50 bpm at rest) Hypotension (systolic BP <90 mmHg) Sick sinus syndrome Second-degree or third-degree heart block Phaeochromocytoma (unless already on alpha-blocker) Metabolic acidosis Prinzmetal's angina Concurrent intravenous verapamil or diltiazem Known hypersensitivity to propranolol
Cautions	Diabetes (masks hypoglycaemia symptoms) First-degree heart block Portal hypertension

	Peripheral vascular disease Psoriasis Raynaud's Myasthenia gravis History of anaphylaxis Depression Hepatic/renal impairment Elderly Do not stop abruptly
Actions if patient is excluded or declines treatment	Advise on alternative treatment options and how these can be accessed. Document any advice given and the decision reached. Inform or refer to the GP as appropriate.

Details of the medicine

Name, form and strength of medicine	Propranolol 10mg tablets
Legal category	POM
Storage	Below 25°C, protect from light
Route/method of administration	Oral
Dose and frequency	10-40mg taken 30-60 minutes before anxiety-provoking situation Alternatively, 10-40mg two to three times daily for ongoing situational anxiety Maximum 120mg daily
Quantity to be administered and/or supplied	Up to 56 tablets (28-day supply at twice daily dosing)
Maximum or minimum treatment period	Review at 4 weeks; ongoing use should be reviewed regularly. Consider gradual dose reduction if discontinuing.
Adverse effects	Common: fatigue, cold extremities, bradycardia, GI disturbance, sleep disturbance, dizziness Uncommon: bronchospasm, hypotension, heart failure exacerbation Rare: alopecia, visual disturbance, psoriasiform rash

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	Record the following information:

	• that valid informed consent was given • name of individual, address, date of birth and GP • name of healthcare practitioner • name and brand of medication • date of supply dose, form and route quantity supplied • advice given, including advice given if excluded or declines treatment • details of any adverse drug reactions and actions taken supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: • Adults aged 18 years and over - keep records for 8 years • Children aged under 18 years - keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)
--	--

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medication.
Follow-up advice to be given to patient or carer	Do not stop suddenly, report breathlessness/wheeze, may cause cold hands/feet, not a cure for anxiety - consider psychological therapy, avoid alcohol (additive CNS depression)

Key references

• NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 https://www.nice.org.uk/guidance/mg2 • NICE MPG2 Patient group directions: competency framework for health professionals using patient group directions. Updated March 2017 https://www.nice.org.uk/guidance/mpg2/resources • NICE CKS Guidance: Anxiety disorders • British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines
--

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which			
--	--	--	--

this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature
Valid from date	Expiry date		
1/11/25	31/7/27		

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date
Name of professional group signatory	Job title & name of organisation	Signature	Date

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor		2/11/25

	GMC: 6047293		
Chris Pilkington	Pharmacist GPhC: 2046322		2/11/25

Change history

Version number	Change details	Date
001	Development & issue of new PGD	1st November 2025

Patient Group Direction
for the supply/administration of
Propranolol 40mg tablets
for the treatment of
Anxiety (Situational and Somatic Symptoms)

Healthcare professionals covered and training

	Healthcare professionals allowed to work under this PGD and requirements
Qualifications and Registration	• Pharmacist registered and practicing with the GPhC • Pharmacy technician registered and practicing with the GPhC
Training and assessment	Pharmacists and Pharmacy Technicians must have completed training relevant to this condition to use this PGD. This training will be documented and overseen by the Get Real Health team. • All users must be familiar with the SPC for the products as well as BNF advice and any applicable national guidelines • All users must previously have used PGDs to supply medication • Must work in compliance with the SOPs of their own employer • All users must have appropriate indemnity in place to cover this service
Further training and responsibilities	• All users must be up to date with CPD requirements and appraisal • All users must be up to date with MHRA and safety alerts with regards to medical products • All users must understand the concepts of capacity and consent and be aware of the Mental Capacity Act 2005.

PGD Indication	Symptomatic relief of situational/performance anxiety (physical symptoms) in adults
Inclusion criteria	Adults aged 18 years and over Physical symptoms of situational anxiety (palpitations, tremor, sweating) Informed consent
Exclusion criteria	Asthma or history of bronchospasm Uncontrolled heart failure Cardiogenic shock Severe bradycardia (heart rate <50 bpm at rest)

	Hypotension (systolic BP <90 mmHg) Sick sinus syndrome Second-degree or third-degree heart block Phaeochromocytoma (unless already on alpha-blocker) Metabolic acidosis Prinzmetal's angina Concurrent intravenous verapamil or diltiazem Known hypersensitivity to propranolol
Cautions	Diabetes (masks hypoglycaemia symptoms) First-degree heart block Portal hypertension Peripheral vascular disease Psoriasis Raynaud's Myasthenia gravis History of anaphylaxis Depression Hepatic/renal impairment Elderly Do not stop abruptly
Actions if patient is excluded or declines treatment	Advise on alternative treatment options and how these can be accessed. Document any advice given and the decision reached. Inform or refer to the GP as appropriate.

Details of the medicine

Name, form and strength of medicine	Propranolol 40mg tablets
Legal category	POM
Storage	Below 25°C, protect from light
Route/method of administration	Oral
Dose and frequency	10-40mg taken 30-60 minutes before anxiety-provoking situation Alternatively, 10-40mg two to three times daily for ongoing situational anxiety Maximum 120mg daily
Quantity to be administered	Up to 56 tablets (28-day supply at twice daily dosing)

and/or supplied	
Maximum or minimum treatment period	Review at 4 weeks; ongoing use should be reviewed regularly. Consider gradual dose reduction if discontinuing.
Adverse effects	Common: fatigue, cold extremities, bradycardia, GI disturbance, sleep disturbance, dizziness Uncommon: bronchospasm, hypotension, heart failure exacerbation Rare: alopecia, visual disturbance, psoriasiform rash

Yellow card reporting	Report suspected adverse effects via https://yellowcard.mhra.gov.uk and inform the GP as appropriate.
Records to be kept	Record the following information: • that valid informed consent was given • name of individual, address, date of birth and GP • name of healthcare practitioner • name and brand of medication • date of supply dose, form and route quantity supplied • advice given, including advice given if excluded or declines treatment • details of any adverse drug reactions and actions taken supplied via PGD Records should be signed and dated. All records should be clear, legible and contemporaneous. Electronic records can be made. A record of all individuals receiving treatment under this PGD should also be kept for audit purposes as follows: • Adults aged 18 years and over - keep records for 8 years • Children aged under 18 years - keep records until the 25th birthday (if the patient was 17 years when treatment was finished, keep records until the 26th birthday)

Patient information

Written information to be given to patient or carer	Supply the patient information leaflet (PIL) provided with the medication.
Follow-up advice to be given to patient or carer	Do not stop suddenly, report breathlessness/wheeze, may cause cold hands/feet, not a cure for anxiety - consider psychological therapy, avoid alcohol (additive CNS depression)

Key references

- NICE Medicines Practice Guideline 2 (MPG2): Patient Group Directions. Published March 2017 <https://www.nice.org.uk/guidance/mg2>
- NICE MPG2 Patient group directions: competency framework for health professionals using

patient group directions. Updated March 2017
<https://www.nice.org.uk/guidance/mpg2/resources>

- NICE CKS Guidance: Anxiety disorders
- British National Formulary (BNF) Key information on the selection, prescribing, dispensing and administration of medicines

Agreement to practise by the Registered Healthcare Professional

By signing this PGD you are confirming that you agree to its contents and that you will work within it. PGDs do not remove inherent professional obligations or accountability. It is the responsibility of each healthcare professional to practise only within the bounds of their own competence and professional code of conduct.

Name and address of pharmacy premises / clinic premises to which this PGD relates			
Name of healthcare professional	Job title	Registration number	Signature
Valid from date	Expiry date		
1/11/25	31/7/27		

PGD adoption and authorisation by the employer/clinical lead

The employer has ensured that the person using this PGD, has the knowledge and skills to safely provide the service which will encompass the supply or administration of a medication or medicine.

This section must be signed by the superintendent or clinical lead responsible for this service.

Name of superintendent / clinical lead	Job title & name of organisation	Signature	Date

	(GPhC: 2046322)		
Name of professional group signatory	Job title & name of organisation	Signature	Date
	(GPhC: 2046322)		

Signed on behalf of Get Real Health

Name	Qualification	Signature	Date
Nitin Shori	Medical Doctor GMC: 6047293		2/11/25
Chris Pilkington	Pharmacist GPhC: 2046322		2/11/25

Change history

Version number	Change details	Date
001	Development & issue of new PGD	1st November 2025